

CPRD Bibliography December 2017

- Zhong VW, Juhaeri J, Cole SR, Kontopantelis E, Shay CM, Gordon-Larsen P, et al. Incidence and Trends in Hypoglycemia Hospitalization in Adults With Type 1 and Type 2 Diabetes in England, 1998-2013: A Retrospective Cohort Study. *Diabetes care*. 2017.
- Zghebi SS, Steinke DT, Rutter MK, Ashcroft DM. A longitudinal observational study on the patterns of comorbid clinical conditions in patients with and patients without Type 2 diabetes. 2017.
- Zghebi SS, Steinke DT, Carr MJ, Rutter MK, Emsley RA, Ashcroft DM. Examining Trends in Type 2 Diabetes Incidence, Prevalence and Mortality in the UK between 2004 and 2014. *Diabetes Obes Metab*. 2017.
- Yu O, Azoulay L, Yin H, Filion KB, Suissa S. Sulfonylureas as Initial Treatment for Type 2 Diabetes and the Risk of Severe Hypoglycemia. *Am J Med*. 2017.
- Yu D, Jordan KP, Bedson J, Englund M, Blyth F, Turkiewicz A, et al. Population trends in the incidence and initial management of osteoarthritis: age-period-cohort analysis of the Clinical Practice Research Datalink, 1992-2013. *Rheumatology (Oxford)*. 2017;56(11):1902-17.
- Young GJ, Harrison S, Turner EL, Walsh EI, Oliver SE, Ben-Shlomo Y, et al. Prostate-specific antigen (PSA) testing of men in UK general practice: a 10-year longitudinal cohort study. *BMJ Open*. 2017;7(10):e017729.
- Yates TA, Tomlinson LA, Bhaskaran K, Langan S, Thomas S, Smeeth L, et al. Lansoprazole use and tuberculosis incidence in the United Kingdom Clinical Practice Research Datalink: A population based cohort. *PLoS Med*. 2017;14(11):e1002457.
- Xiao X, Fischbach S, Zhang T, Chen C, Sheng Q, Zimmerman R, et al. SMAD3/stat3 signaling mediates beta-cell epithelial-mesenchymal transition in chronic pancreatitis-related diabetes. *Diabetes*. 2017;66(10):2646-58.
- Wu JW, Azoulay L, Majdan A, Boivin J-F, Pollak M, Suissa S. Long-Term Use of Long-Acting Insulin Analogs and Breast Cancer Incidence in Women With Type 2 Diabetes. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2017;JCO2017734491.
- Wu JW, Azoulay L, Boivin J-F, Suissa S, Pollak M. Long-term use of long-acting insulin analogs and breast cancer incidence in women with type 2 diabetes. *Journal of Clinical Oncology*. 2017;35(32):3647-53.
- Wong J, Donegan K, Seabroke S, Bryan P. Evaluation of compliance to the timing of childhood vaccinations following introduction of the meningococcal B vaccine in the UK. *Drug Safety*. 2017;40(10):1000-1.
- Wing K, Bhaskaran K, Pealing L, Root A, Smeeth L, van Staa TP, et al. Quantification of the risk of liver injury associated with flucloxacillin: a UK population-based cohort study. *J Antimicrob Chemother*. 2017;72(9):2636-46.
- Wilson JC, Sarsour K, Collinson N, Tuckwell K, Musselman D, Klearman M, et al. Serious adverse effects associated with glucocorticoid therapy in patients with giant cell arteritis (GCA): A nested case-control analysis. *Semin Arthritis Rheum*. 2017;46(6):819-27.
- Wilson JC, Sarsour K, Collinson N, Tuckwell K, Musselman D, Klearman M, et al. Incidence of outcomes potentially associated with corticosteroid therapy in patients with giant cell arteritis. *Seminars in arthritis and rheumatism*. 2017;46(5):650-6.

- Wilson CL, Rhodes KM, Payne RA. Financial incentives improve recognition but not treatment of cardiovascular risk factors in severe mental illness. *PloS one*. 2017;12(6):e0179392.
- Williams R, de Vries F, Kothny W, Serban C, Lopez-Leon S, Chu C, et al. Cardiovascular safety of vildagliptin in patients with type 2 diabetes: a European multi-database, non-interventional post-authorization safety study. *Diabetes, Obesity and Metabolism*. 2017.
- Wilding JPH, Godec T, Khunti K, Pocock S, Fox R, Smeeth L, et al. Changes in HbA1c and weight after initiation of second-line therapy in patients with type 2 diabetes in clinical practice. *Diabetologia*. 2017;60(1):S301.
- Wilding J, Bailey C, Rigney U, Blak B, Kok M, Emmas C. Dapagliflozin therapy for type 2 diabetes in primary care: Changes in HbA1c, weight and blood pressure over 2 years follow-up. *Primary care diabetes*. 2017.
- Wilding J, Bailey C, Rigney U, Blak B, Kok M, Emmas C. Dapagliflozin therapy for Type 2 diabetes in primary care: Effects on HbA1c, weight and blood pressure over two years follow-up. 2017.
- West J, Card TR, Aithal GP, Fleming KM. Risk of hepatocellular carcinoma among individuals with different aetiologies of cirrhosis: a population-based cohort study. *Aliment Pharmacol Ther*. 2017.
- Weir S, Samnaliev M, Kuo T-C, Ni Choitir C, Tierney TS, Cumming D, et al. The incidence and healthcare costs of persistent postoperative pain following lumbar spine surgery in the UK: a cohort study using the Clinical Practice Research Datalink (CPRD) and Hospital Episode Statistics (HES). *BMJ Open*. 2017;7(9):e017585.
- Weir S, Samnaliev M, Kuo T-C, Choitir CN, Tierney TS, Taylor RS, et al. Does instrumented spinal surgery have an increased rate of persistent post-operative pain (PPP)? *The Spine Journal*. 2017;17(3):S11.
- Weir S, Samnaliev M, Kuo T-C, Choitir CN, Tierney TS, Taylor RS, et al. The rate of persistent post-operative pain (FBSS) in revision lumbar spinal surgery. *The Spine Journal*. 2017;17(3):S10.
- Weinstein RB, Ryan P, Berlin JA, Matcho A, Schuemie M, Swerdel J, et al. Channeling in the Use of Nonprescription Paracetamol and Ibuprofen in an Electronic Medical Records Database: Evidence and Implications. *Drug safety*. 2017.
- Watson J, Nicholson BD, Hamilton W, Price S. Identifying clinical features in primary care electronic health record studies: methods for codelist development. *BMJ Open*. 2017;7(11):e019637.
- Wang Y, Pfeiffer RM, Alsaggaf R, Meeraus W, Gage JC, Anderson LA, et al. Risk of skin cancer among patients with myotonic dystrophy type 1 based on primary care physician data from the U.K. *Clinical Practice Research Datalink*. *Int J Cancer*. 2017.
- Walker S, Hamilton W. Risk of cervical cancer in symptomatic women aged ≥ 40 in primary care: A case-control study using electronic records. *European journal of cancer care*. 2017;26(3).
- Walker JL, Andrews NJ, Mathur R, Smeeth L, Thomas SL. Trends in the burden of varicella in UK general practice. *Epidemiol Infect*. 2017:1-5.
- Vuik SI, Fontana G, Mayer E, Darzi A. Do hospitalisations for ambulatory care sensitive conditions reflect low access to primary care? An observational cohort study of primary care usage prior to hospitalisation. *BMJ Open*. 2017;7(8):e015704.
- Viswanathan A, Spera C, Mullins A, Covert D, Banhazi J, McDwyer P, et al. Resource Utilization Among Glaucoma Patients in the UK Treated with Beta-Blocker and Non-Beta-Blocker Adjunctive Therapy: A

Retrospective Cohort Analysis. *Adv Ther.* 2017.

- Verstraeten T, Cattaert T, Harris J, Lopman B, Tam CC, Ferreira G. Estimating the Burden of Medically Attended Norovirus Gastroenteritis: Modeling Linked Primary Care and Hospitalization Datasets. *J Infect Dis.* 2017;216(8):957-65.
- Verstraeten T, Cattaert T, Harris J, Lopman B, Tam CC, Ferreira G. Estimating the burden of medically-attended norovirus gastroenteritis: modelling linked primary care and hospitalization datasets. *J Infect Dis.* 2017.
- Verstraeten T, Cattaert T, Ferreira G. Gender inequality in acute gastro-enteritis rates in England. *Value in Health.* 2017;20(9):A781.
- Venkatesan S, Myles PR, Manning HJ, Mozid AM, Andersson C, Jørgensen ME, et al. Cohort study of preoperative blood pressure and risk of 30-day mortality after elective non-cardiac surgery. *British journal of anaesthesia.* 2017.
- van Dalem J, Brouwers MC, Stehouwer CDA, Krings A, Klungel OH, Driessen JH, et al. Risk of a first ever acute myocardial infarction and all-cause mortality with sulphonylurea treatment: a population-based cohort study. *Diabetes Obes Metab.* 2017.
- Van Boven J, Benhaddi H, Roche N, Bjermer L, Miravittles M, Voorham J, et al. Real-world cost effectiveness comparison of symbicort turbuhaler and duoresp spiromax in UK patients with asthma or chronic obstructive pulmonary disease. *Value in Health.* 2017;20(9):A645.
- Toth PP, Danese M, Villa G, Qian Y, Beaubrun A, Lira A, et al. Estimated burden of cardiovascular disease and value-based price range for evolocumab in a high-risk, secondary-prevention population in the US payer context. *J Med Econ.* 2017:1-10.
- Tomkins S, Chapman C, Myland M, Tham R, de Nobrega R, Jackson B, et al. Treating iron deficiency in patients with gastrointestinal disease: Risk of re-attendance in secondary care. *PloS one.* 2017;12(12):e0189952.
- Tillett W, Charlton R, Nightingale A, Snowball J, Green A, Smith C, et al. Interval between onset of psoriasis and psoriatic arthritis comparing the UK Clinical Practice Research Datalink with a hospital-based cohort. *Rheumatology (Oxford).* 2017.
- Thompson A, Wright AK, Ashcroft DM, van Staa TP, Pirmohamed M. Epidemiology of alcohol dependence in UK primary care: Results from a large observational study using the Clinical Practice Research Datalink. *PLOS ONE.* 2017;12(3):e0174818.
- Thompson A, Ashcroft DM, Owens L, van Staa TP, Pirmohamed M. Drug therapy for alcohol dependence in primary care in the UK: A Clinical Practice Research Datalink study. *PLOS ONE.* 2017;12(3):e0173272.
- Taylor GMJ, Taylor AE, Thomas KH, Jones T, Martin RM, Munafò MR, et al. The effectiveness of varenicline versus nicotine replacement therapy on long-term smoking cessation in primary care: a prospective cohort study of electronic medical records. *Int J Epidemiol.* 2017;46(6):1948-57.
- Tate AR, Dungey S, Glew S, Beloff N, Williams R, Williams T. Quality of recording of diabetes in the UK: how does the GP's method of coding clinical data affect incidence estimates? Cross-sectional study using the CPRD database. *BMJ Open.* 2017;7(1):e012905.
- Taqi A, Otete H, Knaggs RD. Trends in opioid prescribing in primary care patients with knee osteoarthritis: A

population based observational study. *Value in Health*. 2017;20(9):A540-A1.

- Tammes P, Purdy S, Salisbury C, MacKichan F, Lasserson D, Morris RW. Continuity of Primary Care and Emergency Hospital Admissions Among Older Patients in England. *Annals of family medicine*. 2017;15(6):515-22.
- Tamimi I, Nicolau B, Eimar H, Arekunnath Madathil S, Kezouh A, Karp I, et al. Acetylcholinesterase inhibitors and the risk of osteoporotic fractures: nested case-control study. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2017.
- Suruki RY, Daugherty JB, Boudiaf N, Albers FC. The frequency of asthma exacerbations and healthcare utilization in patients with asthma from the UK and USA. *BMC Pulm Med*. 2017;17(1):74.
- Suissa S, Dell'Aniello S, Ernst P. Concurrent use of long-acting bronchodilators in COPD and the risk of adverse cardiovascular events. *Eur Respir J*. 2017;49(5).
- Stolz D, Kostikas K, Loeffroth E, Fogel R, Clemens A, Gutzwiller FS, et al. Gender differences in COPD exacerbations: Analysis from the CPRD database. *Thorax*. 2017;72:A186.
- Stolk LM, de Vries F, Ebbelaar C, de Boer A, Schalekamp T, Souverein P, et al. Risk of myocardial infarction in patients with atrial fibrillation using vitamin K antagonists, aspirin or direct acting oral anticoagulants: Myocardial infarction with VKAs, aspirin or DOACS. *British Journal of Clinical Pharmacology*. 2017.
- Stocks SJ, Kontopantelis E, Webb RT, Avery AJ, Burns A, Ashcroft DM. Antipsychotic Prescribing to Patients Diagnosed with Dementia Without a Diagnosis of Psychosis in the Context of National Guidance and Drug Safety Warnings: Longitudinal Study in UK General Practice. *Drug safety*. 2017.
- Stewart D, Han L, Doran T, McCambridge J. Alcohol consumption and all-cause mortality: an analysis of general practice database records for patients with long-term conditions. *J Epidemiol Community Health*. 2017;71(8):729-35.
- Stevens S, Bankhead C, Mukhtar T, Perera-Salazar R, Holt TA, Salisbury C, et al. Patient-level and practice-level factors associated with consultation duration: a cross-sectional analysis of over one million consultations in English primary care. *BMJ Open*. 2017;7(11):e018261.
- Stapley SA, Rubin GP, Alsina D, Shephard EA, Rutter MD, Hamilton WT. Clinical features of bowel disease in patients aged <50 years in primary care: a large case-control study. *British Journal of General Practice*. 2017;bjgp17X690425.
- Springate DA, Parisi R, Olier I, Reeves D, Kontopantelis E. rEHR: An R package for manipulating and analysing Electronic Health Record data. *PLOS ONE*. 2017;12(2):e0171784.
- Spaetgens B, de Vries F, Driessen JHM, Leufkens HG, Souverein PC, Boonen A, et al. Risk of infections in patients with gout: a population-based cohort study. *Sci Rep*. 2017;7(1):1429.
- Skrtic S, Cabrera C, Olsson M, Schneck V, Lind M. Contemporary risk estimates of three HbA1c variables in relation to heart failure following diagnosis of type 2 diabetes. *Heart*. 2017;103(5):353-8.
- Sinyavskaya L, Gauthier S, Renoux C, Dell'Aniello S, Suissa S, Brassard P. Comparative effect of statins on the risk of incident Alzheimer disease. *Neurology*. 2017.
- Sinnott S-J, Smeeth L, Williamson E, Douglas IJ. Trends for prevalence and incidence of resistant

hypertension: population based cohort study in the UK 1995-2015. *BMJ*. 2017;358:j3984.

- Shields BM, Dennis J, Henley W, Weedon M, Lonergan M, Rodgers L, et al. Predictors of response to SGLT2-inhibitors and DPP4-inhibitors. *Diabetologia*. 2017;60(1):S404.
- Sheppard T, Tamblyn R, Abrahamowicz M, Lunt M, Sperrin M, Dixon WG. A comparison of methods for estimating the temporal change in a continuous variable: Example of HbA1c in patients with diabetes. *Pharmacoepidemiol Drug Saf*. 2017.
- Shepherd C, Koepp M, Myland M, Patel K, Miglio C, Siva V, et al. Understanding the health economic burden of patients with tuberous sclerosis complex (TSC) with epilepsy: a retrospective cohort study in the UK Clinical Practice Research Datalink (CPRD). *BMJ Open*. 2017;7(10):e015236.
- Shah A, Judge A, Delmestri A, Edwards K, Arden NK, Prieto-Alhambra D, et al. Incidence of shoulder dislocations in the UK, 1995-2015: a population-based cohort study. *BMJ Open*. 2017;7(11):e016112.
- Seliger C, Meier CR, Becker C, Jick SS, Proescholdt M, Bogdahn U, et al. Diabetes, use of metformin, and the risk of meningioma. *PloS one*. 2017;12(7):e0181089.
- Seliger C, Meier CR, Becker C, Jick SS, Proescholdt M, Bogdahn U, et al. Metabolic syndrome in relation to risk of meningioma. *Oncotarget*. 2017;8(2):2284-92.
- Seabroke S, Bryan P. Safety of meningococcal group B vaccine: Experience from the UK. *Drug Safety*. 2017;40(10):1003.
- Schneider C, Bodmer M, Jick SS, Meier CR. Colorectal cancer and markers of anemia. *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation (ECP)*. 2017.
- Schmidt SAJ, Langan SM, Pedersen HS, Schønheyder HC, Thomas SL, Smeeth L, et al. Mood Disorders and Risk of Herpes Zoster in Two Population-Based Case-Control Studies in Denmark and the United Kingdom. *Am J Epidemiol*. 2017.
- Schmidt M, Mansfield KE, Bhaskaran K, Nitsch D, Sørensen HT, Smeeth L, et al. Serum creatinine elevation after renin-angiotensin system blockade and long term cardiorenal risks: cohort study. *BMJ*. 2017;j791.
- Schmidt M, Mansfield KE, Bhaskaran K, Nitsch D, Sørensen HT, Smeeth L, et al. Adherence to guidelines for creatinine and potassium monitoring and discontinuation following renin-angiotensin system blockade: a UK general practice-based cohort study. *BMJ Open*. 2017;7(1):e012818.
- Sarginson J, Webb RT, Stocks SJ, Esmail A, Garg S, Ashcroft DM. Temporal trends in antidepressant prescribing to children in UK primary care, 2000-2015. *J Affect Disord*. 2017;210:312-8.
- Sanni OB, Mc Menamin ÚC, Cardwell CR, Sharp L, Murray LJ, Coleman HG. Commonly used medications and endometrial cancer survival: a population-based cohort study. *Br J Cancer*. 2017;117(3):432-8.
- Saine ME, Gizaw M, Carbonari DM, Newcomb CW, Roy JA, Cardillo S, et al. Validity of diagnostic codes to identify hospitalizations for infections among patients treated with oral anti-diabetic drugs. *Pharmacoepidemiol Drug Saf*. 2017.
- Rothnie KJ, Chandan JS, Goss HG, Müllerová H, Quint JK. Validity and interpretation of spirometric recordings to diagnose COPD in UK primary care. *International Journal of COPD*. 2017;12:1663-8.
- Ronaldson SJ, Raghunath A, Torgerson DJ, Van Staa T. Cost-effectiveness of antibiotics for COPD

management: observational analysis using CPRD data. *ERJ open research*. 2017;3(2).

- Rodgers LR, Weedon MN, Henley WE, Hattersley AT, Shields BM. Cohort profile for the MASTERMIND study: using the Clinical Practice Research Datalink (CPRD) to investigate stratification of response to treatment in patients with type 2 diabetes. *BMJ Open*. 2017;7(10):e017989.
- Roddy E, Muller S, Paskins Z, Hider SL, Blagojevic-Bucknall M, Mallen CD. Incident acute pseudogout and prior bisphosphonate use: Matched case-control study in the UK-Clinical Practice Research Datalink. *Medicine*. 2017;96(12):e6177.
- Robbins TD, Tucker RCS, Arvanitis TN. Secondary use of Electronic Health Records (EHRs) in diabetes research: to what extent have we unlocked their potential? 2017.
- Riera-Montes M, Cattaert T, Verstraeten T. Rotavirus vaccination may reduce acute GA gastroenteritis rates across all age groups in England. *Value in Health*. 2017;20(9):A780.
- Renoux C, Dell'Aniello S, Brenner B, Suissa S. Bias from depletion of susceptibles: the example of hormone replacement therapy and the risk of venous thromboembolism: Depletion of Susceptible Bias: HRT and VTE. *Pharmacoepidemiol Drug Saf*. 2017.
- Ravindrarajah R, Hazra NC, Hamada S, Charlton J, Jackson SHD, Dregan A, et al. Systolic Blood Pressure Trajectory, Frailty and All-Cause Mortality Over 80 Years of Age. Cohort Study Using Electronic Health Records. *Circulation*. 2017.
- Ravindrarajah R, Dregan A, Hazra NC, Hamada S, Jackson SHD, Gulliford MC. Declining blood pressure and intensification of blood pressure management among people over 80 years: cohort study using electronic health records. *Journal of hypertension*. 2017;35(6):1276-82.
- Ramtoola S, Ploug UJ, Kragh N, Nyeland ME. Characteristics of newly diagnosed adults with Type 1 diabetes in the UK and their evolution over the first 5 years. 2017.
- Rajaram S, Wiecek W, Lawson R, Blak B, Zhao Y, Hackett J, et al. A retrospective observational analysis of post-pandemic influenza-related outcomes in the United Kingdom, 2010-2014. *Hum Vaccin Immunother*. 2017:0.
- Rahimi K, Mohseni H, Otto CM, Conrad N, Tran J, Nazarzadeh M, et al. Elevated blood pressure and risk of mitral regurgitation: A longitudinal cohort study of 5.5 million United Kingdom adults. *PLoS Med*. 2017;14(10):e1002404.
- Pufulete M, Maishman R, Dabner L, Mohiuddin S, Hollingworth W, Rogers CA, et al. Effectiveness and cost-effectiveness of serum B-type natriuretic peptide testing and monitoring in patients with heart failure in primary and secondary care: an evidence synthesis, cohort study and cost-effectiveness model. *Health Technol Assess*. 2017;21(40):1-150.
- Perry DC, Metcalfe D, Costa ML, Van Staa T. A nationwide cohort study of slipped capital femoral epiphysis. *Arch Dis Child*. 2017.
- Parisi R, Webb RT, Carr MJ, Moriarty KJ, Kleyn CE, Griffiths CEM, et al. Alcohol-Related Mortality in Patients With Psoriasis: A Population-Based Cohort Study. *JAMA dermatology*. 2017.
- Palmer KT, D'Angelo S, Harris EC, Linaker C, Sayer AA, Gale CR, et al. Sleep disturbance and the older worker: findings from the Health and Employment after Fifty study. *Scand J Work Environ Health*. 2017.

- Palmer KT, D'Angelo S, Harris EC, Linaker C, Gale CR, Evandrou M, et al. Frailty, prefrailty and employment outcomes in Health and Employment After Fifty (HEAF) Study. *Occup Environ Med*. 2017.
- Oshagbemi OA, Burden AM, Braeken DCW, Henskens Y, Wouters EFM, Driessen JHM, et al. Stability of Blood Eosinophils in COPD and Controls and the Impact of Gender, Age, Smoking and Baseline Counts. *Am J Respir Crit Care Med*. 2017.
- Nwaru BI, Simpson CR, Sheikh A, Kotz D. External validation of a COPD prediction model using population-based primary care data: a nested case-control study. *Scientific Reports*. 2017;7:44702.
- Numbere B, Fleming KM, Walker A, Card TR. Adrenergic blockers and the risk for common solid cancers: a case-control study. *European Journal of Cancer Prevention*. 2017;26(1):86-93.
- Nissen F, Morales DR, Mullerova H, Smeeth L, Douglas IJ, Quint JK. Validation of asthma recording in the Clinical Practice Research Datalink (CPRD). *BMJ Open*. 2017;7(8):e017474.
- Nightingale AL, Davidson JE, Molta CT, Kan HJ, McHugh NJ. Presentation of SLE in UK primary care using the Clinical Practice Research Datalink. *Lupus Sci Med*. 2017;4(1):e000172.
- Newlove-Delgado T, Ford TJ, Hamilton W, Stein K, Ukoumunne OC. Prescribing of medication for attention deficit hyperactivity disorder among young people in the Clinical Practice Research Datalink 2005-2013: analysis of time to cessation. *Eur Child Adolesc Psychiatry*. 2017.
- Mounce LTA, Price S, Valderas JM, Hamilton W. Comorbid conditions delay diagnosis of colorectal cancer: a cohort study using electronic primary care records. *Br J Cancer*. 2017.
- Mosholder AD, Lee J-Y, Zhou EH, Kang EM, Ghosh M, Izem R, et al. Long-term risk of acute myocardial infarction, stroke and death with outpatient use of clarithromycin: a retrospective cohort study. *Am J Epidemiol*. 2017.
- Morris M, Woods LM, Bhaskaran K, Rachev B. Do pre-diagnosis primary care consultation patterns explain deprivation-specific differences in net survival among women with breast cancer? An examination of individually-linked data from the UK West Midlands cancer registry, national screening programme and Clinical Practice Research Datalink. *BMC Cancer*. 2017;17(1).
- Morgan CL, Inzucchi S, Jenkins-Jones S, Puelles J, Currie C. Impact of pioglitazone on stroke outcomes: A real world database analysis. *Diabetologia*. 2017;60(1):S139-S40.
- Morgan C, Webb RT, Carr MJ, Kontopantelis E, Green J, Chew-Graham CA, et al. Incidence, clinical management, and mortality risk following self harm among children and adolescents: cohort study in primary care. *BMJ*. 2017;359:j4351.
- Morales DR, Lipworth BJ, Donnan PT, Jackson C, Guthrie B. Respiratory effect of beta-blockers in people with asthma and cardiovascular disease: population-based nested case control study. *BMC Med*. 2017;15(1):18.
- Moore E, Newson R, Joshi M, Palmer T, Rothnie KJ, Singh S, et al. Effects of pulmonary rehabilitation on exacerbation number and severity in people with COPD: An historical cohort study using electronic health records. *Chest*. 2017.
- Miani C, Martin A, Exley J, Doble B, Wilson E, Payne R, et al. Clinical effectiveness and cost-effectiveness of issuing longer versus shorter duration (3-month vs. 28-day) prescriptions in patients with chronic conditions: systematic review and economic modelling. *Health Technol Assess*. 2017;21(78):1-128.

- Mehta HB, Mehta V, Goodwin JS. Association of Hypoglycemia With Subsequent Dementia in Older Patients With Type 2 Diabetes Mellitus. *J Gerontol A Biol Sci Med Sci*. 2017;72(8):1110-6.
- McGrogan A, Snowball J, Charlton RA. Statins during pregnancy: a cohort study using the General Practice Research Database to investigate pregnancy loss. *Pharmacoepidemiol Drug Saf*. 2017.
- McDonald L, Ramagopalan S, Burns L, Postema R. Identifying patients with lupus nephritis in the united kingdom (UK) clinical practice research datalink (CPRD). *Value in Health*. 2017;20(9):A743.
- Mathur R, Bhaskaran K, Edwards E, Lee H, Chaturvedi N, Smeeth L, et al. Population trends in the 10-year incidence and prevalence of diabetic retinopathy in the UK: a cohort study in the Clinical Practice Research Datalink 2004–2014. *BMJ Open*. 2017;7(2):e014444.
- Martín-Merino E, Petersen I, Hawley S, Álvarez-Gutierrez A, Khalid S, Llorente-Garcia A, et al. Risk of venous thromboembolism among users of different anti-osteoporosis drugs: a population-based cohort analysis including over 200,000 participants from Spain and the UK. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2017.
- Mars B, Heron J, Gunnell D, Martin RM, Thomas KH, Kessler D. Prevalence and patterns of antidepressant switching amongst primary care patients in the UK. *Journal of psychopharmacology (Oxford, England)*. 2017;31(5):553-60.
- Margulis AV, Fortuny J, Kaye JA, Calingaert B, Reynolds M, Plana E, et al. Validation of Cancer Cases Using Primary Care, Cancer Registry, and Hospitalization Data in the UK. *Epidemiology*. 2017.
- Maguire A, Johnson ME, Denning DW, Ferreira GLC, Cassidy A. Identifying rare diseases using electronic medical records: the example of allergic bronchopulmonary aspergillosis. *Pharmacoepidemiol Drug Saf*. 2017.
- Loo SY, Dell'Aniello S, Huiart L, Renoux C. Trends in the prescription of novel oral anticoagulants in UK primary care. *British journal of clinical pharmacology*. 2017.
- Little I, Vinogradova Y, Orton E, Kai J, Qureshi N. Venous thromboembolism in adults screened for sickle cell trait: a population-based cohort study with nested case–control analysis. *BMJ Open*. 2017;7(3):e012665.
- Liang H, Manne S, Shick J, Lissos T, Dolin P. Incidence, prevalence, and natural history of primary sclerosing cholangitis in the United Kingdom. *Medicine*. 2017;96(24):e7116.
- Li L, Neogi T, Jick S. A cohort study of comorbidity in patients with granulomatosis with polyangiitis. *Rheumatology (Oxford)*. 2017.
- Li L, Neogi T, Jick S. Giant cell arteritis and vascular disease–risk factors and outcomes: a cohort study using UK Clinical Practice Research Datalink. *Rheumatology (Oxford)*. 2017.
- Leite A, Thomas SL, Andrews NJ. Do delays in data availability limit the implementation of near real-time vaccine safety surveillance using the Clinical Practice Research Datalink? *Pharmacoepidemiol Drug Saf*. 2017.
- Leite A, Thomas SL, Andrews NJ. Implementing near real-time vaccine safety surveillance using the Clinical Practice Research Datalink (CPRD). *Vaccine*. 2017.
- Leite A, Andrews NJ, Thomas SL. Assessing recording delays in general practice records to inform near real-time vaccine safety surveillance using the Clinical Practice Research Datalink (CPRD). *Pharmacoepidemiol*

Drug Saf. 2017.

- Leal J, Gray AM, Hawley S, Prieto-Alhambra D, Delmestri A, Arden NK, et al. Cost-Effectiveness of Orthogeriatric and Fracture Liaison Service Models of Care for Hip Fracture Patients: A Population-Based Study: COST-EFFECTIVENESS OF FRACTURE LIAISON SERVICE MODELS. *Journal of Bone and Mineral Research*. 2017;32(2):203-11.
- Larivée N, Suissa S, Eberg M, Joseph L, Eisenberg MJ, Abenhaim HA, et al. Drospirenone-containing combined oral contraceptives and the risk of arterial thrombosis: a population-based nested case-control study. *BJOG : an international journal of obstetrics and gynaecology*. 2017;124(11):1672-9.
- Larivée N, Suissa S, Coulombe J, Tagalakis V, Filion KB. Drospirenone-Containing Oral Contraceptive Pills and the Risk of Venous Thromboembolism: An Assessment of Risk in First-Time Users and Restarters. *Drug Safety*. 2017;40(7):583-96.
- Langham S, Wright A, Shaw H, Bryson S, O'Leary S. Budget impact analysis of oral glycopyrronium bromide (Sialanar™) for the symptomatic treatment of severe sialorrhoea (Drooling) in the UK setting. *Value in Health*. 2017;20(9):A720.
- Langham J, Langham S. European real world data sources to characterise flares in patients with systemic lupus erythematosus: A review. *Value in Health*. 2017;20(9):A571.
- Lane S, Lynn E, Shakir S. Measuring the impact of product withdrawals and other major pharmacovigilance actions on public health burden in the EU: Methodological considerations. *Drug Safety*. 2017;40(10):953-4.
- Lane DA, Skjøth F, Lip GYH, Larsen TB, Kotecha D. Temporal Trends in Incidence, Prevalence, and Mortality of Atrial Fibrillation in Primary Care. *Journal of the American Heart Association*. 2017;6(5).
- Landis SH, Suruki R, Hilton E, Compton C, Galwey NW. Stability of Blood Eosinophil Count in Patients with COPD in the UK Clinical Practice Research Datalink. *COPD*. 2017:1-7.
- Lan W-C, Yu K-H, Luo S-F, Kuo C-F, Chan T-M. Effect of urate-lowering treatment on the risk of urolithiasis in people with gout. *Arthritis and Rheumatology*. 2017;69.
- Lamy FX, Saragoussi D, Johnson ME, Guiraud-Diawara A, Jørgensen KT, Loze JY, et al. The use of adjunctive antipsychotics to treat depression in UK primary care. *Current Medical Research and Opinion*. 2017:1-8.
- Lamy FX, Saragoussi D, Johnson ME, Guiraud-Diawara A, Jorgensen KT, Loze JY, et al. The use of adjunctive antipsychotics to treat depression in UK primary care. *Curr Med Res Opin*. 2017;33(5):891-8.
- Lambrelli D, Merinopoulou E, Donaldson R, Mardekian J, Hlavacek P, Tarasenko L, et al. Real-world treatment patterns among patients initiating on statins in England. *Value in Health*. 2017;20(9):A630-A1.
- Lacoïn L, Lumley M, Ridha E, Pereira M, McDonald L, Ramagopalan S, et al. Evolving landscape of stroke prevention in atrial fibrillation within the UK between 2012 and 2016: a cross-sectional analysis study using CPRD. *BMJ Open*. 2017;7(9):e015363.
- Kurdi AI, Chen L-C, Elliott RA. Exploring factors associated with patients' adherence to antihypertensive drugs among people with primary hypertension in the United Kingdom. *Journal of hypertension*. 2017.
- Klil-Drori AJ, Yin H, Abenhaim HA, Galbaud du Fort G, Azoulay L. Prolactin-Elevating Antipsychotics and the Risk of Endometrial Cancer. *The Journal of Clinical Psychiatry*. 2017.

- Kiconco S, Abdul Sultan A, Grainge MJ. Recurrence risk of venous thromboembolism and hormone use in women from England: a cohort study using clinical practice research datalink. *Br J Haematol*. 2017.
- Khosrow-Khavar F, Yin H, Barkun A, Bouganim N, Azoulay L. Aromatase Inhibitors and the Risk of Colorectal Cancer in Post-Menopausal Women with Breast Cancer. *Annals of oncology : official journal of the European Society for Medical Oncology*. 2017.
- Khan UT, Walker AJ, Baig S, Card TR, Kirwan CC, Grainge MJ. Venous thromboembolism and mortality in breast cancer: Cohort study with systematic review and meta-analysis. *BMC Cancer*. 2017;17(1).
- Kerkhof M, Tran TN, Soriano JB, Golam S, Gibson D, Hillyer EV, et al. Healthcare resource use and costs of severe, uncontrolled eosinophilic asthma in the UK general population. *Thorax*. 2017.
- Kaye JA, Margulis AV, Fortuny J, McQuay LJ, Plana E, Bartsch JL, et al. Cancer Incidence after Initiation of Antimuscarinic Medications for Overactive Bladder in the United Kingdom: Evidence for Protopathic Bias. *Pharmacotherapy*. 2017.
- Judge A, Garriga C, Arden NK, Lovestone S, Prieto-Alhambra D, Cooper C, et al. Protective effect of antirheumatic drugs on dementia in rheumatoid arthritis patients. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2017;3(4):612-21.
- Joseph RM, Soames J, Wright M, Sultana K, van Staa TP, Dixon WG. Supplementing electronic health records through sample collection and patient diaries: A study set within a primary care research database. *Pharmacoepidemiol Drug Saf*. 2017.
- Joseph RM, Movahedi M, Dixon WG, Symmons DP. Risks of smoking and benefits of smoking cessation on hospitalisations for cardiovascular events and respiratory infection in patients with rheumatoid arthritis: a retrospective cohort study using the Clinical Practice Research Datalink. *RMD open*. 2017;3(2):e000506.
- Jick S, Li L, Gastanaga VM, Liede A, Hernandez RK. Prevalence of hypercalcemia of malignancy among pediatric cancer patients in the UK Clinical Practice Research Datalink database. *Clin Epidemiol*. 2017;9:339-43.
- Jenkins-Jones S, Currie C, Tomazos IC, Donato BM, Bishop N, Eastell R. Characterizing the epidemiology of hypophosphatasia in a UK population. *Calcif Tissue Int*. 2017;100(1):S141-S2.
- Jain A, van Hoek AJ, Walker JL, Mathur R, Smeeth L, Thomas SL. Identifying social factors amongst older individuals in linked electronic health records: An assessment in a population based study. *PLoS one*. 2017;12(11):e0189038.
- Jain A, McDonald HI, Nitsch D, Tomlinson L, Thomas SL. Risk factors for developing acute kidney injury in older people with diabetes and community-acquired pneumonia: a population-based UK cohort study. *BMC nephrology*. 2017;18(1):142.
- Iwagami M, Tomlinson LA, Mansfield KE, McDonald HI, Smeeth L, Nitsch D. Prevalence, incidence, indication, and choice of antidepressants in patients with and without chronic kidney disease: a matched cohort study in UK Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf*. 2017.
- Iwagami M, Tomlinson LA, Mansfield KE, Casula A, Caskey FJ, Aitken G, et al. Validity of estimated prevalence of decreased kidney function and renal replacement therapy from primary care electronic health records compared with national survey and registry data in the United Kingdom. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal*

Association. 2017;32(suppl_2):ii142-ii50.

- Ingram JR, Jenkins-Jones S, Knipe DW, Morgan CLI, Cannings-John R, Piguet V. Population-based Clinical Practice Research Datalink study using algorithm modelling to identify the true burden of hidradenitis suppurativa. *Br J Dermatol*. 2017.
- Humphreys JH, Warner A, Costello R, Lunt M, Verstappen SMM, Dixon WG. Quantifying the hepatotoxic risk of alcohol consumption in patients with rheumatoid arthritis taking methotrexate. *Annals of the Rheumatic Diseases*. 2017:annrheumdis-2016-210629.
- Holden SE, Morgan CL, Qiao Q, Jenkins-Jones S, Berni ER, Currie CJ. Healthcare resource utilization and related financial costs associated with glucose lowering with either exenatide or basal insulin: a retrospective cohort study. *Diabetes, Obesity and Metabolism*. 2017.
- Holden SE, Jenkins-Jones S, Morgan CL, Peters JR, Scherthaner G, Currie CJ. Prevalence, glucose control and relative survival of people with Type 2 diabetes in the UK from 1991 to 2013. *Diabetic Medicine*. 2017.
- Holbrook T, Tang Y, Das R, Shankar RR, Tunceli K, Williams J, et al. Direct medical costs of severe hypoglycaemic events in patients with type 2 diabetes in England: A retrospective database study. *Int J Clin Pract*. 2017.
- Hicks BM, Yin H, Sinyavskaya L, Suissa S, Azoulay L, Brassard P. Metformin and the incidence of viral associated cancers in patients with type 2 diabetes. *Int J Cancer*. 2017.
- Hicks BM, Yin H, Bladou F, Ernst P, Azoulay L. Androgen deprivation therapy for prostate cancer and the risk of hospitalisation for community-acquired pneumonia. *Thorax*. 2017;72(7):596-7.
- Hicks BM, Klil-Drori AJ, Yin H, Campeau L, Azoulay L. Androgen Deprivation Therapy and the Risk of Anemia in Men with Prostate Cancer. *Epidemiology*. 2017;28(5):712-8.
- Herrington WG, Smith M, Bankhead C, Matsushita K, Stevens S, Holt T, et al. Body-mass index and risk of advanced chronic kidney disease: Prospective analyses from a primary care cohort of 1.4 million adults in England. *PLOS ONE*. 2017;12(3):e0173515.
- He X, Zhao Z, Curtis BH, Faries DE, Fu H, Peng X. Machine-learning algorithms for personalized medicine-applications for patients with type 2 diabetes initiating injectable treatments. *Diabetes*. 2017;66:A251.
- Hazra NC, Rudisill C, Gulliford MC. Determinants of health care costs in the senior elderly: age, comorbidity, impairment, or proximity to death? *The European journal of health economics : HEPAC : health economics in prevention and care*. 2017.
- Hawley S, Cordtz R, Dreyer L, Edwards CJ, Arden NK, Delmestri A, et al. Association between NICE guidance on biologic therapies with rates of hip and knee replacement among rheumatoid arthritis patients in England and Wales: An interrupted time-series analysis. *Seminars in arthritis and rheumatism*. 2017.
- Hansson E, Forbes HJ, Langan SM, Smeeth L, Bhaskaran K. Herpes zoster risk after 21 specific cancers: population-based case-control study. *Br J Cancer*. 2017.
- Hamilton G, Diels J, Schubert A. Characteristics of patients with type 2 diabetes initiating canagliflozin compared with other treatment regimens in the UK. *Diabetologia*. 2017;60(1):S401.
- Hall N, Stauble F, Zahra M, Thavaneswaran S, Eldabe S. Impact of timing of spinal cord stimulation on medication utilisation long term: Results from the CPRD database. *British Journal of Pain*. 2017;11(2):44.

- Hagberg KW, Jick SS. Validation of autism spectrum disorder diagnoses recorded in the clinical practice research datalink, 1990-2014. *Clin Epidemiol.* 2017;9:475-82.
- Hagberg KW, Divan HA, Nickel JC, Jick SS. Risk of Incident Antidepressant-Treated Depression Associated with Use of 5- α -Reductase Inhibitors Compared with Use of α -Blockers in Men with Benign Prostatic Hyperplasia: A Population-Based Study Using the Clinical Practice Research Datalink. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy.* 2017.
- Hagberg KW, Divan H, Fang SC, Nickel JC, Jick S. Risk of gynecomastia and breast cancer associated with the use of 5-alpha reductase inhibitors for benign prostatic hyperplasia. *Clin Epidemiol.* 2017;Volume 9:83-91.
- Gulliford M, Ravindrarajah R, Hamada S, Jackson S, Charlton J. Inception and deprescribing of statins in people aged over 80 years: cohort study. *Age Ageing.* 2017;46(6):1001-5.
- Griffiths RI, Valderas JM, McFadden EC, Bankhead CR, Lavery BA, Khan NF, et al. Outcomes of preexisting diabetes mellitus in breast, colorectal, and prostate cancer. *Journal of cancer survivorship : research and practice.* 2017.
- Grandi SM, Vallée-Pouliot K, Reynier P, Eberg M, Platt RW, Arel R, et al. Hypertensive Disorders in Pregnancy and the Risk of Subsequent Cardiovascular Disease. *Paediatr Perinat Epidemiol.* 2017.
- Gorricho J, Garjón J, Alonso A, Celaya MC, Saiz LC, Erviti J, et al. Use of oral antidiabetic agents and risk of community-acquired pneumonia: a nested case-control study: Use of oral antidiabetic agents and risk of community-acquired pneumonia. *British Journal of Clinical Pharmacology.* 2017.
- Gordon J, McEwan P, Evans M, Puelles J, Sinclair A. Managing glycaemia in older people with type 2 diabetes: A retrospective, primary care-based cohort study, with economic assessment of patient outcomes. *Diabetes Obes Metab.* 2017;19(5):644-53.
- Gieling E, van den Ham HA, van Onzenoort H, Bos J, Kramers C, de Boer A, et al. Risk of major bleeding and stroke associated with the use of VKAs, NOACs and aspirin in patients with atrial fibrillation: a cohort study: Risk of bleeding and stroke of anti-thrombotic treatment. *British Journal of Clinical Pharmacology.* 2017.
- Gielen V, Calado F, Balas B, Proenca CC, Bruce Wirta S, McMahon P, et al. Hospitalisations among a cohort of patients with heart failure versus an age-and sex-matched cohort without heart failure in England: A comparative study using CPRD data. *Value in Health.* 2017;20(9):A512.
- Gentil M-L, Cuggia M, Fiquet L, Hagenbourger C, Le Berre T, Banâtre A, et al. Factors influencing the development of primary care data collection projects from electronic health records: a systematic review of the literature. *BMC Med Inform Decis Mak.* 2017;17(1):139.
- Gamble J-M, Donnan JR, Chibrikov E, Twells LK, Midodzi WK, Majumdar SR. The risk of fragility fractures in new users of dipeptidyl peptidase-4 inhibitors compared to sulfonylureas and other anti-diabetic drugs: A cohort study. *Diabetes research and clinical practice.* 2017;136:159-67.
- Frey N, Jossi J, Bodmer M, Bircher A, Jick SS, Meier CR, et al. The Epidemiology of Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis in the UK. *Journal of Investigative Dermatology.* 2017.
- Frey N, Hügle T, Jick SS, Meier CR, Spöndlin J. Hyperlipidaemia and incident osteoarthritis of the hand: a population-based case-control study. *Osteoarthritis and Cartilage.* 2017.
- Frey N, Bodmer M, Bircher A, Rüegg S, Jick SS, Meier CR, et al. The risk of Stevens-Johnson syndrome and toxic epidermal necrolysis in new users of antiepileptic drugs. *Epilepsia.* 2017.

- Frey N, Bircher A, Bodmer M, Jick SS, Meier CR, Spoendlin J. Antibiotic drug use and the risk of Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis A Population-Based Case-Control Study. *J Invest Dermatol*. 2017.
- Frey N, Bircher A, Bodmer M, Jick SS, Meier CR, Spoendlin J. Validation of Stevens–Johnson syndrome or toxic epidermal necrolysis diagnoses in the Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf*. 2017;26(4):429-36.
- Franklin M, Davis S, Horspool M, Kua WS, Julious S. Economic Evaluations Alongside Efficient Study Designs Using Large Observational Datasets: the PLEASANT Trial Case Study. *Pharmacoeconomics*. 2017.
- François C, Tanasescu A, Lamy F-X, Despiegel N, Falissard B, Chalem Y, et al. Creating an index to measure health state of depressed patients in automated healthcare databases: the methodology. *Journal of market access & health policy*. 2017;5(1):1372025.
- Francis NA, Entwistle K, Santer M, Layton AM, Eady EA, Butler CC. The management of acne vulgaris in primary care: a cohort study of consulting and prescribing patterns using the Clinical Practice Research Datalink. *Br J Dermatol*. 2017;176(1):107-15.
- Ferreira GLC, Marano C, De Moerlooze L, Guignard A, Feng Y, El Hahi Y, et al. Incidence and prevalence of hepatitis B in patients with diabetes mellitus in the UK: a population-based cohort study using the UK Clinical Practice Research Datalink. *Journal of viral hepatitis*. 2017.
- Fernandes G, Hannachi H, Liu J, Wang T, McNeill A, Rajpathak S. Distance from glycaemic goal at the time of treatment intensification in patients with type 2 diabetes failing metformin monotherapy in the UK. *Diabetologia*. 2017;60(1):S302.
- Fazeli Farsani S, Raabe C, Girman CJ, Spencer W, George J, Verpillat P, et al. Characteristics of patients initiating empagliflozin or other noninsulin glucose lowering drugs (GLDs) in the United Kingdom (UK). *Diabetes*. 2017;66:A628.
- Farley A, Koshiaris C, Oke J, Ryan R, Szatkowski L, Stevens R, et al. Physician Support of Smoking Cessation After Diagnosis of Lung, Bladder, or Upper Aerodigestive Tract Cancer. *Annals of family medicine*. 2017;15(5):443-50.
- Eberg M, Platt RW, Filion KB. The Estimation of Gestational Age at Birth in Database Studies. *Epidemiology*. 2017.
- Durham TA, Hassmiller Lich K, Viera AJ, Fine JP, Mukherjee J, Weinberger M, et al. Utilization of Standard and Target-Specific Oral Anticoagulants Among Adults in the United Kingdom With Incident Atrial Fibrillation. *Am J Cardiol*. 2017.
- Douros A, Yin H, Yu OHY, Filion KB, Azoulay L, Suissa S. Pharmacologic Differences of Sulfonylureas and the Risk of Adverse Cardiovascular and Hypoglycemic Events. *Diabetes care*. 2017.
- Dollerup J, Vestbo J, Murray-Thomas T, Kaplan A, Martin RJ, Pizzichini E, et al. Cardiovascular risks in smokers treated with nicotine replacement therapy: a historical cohort study. *Clin Epidemiol*. 2017;9:231-43.
- Doble B, Payne R, Harshfield A, Wilson ECF. Retrospective, multicohort analysis of the Clinical Practice Research Datalink (CPRD) to determine differences in the cost of medication wastage, dispensing fees and prescriber time of issuing either short (<60 days) or long (≥60 days) prescription lengths in primary care for common, chronic conditions in the UK. *BMJ Open*. 2017;7(12):e019382.

- Diels J, Schubert A, Hamilton G. Time to next therapy for patients with type 2 diabetes in the UK: Canagliflozin compared with other antihyperglycaemic agents. *Diabetologia*. 2017;60(1):S19.
- Dennis JM, Shields BM, Henley WE, Knight BA, McDonald TJ, Hill AV, et al. Clinical markers of insulin resistance predict reduced glycaemic response with DPP4-inhibitors: A MASTERMIND stratified medicine study. 2017.
- Dennis JM, Henley WE, Weedon MN, Lonergan M, Rodgers LR, Jones AG, et al. A calculator to predict durability of HbA1c response with DPP4 inhibitors, sulfonylureas and thiazolidinediones: A MASTERMIND precision medicine study. *Diabetologia*. 2017;60(1):S388.
- Dennis JM, Henley WE, Weedon M, Lonergan M, Jones AG, Sattar N, et al. Development of an online risk calculator to predict durability of good glycaemic control with sulfonylurea and thiazolidinedione therapy: A MASTERMIND stratified medicine study. 2017.
- De Jong HJ, van Staa TP, Lalmohamed A, de Vries F, Vandebriel RJ, Van Loveren H, et al. Pattern of risks of systemic lupus erythematosus among statin users: a population-based cohort study. *Annals of the rheumatic diseases*. 2017;76(10):1723-30.
- Davies NM, Thomas KH, Taylor AE, Taylor GMJ, Martin RM, Munafò MR, et al. How to compare instrumental variable and conventional regression analyses using negative controls and bias plots. *Int J Epidemiol*. 2017;46(6):2067-77.
- Daugherty J, Lin X, Baxter R, Suruki R, Bradford E. The impact of long-term systemic glucocorticoid use in severe asthma: A UK retrospective cohort analysis. *The Journal of asthma : official journal of the Association for the Care of Asthma*. 2017:1-8.
- Danese MD, Gleeson M, Kutikova L, Griffiths RI, Khunti K, Seshasai SRK, et al. Management of lipid-lowering therapy in patients with cardiovascular events in the UK: a retrospective cohort study. *BMJ Open*. 2017;7(5):e013851.
- Danese MD, Gleeson M, Griffiths RI, Catterick D, Kutikova L. Methods for estimating costs in patients with hyperlipidemia experiencing their first cardiovascular event in the United Kingdom. *J Med Econ*. 2017:1-7.
- Danese M, Sidelnikov E, Kutikova L. The prevalence of patients at very high risk for cardiovascular events in the United Kingdom. *Value in Health*. 2017;20(9):A605.
- Danese M, Kutikova L, Catterick D, Sorio-Vilela F, Gleeson M, Khunti K, et al. Characteristics and risk factor burden of patients receiving lipid modifying therapy in The United Kingdom. *Value in Health*. 2017;20(9):A605-A6.
- Currie CJ, Holden S, Jenkins-Jones S, Morgan CL, Voss B, Rajpathak SN, et al. The impact of differing glucose-lowering regimens on the pattern of association between glucose control and survival. *Diabetes Obes Metab*. 2017.
- Curran D, Hunjan M, El Ghachi A, El Hahi Y, Bianco V, Ferreira G. Herpes zoster related healthcare burden and costs in both immunocompromised (IC) and IC-free populations in the United Kingdom. *Value in Health*. 2017;20(9):A786.
- Cooper AJM, Willis J, Fuller J, Benecke H, Leighton-Scott J, Andersohn F, et al. Prevalence and Incidence Trends for Diagnosed Prescription Opioid Use Disorders in the United Kingdom. *Pain and therapy*. 2017.
- Conradt C, Guo D, Miclea A, Nisslein T, Ismail C, Chatamra K, et al. Increased Prevalence of Polyneuropathy

in Parkinson's Disease Patients: An Observational Study. *Journal of Parkinson's disease*. 2017.

- Conrad N, Judge A, Tran J, Mohseni H, Hedgecote D, Crespillo AP, et al. Temporal trends and patterns in heart failure incidence: a population-based study of 4 million individuals. *Lancet*. 2017.
- Connell OL, Rothnie KJ, Quint JK. Risk of stroke associated with acute exacerbations of chronic obstructive pulmonary disease (COPD): A self-controlled case series. *Thorax*. 2017;72:A137-A8.
- Collin SM, Bakken IJ, Nazareth I, Crawley E, White PD. Trends in the incidence of chronic fatigue syndrome and fibromyalgia in the UK, 2001-2013: a Clinical Practice Research Datalink study. *J R Soc Med*. 2017;110(6):231-44.
- Collin SM, Bakken IJ, Nazareth I, Crawley E, White PD. Health care resource use by patients before and after a diagnosis of chronic fatigue syndrome (CFS/ME): a clinical practice research datalink study. *BMC Fam Pract*. 2017;18(1):60.
- Chiquette E, Oral EA, Garg A, Araújo-Vilar D, Dhankhar P. Estimating the prevalence of generalized and partial lipodystrophy: Findings and challenges. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. 2017;10:375-83.
- Chen T-C, Chen L-C, Knaggs RD. A 15-year overview of increasing tramadol utilisation and associated mortality and the impact of tramadol classification in the United Kingdom. *Pharmacoepidemiol Drug Saf*. 2017.
- Chen J-S, Kuo C-F, Tsai P-H, Luo S-F, Yu K-H. Urate-lowering treatment and risk of total joint replacement in patients with incident gout: A population-based cohort and nested case-control study. *Arthritis and Rheumatology*. 2017;69.
- Chaudhry Z, Mannan F, Gibson-White A, Syed U, Ahmed S, Kousoulis A, et al. Outputs and Growth of Primary Care Databases in the United Kingdom: Bibliometric Analysis. *J Innov Health Inform*. 2017;24(3):942.
- Charlton RA, McGrogan A, Snowball J, Yates LM, Wood A, Clayton-Smith J, et al. Sensitivity of the UK Clinical Practice Research Datalink to Detect Neurodevelopmental Effects of Medicine Exposure in Utero: Comparative Analysis of an Antiepileptic Drug-Exposed Cohort. *Drug Safety*. 2017.
- Charlton R, Green A, Shaddick G, Snowball J, Nightingale A, Tillett W, et al. Risk of uveitis and inflammatory bowel disease in people with psoriatic arthritis: a population-based cohort study. *Annals of the rheumatic diseases*. 2017.
- Charlton J, Ravindrarajah R, Hamada S, Jackson SH, Gulliford MC. Trajectory of Total Cholesterol in The Last Years of Life Over Age 80 Years. Cohort Study of 99,758 Participants. *J Gerontol A Biol Sci Med Sci*. 2017.
- Chapple CR, Nazir J, Hakimi Z, Bowditch S, Fatoye F, Guelfucci F, et al. Persistence and Adherence with Mirabegron versus Antimuscarinic Agents in Patients with Overactive Bladder: A Retrospective Observational Study in UK Clinical Practice. *European Urology*. 2017.
- Chalmers JD, Tebboth A, Gayle A, Ternouth A, Ramscar N. Determinants of initial inhaled corticosteroid use in patients with GOLD A/B COPD: a retrospective study of UK general practice. *NPJ Prim Care Respir Med*. 2017;27(1):43.
- Cattaert T, Ferreira G, Verstraeten T. Increased episode rate of medically-attended acute gastroenteritis in diabetics in England. *Value in Health*. 2017;20(9):A781.

- Caspard H, Wise RP, Steffey A, Brody RS. Incidence of live-attenuated influenza vaccine administration beyond expiry date in children and adolescents aged 2-17 years in the UK: a population-based cohort study. *BMJ Open*. 2017;7(7):e016520.
- Carreira H, Williams R, Müller M, Harewood R, Bhaskaran K. Adverse mental health outcomes in breast cancer survivors compared to women who did not have cancer: Systematic review protocol. *Systematic Reviews*. 2017;6(1).
- Carr MJ, Ashcroft DM, Kontopantelis E, While D, Awenat Y, Cooper J, et al. Premature Death Among Primary Care Patients With a History of Self-Harm. *Annals of family medicine*. 2017;15(3):246-54.
- But A, De Bruin ML, Bazelier MT, Hjellvik V, Andersen M, Auvinen A, et al. Cancer risk among insulin users: comparing analogues with human insulin in the CARING five-country cohort study. *Diabetologia*. 2017.
- Busby J, McMenemy Ú, Spence A, Johnston BT, Hughes C, Cardwell CR. Angiotensin receptor blocker use and gastro-oesophageal cancer survival: a population-based cohort study. *Aliment Pharmacol Ther*. 2017.
- Burn E, Arden NK, Edwards C, Cooper C, Murray D, Prieto-Alhambra D, et al. Lifetime risk of revision following joint replacement: Evidence from routinely-collected data. *Value in Health*. 2017;20(9):A526.
- Bruderer SG, Meier CR, Jick SS, Bodmer M. The association between thyroid disorders and incident gout: population-based case-control study. *Clin Epidemiol*. 2017;9:205-15.
- Bruderer SG, Bodmer D, Stohler NA, Jick SS, Meier CR. Population-Based Study on the Epidemiology of Ménière's Disease. *Audiology & neuro-otology*. 2017;22(2):74-82.
- Browne J, Edwards DA, Rhodes KM, Brimicombe DJ, Payne RA. Association of comorbidity and health service usage among patients with dementia in the UK: a population-based study. *BMJ Open*. 2017;7(3):e012546.
- Bronsveld HK, Peeters PJHL, de Groot MCH, de Boer A, Schmidt MK, De Bruin ML. Trends in breast cancer incidence among women with type-2 diabetes in British general practice. *Primary Care Diabetes*. 2017.
- Braeken DC, Rohde GG, Franssen FM, Driessen JH, van Staa TP, Souverein PC, et al. Risk of community-acquired pneumonia in chronic obstructive pulmonary disease stratified by smoking status: a population-based cohort study in the United Kingdom. *Int J Chron Obstruct Pulmon Dis*. 2017;12:2425-32.
- Bowman K, Delgado J, Henley WE, Masoli JA, Kos K, Brayne C, et al. Obesity in Older People With and Without Conditions Associated With Weight Loss: Follow-up of 955,000 Primary Care Patients. *J Gerontol A Biol Sci Med Sci*. 2017;72(2):203-9.
- Bouras G, Markar SR, Burns EM, Huddy JR, Bottle A, Athanasiou T, et al. The psychological impact of symptoms related to esophagogastric cancer resection presenting in primary care: A national linked database study. *Eur J Surg Oncol*. 2017;43(2):454-60.
- Bouras G, Burns EM, Howell AM, Bottle A, Athanasiou T, Darzi A. Linked hospital and primary care database analysis of the impact of short-term complications on recurrence in laparoscopic inguinal hernia repair. *Hernia*. 2017.
- Bottle A, Kim D, Aylin P, Cowie MR, Majeed A, Hayhoe B. Routes to diagnosis of heart failure: observational study using linked data in England. *Heart*. 2017.
- Borghs S, Thieffry S, Noack-Rink M, Dedeken P, Hong LS, Byram L, et al. Health care cost associated with the use of enzyme-inducing and non-enzyme-active antiepileptic drugs in the UK: a long-term retrospective

matched cohort study. *BMC Neurology*. 2017;17(1).

- Bloom CI, Nissen F, Douglas IJ, Smeeth L, Cullinan P, Quint JK. Exacerbation risk and characterisation of the UK's asthma population from infants to old age. *Thorax*. 2017.
- Bloechliger M, Reinau D, Spoenclin J, Chang SC, Kuhlbusch K, Heaney LG, et al. Adverse events profile of oral corticosteroids among asthma patients in the UK. *Thorax*. 2017;72:A190.
- Ble A, Hughes PM, Delgado J, Masoli JA, Bowman K, Zirk-Sadowski J, et al. Safety and Effectiveness of Statins for Prevention of Recurrent Myocardial Infarction in 12 156 Typical Older Patients: A Quasi-Experimental Study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2017;72(2):243-50.
- Birks J, Bankhead C, Holt TA, Fuller A, Patnick J. Evaluation of a prediction model for colorectal cancer: retrospective analysis of 2.5 million patient records. *Cancer medicine*. 2017.
- Biétry FA, Hug B, Reich O, Susan JS, Meier CR. Iron supplementation in Switzerland - A bi-national, descriptive and observational study. *Swiss medical weekly*. 2017;147:w14444.
- Berni E, de Voogd H, Halcox JP, Butler CC, Bannister CA, Jenkins-Jones S, et al. Risk of cardiovascular events, arrhythmia and all-cause mortality associated with clarithromycin versus alternative antibiotics prescribed for respiratory tract infections: a retrospective cohort study. *BMJ Open*. 2017;7(1):e013398.
- Bennett H, McEwan P, Ayoubkhani D, Evans M, Qin L, Kim K, et al. Development of a mortality risk prediction tool for patients with heart failure or chronic kidney disease at risk of hyperkalaemia. *Value in Health*. 2017;20(9):A682.
- Bennett H, Ayoubkhani D, Evans M, McEwan P, Qin L, Grandy S, et al. Quantifying the relationship between hyperkalaemia and outcomes in patients with heart failure or chronic kidney disease. *Value in Health*. 2017;20(9):A682.
- Becker C, Jick SS, Meier CR, Bodmer M. Metformin and the risk of renal cell carcinoma: a case-control analysis. *European journal of cancer prevention : the official journal of the European Cancer Prevention Organisation (ECP)*. 2017;26(3):257-62.
- Becker C, Jick SS, Meier CR. Selective Serotonin Reuptake Inhibitors and Cataract Risk: A Case-Control Analysis. *Ophthalmology*. 2017.
- Bayliss LE, Culliford D, Monk AP, Glyn-Jones S, Prieto-Alhambra D, Judge A, et al. The effect of patient age at intervention on risk of implant revision after total replacement of the hip or knee: a population-based cohort study. *The Lancet*. 2017.
- Barker I, Steventon A, Deeny SR. Association between continuity of care in general practice and hospital admissions for ambulatory care sensitive conditions: cross sectional study of routinely collected, person level data. *BMJ*. 2017;356:j84.
- Baricault B, Palmaro A, Lapeyre-Mestre M. Mortality related to benzodiazepines or other psychotropic drugs in patients with cancer. *Clin Ther*. 2017;39(8):e12-e3.
- Ban L, Sprigg N, Abdul Sultan A, Nelson-Piercy C, Bath PM, Ludvigsson JF, et al. Incidence of First Stroke in Pregnant and Nonpregnant Women of Childbearing Age: A Population-Based Cohort Study From England. *Journal of the American Heart Association*. 2017;6(4).
- Baker R, Kendrick D, Tata LJ, Orton E. Association between maternal depression and anxiety episodes and

rates of childhood injuries: a cohort study from England. *Injury Prevention*. 2017;injuryprev-2016-042294.

- Bailey SE, Ukoumunne OC, Shephard EA, Hamilton W. Clinical relevance of thrombocytosis in primary care: a prospective cohort study of cancer incidence using English electronic medical records and cancer registry data. *Br J Gen Pract*. 2017;67(659):e405-e13.
- Badrick EL, Lennon H, Sperrin M, Van Staa T, O'Reilly D, Buchan I, et al. Can we identify people with higher pancreatic cancer risk who present with type 2 diabetes? *Diabetologia*. 2017;60(1):S562.
- Atkins J, Delgado J, Kuchel GA, Melzer D. Aging outcomes with nearly ideal simple seven cardiovascular health risks. *Journal of the American Geriatrics Society*. 2017;65:S115.
- Arhi C, Burns E, Bouras G, Aylin P, Darzi A, Ziprin P. Complications following colonic resection delay adjuvant chemotherapy treatment and reduce overall survival: A nationwide study based on routinely collected primary and secondary data. *Colorectal Disease*. 2017;19:46.
- Arden N, Altman D, Beard D, Carr A, Clarke N, Collins G, et al. Lower limb arthroplasty: can we produce a tool to predict outcome and failure, and is it cost-effective? An epidemiological study. Lower limb arthroplasty: can we produce a tool to predict outcome and failure, and is it cost-effective? An epidemiological study. Southampton (UK): NIHR Journals Library
- Copyright (c) Queen's Printer and Controller of HMSO 2017. This work was produced by Arden et al. under the terms of a commissioning contract issued by the Secretary of State for Health. This issue may be freely reproduced for the purposes of private research and study and extracts (or indeed, the full report) may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.; 2017.
- Ankarfeldt MZ, Thorsted BL, Groenwold RH, Adalsteinsson E, Ali MS, Klungel OH. Assessment of channeling bias among initiators of glucose-lowering drugs: A UK cohort study. *Clin Epidemiol*. 2017;9:19-30.
- Allen C, Meeraus W, Donegan K. Comparative risk of cause-specific mortality in older patients prescribed codeine or tramadol for non-malignant pain: Retrospective cohort study. *Drug Safety*. 2017;40(10):993-4.
- Allen C, Donegan K. The impact of regulatory action on the co-prescribing of renin-angiotensin system blockers in UK primary care. *Pharmacoepidemiol Drug Saf*. 2017.
- Alexandridou M, Bollaerts K. Zoster vaccine effectiveness against incident herpes zoster and post-herpetic neuralgia in elderly in the UK. *Value in Health*. 2017;20(9):A780.
- Alexakis C, Saxena S, Chhaya V, Cecil E, Curcin V, Pollok R. Do Thiopurines Reduce the Risk of Surgery in Elderly Onset Inflammatory Bowel Disease? A 20-Year National Population-Based Cohort Study. *Inflamm Bowel Dis*. 2017;23(4):672-80.
- Alatorre CI, Hoogwerf BJ, Deeg MA, Nelson DR, Hunter TM, Ng WT, et al. Factors associated with stroke, myocardial infarction, ischemic heart disease, unstable angina, or mortality in patients from real world clinical practice with newly diagnosed type 2 diabetes and early glycemic control. *Current medical research and opinion*. 2017:1-19.
- Ahmadizar F, Souverein P, de Boer A, Maitland-van der Zee AH. Under-treatment of hypertension and hypercholesterolemia in children and adolescents with type 1 diabetes- Long term follow-up on time trends

in the occurrence of cardiovascular disease, risk factors and medications use. *British journal of clinical pharmacology*. 2017.

- Achana FA, Fleming KM, Tata LJ, Sultan AA, Petrou S. Peripartum hysterectomy: an economic analysis of direct healthcare costs using routinely collected data. *BJOG : an international journal of obstetrics and gynaecology*. 2017.
- Achana F, Fleming KM, Tata LJ, Sultan AA, Petrou S. Peripartum hysterectomy: an economic analysis of direct health care costs using routinely collected data. *BJOG : an international journal of obstetrics and gynaecology*. 2017.
- Abrahami D, Yin H, Yu OHY, Pollak MN, Azoulay L. Incretin-based drugs and the incidence of colorectal cancer in patients with type 2 diabetes. *Epidemiology*. 2017.
- Abhishek A, Doherty M, Kuo C-F, Mallen CD, Zhang W, Grainge MJ. Rheumatoid arthritis is getting less frequent-results of a nationwide population-based cohort study. *Rheumatology (Oxford)*. 2017.
- Abdul Sultan A, Mallen C, Hayward R, Muller S, Whittle R, Hotston M, et al. Gout and subsequent erectile dysfunction: a population-based cohort study from England. *Arthritis Res Ther*. 2017;19(1):123.
- Zingone F, Abdul Sultan A, Crooks CJ, Tata LJ, Ciacci C, West J. The risk of community-acquired pneumonia among 9803 patients with coeliac disease compared to the general population: a cohort study. *Aliment Pharmacol Ther*. 2016;44(1):57-67.
- Zghebi SS, Steinke DT, Rutter MK, Emsley RA, Ashcroft DM. Comparative risk of major cardiovascular events associated with second-line antidiabetic treatments: a retrospective cohort study using UK primary care data linked to hospitalization and mortality records. *Diabetes Obes Metab*. 2016;18(9):916-24.
- Yang B, Petrick JL, Chen J, Hagberg KW, Sahasrabuddhe VV, Graubard BI, et al. Associations of NSAID and paracetamol use with risk of primary liver cancer in the Clinical Practice Research Datalink. *Cancer Epidemiol*. 2016;43:105-11.
- Yang B, Hagberg KW, Chen J, Sahasrabuddhe VV, Graubard BI, Jick S, et al. Associations of antibiotic use with risk of primary liver cancer in the Clinical Practice Research Datalink. *Br J Cancer*. 2016;115(1):85-9.
- Wright AK, Kontopantelis E, Emsley R, Buchan I, Sattar N, Rutter MK, et al. Life Expectancy and Cause-Specific Mortality in Type 2 Diabetes: A Population-Based Cohort Study Quantifying Relationships in Ethnic Subgroups. *Diabetes Care*. 2016:dc161616.
- Wing K, Bhaskaran K, Smeeth L, van Staa TP, Klungel OH, Reynolds RF, et al. Optimising case detection within UK electronic health records: use of multiple linked databases for detecting liver injury. *BMJ Open*. 2016;6(9):e012102.
- Windsor C, Herrett E, Smeeth L, Quint JK. No association between exacerbation frequency and stroke in patients with COPD. *Int J Chron Obstruct Pulmon Dis*. 2016;11:217-25.
- Windfuhr K, While D, Kapur N, Ashcroft DM, Kontopantelis E, Carr MJ, et al. Suicide risk linked with clinical consultation frequency, psychiatric diagnoses and psychotropic medication prescribing in a national study of primary-care patients. *Psychol Med*. 2016:1-11.
- Wilson JC, Furlano RI, Jick SS, Meier CR. Inflammatory Bowel Disease and the Risk of Autoimmune Diseases. *J Crohns Colitis*. 2016;10(2):186-93.

- Wilson JC, Furlano RI, Jick SS, Meier CR. A population-based study examining the risk of malignancy in patients diagnosed with inflammatory bowel disease. *J Gastroenterol.* 2016;51(11):1050-62.
- Willame C, Rosillon D, Zima J, Angelo M-G, Stuurman AL, Vroling H, et al. Risk of new onset autoimmune disease in 9- to 25-year-old women exposed to human papillomavirus-16/18 AS04-adjuvanted vaccine in the United Kingdom. *Hum Vaccin Immunother.* 2016:1-10.
- Willame C, Baril L, van den Bosch J, Ferreira GLC, Williams R, Rosillon D, et al. Importance of feasibility assessments before implementing non-interventional pharmacoepidemiologic studies of vaccines: lessons learned and recommendations for future studies. *Pharmacoepidemiol Drug Saf.* 2016.
- Wilke T, Mueller S, Groth A, Berg B, Fuchs A, Sikirica M, et al. Non-Persistence and Non-Adherence of Patients with Type 2 Diabetes Mellitus in Therapy with GLP-1 Receptor Agonists: A Retrospective Analysis. *Diabetes Ther.* 2016;7(1):105-24.
- Wilding J, Bailey C, Rigney U, Blak B, Beekman W, Emmas C. Glycated Hemoglobin, Body Weight and Blood Pressure in Type 2 Diabetes Patients Initiating Dapagliflozin Treatment in Primary Care: A Retrospective Study. *Diabetes Ther.* 2016.
- Wexler NS, Collett L, Wexler AR, Rawlins MD, Tabrizi SJ, Douglas I, et al. Incidence of adult Huntington's disease in the UK: a UK-based primary care study and a systematic review. *BMJ Open.* 2016;6(2):e009070.
- Weir S, Samnaliev M, Kuo T, Tierney T, Walleser Autiero S, Taylor R, et al. Healthcare resource utilisation and cost of Parkinson's disease in the united kingdom. *Value in Health.* 2016;19(7):A428-A9.
- Wei L, MacDonald T, Shimi S. Association between prior appendectomy and/or tonsillectomy in women and subsequent pregnancy rate: a cohort study. *Fertil Steril.* 2016;106(5):1150-6.
- Watson L, Das R, Farquhar R, Langerman H, Barnett AH. Consequences of delaying treatment intensification in type 2 diabetes: evidence from a UK database. *Current Medical Research and Opinion.* 2016;32(9):1465-75.
- Walker VM, Davies NM, Jones T, Kehoe PG, Martin RM. Can commonly prescribed drugs be repurposed for the prevention or treatment of Alzheimer's and other neurodegenerative diseases? Protocol for an observational cohort study in the UK Clinical Practice Research Datalink. *BMJ Open.* 2016;6(12):e012044.
- Walker AJ, West J, Card TR, Crooks C, Kirwan CC, Grainge MJ. When are breast cancer patients at highest risk of venous thromboembolism? A cohort study using English health care data. *Blood.* 2016;127(7):849-57; quiz 953.
- Walker AJ, Baldwin DR, Card TR, Powell HA, Hubbard RB, Grainge MJ. Risk of venous thromboembolism in people with lung cancer: a cohort study using linked UK healthcare data. *Br J Cancer.* 2016;115(1):115-21.
- Villa G, Patel J, Qian Y, Lira A, Taylor B, Danese M. Cardiovascular event rates in patients with high cardiovascular risk in the United Kingdom. *Value in Health.* 2016;19(7):A636.
- Vezyridis P, Timmons S. Evolution of primary care databases in UK: a scientometric analysis of research output. *BMJ Open.* 2016;6(10):e012785.
- Verhaegh BPM, de Vries F, Masclee AaM, Keshavarzian A, de Boer A, Souverein PC, et al. High risk of drug-induced microscopic colitis with concomitant use of NSAIDs and proton pump inhibitors. *Aliment Pharmacol Ther.* 2016;43(9):1004-13.
- Venkatesan S, Okoli GN, Mozid AM, Pickworth TWH, Grocott MPW, Sanders RD, et al. Effects of five

preoperative cardiovascular drugs on mortality after coronary artery bypass surgery: A retrospective analysis of an observational study of 16,192 patients. *Eur J Anaesthesiol.* 2016;33(1):49-57.

- van der Velde RY, Wyers CE, Teesselink E, Geusens PPM, van den Bergh JPW, de Vries F, et al. Trends in oral anti-osteoporosis drug prescription in the United Kingdom between 1990 and 2012: Variation by age, sex, geographic location and ethnicity. *Bone.* 2016;94:50-5.
- van der Velde RY, Wyers CE, Curtis EM, Geusens PPM, van den Bergh JPW, de Vries F, et al. Secular trends in fracture incidence in the UK between 1990 and 2012. *Osteoporos Int.* 2016;27(11):3197-206.
- van Dalem J, Brouwers MCGJ, Stehouwer CDA, Krings A, Leufkens HGM, Driessen JHM, et al. Risk of hypoglycaemia in users of sulphonylureas compared with metformin in relation to renal function and sulphonylurea metabolite group: population based cohort study. *BMJ.* 2016;354:i3625.
- van Brunt K, Curtis B, Ivanyi T, Balogh E, Chalkiadaki C, MacLachlan S, et al. Basal-bolus Therapy in Patients with Type 2 Diabetes Mellitus in the UK: Patient Characteristics, Treatment Patterns and the Effect of Switching to Premixed Insulin. *Diabetes Ther.* 2016.
- Vamos EP, Pape UJ, Curcin V, Harris MJ, Valabhji J, Majeed A, et al. Effectiveness of the influenza vaccine in preventing admission to hospital and death in people with type 2 diabetes. *CMAJ.* 2016;188(14):E342-E51.
- Valovicova V, Nagy L, Brasseur P, Pitman RJ. A CPRD-HES database study of back pain in the UK: Incidence and prevalence. *Value in Health.* 2016;19(7):A579.
- Tyrrell EG, Orton E, Sayal K, Baker R, Kendrick D. Differing patterns in intentional and unintentional poisonings among young people in England, 1998-2014: a population-based cohort study. *J Public Health (Oxf).* 2016.
- Tutt L, Anderson C, Chen L-C, Thornley T. Exploring prescribing patterns to inform community pharmacy services in postbreast cancer survivorship. *International Journal of Pharmacy Practice.* 2016;24:36.
- Tuccori M, Filion KB, Yin H, Yu OH, Platt RW, Azoulay L. Pioglitazone use and risk of bladder cancer: population based cohort study. *BMJ.* 2016;352:i1541.
- Trueman D, Hancock E, Jhund P, McMurray JJ, Petrie M, Deschaseaux-Voinet C, et al. The cost-effectiveness of sacubitril/valsartan for the treatment of chronic heart failure with reduced ejection fraction in England and Scotland: Applying real-world evidence to enhance generalisability to a UK population. *Value in Health.* 2016;19(7):A655.
- Thomas SL, Walker JL, Fenty J, Atkins KE, Elliot AJ, Hughes HE, et al. Impact of the national rotavirus vaccination programme on acute gastroenteritis in England and associated costs averted. *Vaccine.* 2016.
- Tebboth A, Lee S, Scowcroft A, Bingham-Gardiner P, Spencer W, Bolodeoku J, et al. Demographic and Clinical Characteristics of Patients With Type 2 Diabetes Mellitus Initiating Dipeptidyl Peptidase 4 Inhibitors: A Retrospective Study of UK General Practice. *Clin Ther.* 2016;38(8):1825-32.e15.
- Taylor S, Taylor RJ, Lustig RL, Schuck-Paim C, Haguinet F, Webb DJ, et al. Modelling estimates of the burden of respiratory syncytial virus infection in children in the UK. *BMJ Open.* 2016;6(6):e009337.
- Tascilar K, Dell'Aniello S, Hudson M, Suissa S. Statins and Risk of Rheumatoid Arthritis: A Nested Case-Control Study. *Arthritis & Rheumatology (Hoboken, NJ).* 2016;68(11):2603-11.
- Tascilar K, Azoulay L, Dell'Aniello S, Bartels DB, Suissa S. The Use of Telmisartan and the Incidence of Cancer.

Am J Hypertens. 2016.

- Sultan AA, West J, Grainge MJ, Riley RD, Tata LJ, Stephansson O, et al. Development and validation of risk prediction model for venous thromboembolism in postpartum women: multinational cohort study. *BMJ*. 2016;i6253.
- Sultan AA, Mallen C, Hayward R, Muller S, Whittle R, Hotston M, et al. Gout and subsequent risk of incident erectile dysfunction: A population-based cohort study from the united kingdom. *Arthritis and Rheumatology*. 2016;68:2991-2.
- Suissa S, Moodie EEM, Dell'Aniello S. Prevalent new-user cohort designs for comparative drug effect studies by time-conditional propensity scores. *Pharmacoepidemiol Drug Saf*. 2016.
- Suissa S, Dellaniello S, Ernst P. Long-acting bronchodilator initiation in COPD and the risk of adverse cardio-pulmonary events: A population-based comparative safety study. *Chest*. 2016.
- Stevens SL, Stevens RJ, Hobbs FDR, Lasserson DS. Chronic renal disease is not chronic kidney disease: implications for use of the QRISK and Joint British Societies risk scores. *Fam Pract*. 2016;33(1):57-60.
- Stack A, Blak B, Johnson M, Parsons V, Maguire A, Klein AB, et al. Association of gout with risk of advanced chronic kidney disease. *Arthritis and Rheumatology*. 2016;68:4262-3.
- Springate DA, Parisi R, Kontopantelis E, Reeves D, Griffiths CEM, Ashcroft DM. Incidence, prevalence and mortality of patients with psoriasis: a UK population-based cohort study. *Br J Dermatol*. 2016.
- Spoenclin J, Meier C, Jick SS, Meier CR. Bisphosphonate therapy start may transiently increase the risk of tendon rupture in patients with glucocorticoid co-medication: a population-based observational study. *Pharmacoepidemiol Drug Saf*. 2016;25(10):1116-23.
- Spoenclin J, Meier C, Jick SS, Meier CR. Achilles or biceps tendon rupture in women and men with type 2 diabetes: A population-based case-control study. *J Diabetes Complicat*. 2016;30(5):903-9.
- Spoenclin J, Layton JB, Mundkur M, Meier C, Jick SS, Meier CR. The Risk of Achilles or Biceps Tendon Rupture in New Statin Users: A Propensity Score-Matched Sequential Cohort Study. *Drug Safety*. 2016;39(12):1229-37.
- Spoenclin J, Karatas G, Furlano RI, Jick SS, Meier CR. Rosacea in Patients with Ulcerative Colitis and Crohn's Disease: A Population-based Case-control Study. *Inflamm Bowel Dis*. 2016;22(3):680-7.
- Shin J-Y, Eberg M, Ernst P, Filion KB. Statin Potency and the Risk of Hospitalization for Community-Acquired Pneumonia. *British Journal of Clinical Pharmacology*. 2016.
- Shephard EA, Neal RD, Rose PW, Walter FM, Hamilton W. Symptoms of adult chronic and acute leukaemia before diagnosis: large primary care case-control studies using electronic records. *Br J Gen Pract*. 2016;66(644):e182-8.
- Shah A, Prieto-Alhambra D, Hawley S, Delmestri A, Lippett J, Cooper C, et al. Geographic variation in secondary fracture prevention after a hip fracture during 1999-2013: a UK study. *Osteoporos Int*. 2016.
- Seliger C, Ricci C, Meier CR, Bodmer M, Jick SS, Bogdahn U, et al. Diabetes, use of antidiabetic drugs, and the risk of glioma. *Neuro-oncology*. 2016;18(3):340-9.
- Seliger C, Meier CR, Jick SS, Uhl M, Bogdahn U, Hau P, et al. Use of cardiac glycosides and risk of glioma. *J Neurooncol*. 2016;127(2):321-8.

- Seliger C, Meier CR, Becker C, Jick SS, Bogdahn U, Hau P, et al. Use of Selective Cyclooxygenase-2 Inhibitors, Other Analgesics, and Risk of Glioma. *PLoS One*. 2016;11(2):e0149293.
- Schuerch M, Gasse C, Robinson NJ, Alvarez Y, Walls R, Mors O, et al. Impact of varying outcomes and definitions of suicidality on the associations of antiepileptic drugs and suicidality: comparisons from UK Clinical Practice Research Datalink (CPRD) and Danish national registries (DNR). *Pharmacoepidemiol Drug Saf*. 2016;25 Suppl 1:142-55.
- Schmidt SAJ, Vestergaard M, Pedersen HS, Schønheyder HC, Thomas SL, Smeeth L, et al. Partner Bereavement and Risk of Herpes Zoster: Results from Two Population-Based Case-Control Studies in Denmark and the United Kingdom. *Clin Infect Dis*. 2016.
- Rudisill C, Charlton J, Booth HP, Gulliford MC. Are healthcare costs from obesity associated with body mass index, comorbidity or depression? Cohort study using electronic health records. *Clin Obes*. 2016;6(3):225-31.
- Rothnie KJ, Müllerová H, Thomas SL, Chandan JS, Smeeth L, Hurst JR, et al. Recording of hospitalizations for acute exacerbations of COPD in UK electronic health care records. *Clin Epidemiol*. 2016;Volume 8:771-82.
- Rothnie KJ, Müllerová H, Hurst JR, Smeeth L, Davis K, Thomas SL, et al. Validation of the Recording of Acute Exacerbations of COPD in UK Primary Care Electronic Healthcare Records. *PLoS One*. 2016;11(3):e0151357.
- Root AA, Wong AYS, Ghebremichael-Weldeselassie Y, Smeeth L, Bhaskaran K, Evans SJW, et al. Evaluation of the risk of cardiovascular events with clarithromycin using both propensity score and self-controlled study designs. *British Journal of Clinical Pharmacology*. 2016;82(2):512-21.
- Robson JC, Kiran A, Maskell J, Hutchings A, Arden N, Dasgupta B, et al. Which Patients with Giant Cell Arteritis Will Develop Cardiovascular or Cerebrovascular Disease? A Clinical Practice Research Datalink Study. *J Rheumatol*. 2016;43(6):1085-92.
- Renzi C, Lyraztopoulos G, Card T, Chu TPC, Macleod U, Rachet B. Do colorectal cancer patients diagnosed as an emergency differ from non-emergency patients in their consultation patterns and symptoms? A longitudinal data-linkage study in England. *Br J Cancer*. 2016;115(7):866-75.
- Renoux C, Vahey S, Dell'Aniello S, Boivin J-F. Association of Selective Serotonin Reuptake Inhibitors With the Risk for Spontaneous Intracranial Hemorrhage. *JAMA Neurology*. 2016.
- Renoux C, Shin J-Y, Dell'Aniello S, Fergusson E, Suissa S. Prescribing trends of attention-deficit hyperactivity disorder (ADHD) medications in UK primary care, 1995-2015. *British Journal of Clinical Pharmacology*. 2016;82(3):858-68.
- Renoux C, Dell'Aniello S, Khairy P, Marras C, Bugden S, Turin TC, et al. Ventricular tachyarrhythmia and sudden cardiac death with domperidone use in Parkinson's disease. *British Journal of Clinical Pharmacology*. 2016;82(2):461-72.
- Rees F, Doherty M, Lanyon P, Davenport G, Riley RD, Zhang W, et al. Early clinical features in Systemic Lupus Erythematosus: can they be used to achieve earlier diagnosis? A risk prediction model. *Arthritis Care Res (Hoboken)*. 2016.
- Rees F, Doherty M, Grainge MJ, Lanyon P, Davenport G, Zhang W. Mortality in systemic lupus erythematosus in the United Kingdom 1999-2012. *Rheumatology (Oxford)*. 2016;55(5):854-60.
- Rees F, Doherty M, Grainge M, Lanyon P, Davenport G, Zhang W. Burden of Comorbidity in Systemic Lupus Erythematosus in the UK, 1999-2012. *Arthritis Care Res (Hoboken)*. 2016;68(6):819-27.

- Rees F, Doherty M, Grainge M, Davenport G, Lanyon P, Zhang W. The incidence and prevalence of systemic lupus erythematosus in the UK, 1999-2012. *Annals of the Rheumatic Diseases*. 2016;75(1):136-41.
- Rees DA, Jenkins-Jones S, Morgan CL. Contemporary Reproductive Outcomes for Patients With Polycystic Ovary Syndrome: A Retrospective Observational Study. *J Clin Endocrinol Metab*. 2016;101(4):1664-72.
- Rajaram S, Steffey A, Blak B, Hickman M, Christensen H, Caspard H. Uptake of childhood influenza vaccine from 2012-2013 to 2014-2015 in the UK and the implications for high-risk children: a retrospective observational cohort study. *BMJ Open*. 2016;6(8):e010625.
- Quint JK, Millett ERC, Joshi M, Navaratnam V, Thomas SL, Hurst JR, et al. Changes in the incidence, prevalence and mortality of bronchiectasis in the UK from 2004 to 2013: a population-based cohort study. *Eur Respir J*. 2016;47(1):186-93.
- Qiao Q, Morgan CL, Jenkins-Jones S, Berni E, Currie CJ, Holden SE. Healthcare resource utilisation associated with patients treated with either exenatide once weekly or basal insulin: A retrospective UK database analysis. *Value in Health*. 2016;19(7):A670-A1.
- Price SJ, Stapley SA, Shephard E, Barraclough K, Hamilton WT. Is omission of free text records a possible source of data loss and bias in Clinical Practice Research Datalink studies? A case-control study. *BMJ Open*. 2016;6(5):e011664.
- Price D, Wilson AM, Chisholm A, Rigazio A, Burden A, Thomas M, et al. Predicting frequent asthma exacerbations using blood eosinophil count and other patient data routinely available in clinical practice. *J Asthma Allergy*. 2016;9:1-12.
- Peng X, He X, Liu D, Van Brunt K, Balogh E, Idris I. The trend of high dose insulin initiated among diabetes patients in UK. *Value in Health*. 2016;19(7):A683-A4.
- Peeters PJHL, Bazelier MT, Leufkens HGM, Auvinen A, van Staa TP, de Vries F, et al. Insulin glargine use and breast cancer risk: Associations with cumulative exposure. *Acta Oncol*. 2016;55(7):851-8.
- Pearce FA, Grainge MJ, Lanyon PC, Watts RA, Hubbard RB. The incidence, prevalence and mortality of granulomatosis with polyangiitis in the UK Clinical Practice Research Datalink. *Rheumatology*. 2016:kew413.
- Paul SK, Klein K, Majeed A, Khunti K. Association of smoking and concomitant metformin use with cardiovascular events and mortality in people newly diagnosed with type 2 diabetes. *J Diabetes*. 2016;8(3):354-62.
- Parker SE, Jick SS, Werler MM. Intrauterine device use and the risk of pre-eclampsia: a case-control study. *BJOG*. 2016;123(5):788-95.
- Pang M, Schuster T, Filion KB, Eberg M, Platt RW. Targeted Maximum Likelihood Estimation for Pharmacoepidemiologic Research. *Epidemiology*. 2016;27(4):570-7.
- Oyinlola JO, Campbell J, Kousoulis AA. Is real world evidence influencing practice? A systematic review of CPRD research in NICE guidances. *BMC Health Serv Res*. 2016;16:299.
- Olsen JR, Piguet V, Gallacher J, Francis NA. Molluscum contagiosum and associations with atopic eczema in children: a retrospective longitudinal study in primary care. *Br J Gen Pract*. 2016;66(642):e53-8.
- Olier I, Springate DA, Ashcroft DM, Doran T, Reeves D, Planner C, et al. Modelling Conditions and Health Care Processes in Electronic Health Records: An Application to Severe Mental Illness with the Clinical Practice

Research Datalink. PloS One. 2016;11(2):e0146715.

- Offman J, Pesola F, Fitzgerald R, Hamilton W, Sasieni P. Incidence and progression of Barrett's oesophagus in clinical practice: A cohort analysis for planning of the Barrett's ESophagus Trial 3 (BEST3) comparing the Cytosponge-TFF3 test with usual care. *European Journal of Surgical Oncology (EJSO)*. 2016;42(11):S252-S3.
- Nyberg F, Horne L, Morlock R, Nuevo J, Storgard C, Aiyer L, et al. Comorbidity Burden in Trial-Aligned Patients with Established Gout in Germany, UK, US, and France: a Retrospective Analysis. *Adv Ther*. 2016;33(7):1180-98.
- Nielen JTH, Emans PJ, Dagnelie PC, Boonen A, Lalmohamed A, de Boer A, et al. Severity of Diabetes Mellitus and Total Hip or Knee Replacement: A Population-Based Case-Control Study. *Medicine*. 2016;95(20):e3739.
- Nielen JTH, de Vries F, Dagnelie PC, van den Bemt BJF, Emans PJ, Lalmohamed A, et al. Use of thiazolidinediones and the risk of elective hip or knee replacement: a population based case-control study. *British Journal of Clinical Pharmacology*. 2016;81(2):370-8.
- Nielen JTH, Dagnelie PC, Emans PJ, Veldhorst-Janssen N, Lalmohamed A, van Staa T-P, et al. Safety and efficacy of new oral anticoagulants and low-molecular-weight heparins compared with aspirin in patients undergoing total knee and hip replacements. *Pharmacoepidemiol Drug Saf*. 2016;25(11):1245-52.
- Navaratnam V, Millett ERC, Hurst JR, Thomas SL, Smeeth L, Hubbard RB, et al. Bronchiectasis and the risk of cardiovascular disease: a population-based study. *Thorax*. 2016.
- Nakafero G, Sanders RD, Nguyen-Van-Tam JS, Myles PR. The association between benzodiazepines and influenza-like illness-related pneumonia and mortality: a survival analysis using UK Primary Care data. *Pharmacoepidemiol Drug Saf*. 2016;25(11):1263-73.
- Nagy L, Valovicova V, Brasseur P, Pitman RJ. A CPRD-HES database study of back pain in the UK: Patient pathways. *Value in Health*. 2016;19(7):A603.
- Movahedi M, Beauchamp M-E, Abrahamowicz M, Ray DW, Michaud K, Pedro S, et al. Risk of Incident Diabetes Mellitus Associated With the Dosage and Duration of Oral Glucocorticoid Therapy in Patients With Rheumatoid Arthritis. *Arthritis & Rheumatology (Hoboken, NJ)*. 2016;68(5):1089-98.
- Moss S, Melia J, Sutton J, Mathews C, Kirby M. Prostate-specific antigen testing rates and referral patterns from general practice data in England. *Int J Clin Pract*. 2016;70(4):312-8.
- Morlock R, Chevalier P, Klein AB. Rate of hospitalization for heart failure is lower in patients with controlled gout versus uncontrolled gout. *Arthritis and Rheumatology*. 2016;68:1575-6.
- Morgan AD, Herrett E, De Stavola BL, Smeeth L, Quint JK. COPD disease severity and the risk of venous thromboembolic events: a matched case-control study. *Int J Chron Obstruct Pulmon Dis*. 2016;11:899-908.
- Morales DR, Dreischulte T, Lipworth BJ, Donnan PT, Jackson C, Guthrie B. Respiratory effect of beta-blocker eye drops in asthma: population-based study and meta-analysis of clinical trials. *British Journal of Clinical Pharmacology*. 2016;82(3):814-22.
- Moore E, Hashmi M, Sultana K, Chatzidiakou L, Jones RL, Beevers S, et al. Using the clinical practice research datalink (CPRD) to recruit participants from primary care to investigate chronic obstructive pulmonary disease (COPD) exacerbations. *Thorax*. 2016;71:A199-A200.
- Moore E, Chatzidiakou L, Jones RL, Smeeth L, Beevers S, Kelly FJ, et al. Linking e-health records, patient-

reported symptoms and environmental exposure data to characterise and model COPD exacerbations: protocol for the COPE study. *BMJ Open*. 2016;6(7):e011330.

- Moon RJ, Harvey NC, Curtis EM, de Vries F, van Staa T, Cooper C. Ethnic and geographic variations in the epidemiology of childhood fractures in the United Kingdom. *Bone*. 2016;85:9-14.
- Millett ERC, Quint JK, De Stavola BL, Smeeth L, Thomas SL. Improved incidence estimates from linked vs. stand-alone electronic health records. *J Clin Epidemiol*. 2016;75:66-9.
- Merinopoulou E, Raluy-Callado M, Ramagopalan S, MacLachlan S, Khalid JM. COPD exacerbations by disease severity in England. *Int J Chron Obstruct Pulmon Dis*. 2016;11:697-709.
- Menamin ÚCM, Cardwell CR, Hughes CM, Murray LM. Metformin use and survival from lung cancer: A population-based cohort study. *Lung Cancer*. 2016;94:35-9.
- Mehta HB, Mehta V, Goodwin JS. Association of Hypoglycemia With Subsequent Dementia in Older Patients With Type 2 Diabetes Mellitus. *J Gerontol A Biol Sci Med Sci*. 2016.
- Mehta HB, Mehta V, Girman CJ, Adhikari D, Johnson ML. Regression coefficient-based scoring system should be used to assign weights to the risk index. *J Clin Epidemiol*. 2016.
- McGlynn KA, Hagberg K, Chen J, Braunlin M, Graubard BI, Suneja N, et al. Menopausal hormone therapy use and risk of primary liver cancer in the clinical practice research datalink. *Int J Cancer*. 2016;138(9):2146-53.
- Mc Menamin ÚC, Murray LJ, Hughes CM, Cardwell CR. Metformin use and survival after colorectal cancer: A population-based cohort study. *Int J Cancer*. 2016;138(2):369-79.
- Mazzarella L, Botteri E, Matthews A, Disalvatore D, Bagnardi V, Breccia M, et al. Obesity as a risk factor for acute promyelocytic leukemia. Results from population and case-control studies across Western Countries and correlation with gene expression in the TCGA. *Blood*. 2016;128(22).
- Matthews A, Langan SM, Douglas IJ, Smeeth L, Bhaskaran K. Phosphodiesterase Type 5 Inhibitors and Risk of Malignant Melanoma: Matched Cohort Study Using Primary Care Data from the UK Clinical Practice Research Datalink. *PLoS Med*. 2016;13(6):e1002037.
- Matthews A, Herrett E, Gasparrini A, Van Staa T, Goldacre B, Smeeth L, et al. Impact of statin related media coverage on use of statins: interrupted time series analysis with UK primary care data. *BMJ*. 2016;353:i3283.
- Martinez C, Suissa S, Rietbrock S, Katholing A, Freedman B, Cohen AT, et al. Testosterone treatment and risk of venous thromboembolism: population based case-control study. *BMJ*. 2016;355:i5968.
- Martinez C, Katholing A, Wallenhorst C, Freedman SB. Therapy persistence in newly diagnosed non-valvular atrial fibrillation treated with warfarin or NOAC. A cohort study. *Thromb Haemost*. 2016;115(1):31-9.
- Martinez C, Katholing A, Folkerts K, Cohen AT. Risk of recurrent venous thromboembolism after discontinuation of vitamin K antagonist treatment: a nested case-control study. *J Thromb Haemost*. 2016;14(7):1374-83.
- Mars B, Heron J, Kessler D, Davies NM, Martin RM, Thomas KH, et al. Influences on antidepressant prescribing trends in the UK: 1995–2011. *Social Psychiatry and Psychiatric Epidemiology*. 2016.
- Mansfield KE, Nitsch D, Smeeth L, Bhaskaran K, Tomlinson LA. Prescription of renin-angiotensin system blockers and risk of acute kidney injury: a population-based cohort study. *BMJ Open*. 2016;6(12):e012690.

- Mandilaras V, Bouganim N, Yin H, Asselah J, Azoulay L. The use of drugs acting on the renin–angiotensin system and the incidence of pancreatic cancer. *Br J Cancer*. 2016.
- Mahmoudpour SH, Baranova EV, Souverein PC, Asselbergs FW, de Boer A, Maitland-van der Zee AH, et al. Determinants of angiotensin-converting enzyme inhibitor (ACEI) intolerance and angioedema in the UK Clinical Practice Research Datalink. *British Journal of Clinical Pharmacology*. 2016.
- Mackenzie IS, Morant SV, Wei L, Thompson AM, MacDonald TM. Spironolactone use and risk of incident cancers: a retrospective, matched cohort study. *British Journal of Clinical Pharmacology*. 2016.
- Lum KJ, Newcomb CW, Roy JA, Carbonari DM, Saine ME, Cardillo S, et al. Evaluation of methods to estimate missing days' supply within pharmacy data of the Clinical Practice Research Datalink (CPRD) and The Health Improvement Network (THIN). *Eur J Clin Pharmacol*. 2016.
- Lim AWW, Hamilton W, Hollingworth A, Stapley S, Sasieni P. Performance characteristics of visualising the cervix in symptomatic young females: a review of primary care records in females with and without cervical cancer. *Br J Gen Pract*. 2016;66(644):e189-92.
- Lim AW, Landy R, Castanon A, Hollingworth A, Hamilton W, Dudding N, et al. Cytology in the diagnosis of cervical cancer in symptomatic young women: a retrospective review. *Br J Gen Pract*. 2016.
- Lian Y, Yin H, Pollak MN, Carrier S, Platt RW, Suissa S, et al. Phosphodiesterase Type 5 Inhibitors and the Risk of Melanoma Skin Cancer. *European Urology*. 2016.
- Larivée N, Suissa S, Eberg M, Joseph L, Eisenberg MJ, Abenhaim HA, et al. Drospirenone-containing combined oral contraceptives and the risk of arterial thrombosis: a population-based nested case-control study. *BJOG*. 2016.
- Lane A, Metcalfe C, Young GJ, Peters TJ, Blazeby J, Avery KNL, et al. Patient-reported outcomes in the ProtecT randomized trial of clinically localized prostate cancer treatments: study design, and baseline urinary, bowel and sexual function and quality of life. *BJU International*. 2016;118(6):869-79.
- Landis SH, Wurst K, Le HV, Bonar K, Punekar YS. Can Assessment of Disease Burden Prior to Changes in Initial COPD Maintenance Treatment Provide Insight into Remaining Unmet Needs? A Retrospective Database Study in UK Primary Care. *COPD*. 2016:1-6.
- Kuo C-F, Grainge MJ, Mallen C, Zhang W, Doherty M. Impact of gout on the risk of atrial fibrillation. *Rheumatology (Oxford)*. 2016;55(4):721-8.
- Kuiper JG, Van Herk-Sukel MP, Castellsague J, Pottgard A, Anveden Berglind I, Dedman D, et al. Utilization of topical tacrolimus and topical pimecrolimus in Europe: Results from the joint European longitudinal lymphoma and skin cancer evaluation (Joelle) study. *Value in Health*. 2016;19(7):A573.
- Krishnamoorthi R, Borah B, Heien H, Das A, Chak A, Iyer PG. Rates and predictors of progression to esophageal carcinoma in a large population-based Barrett's esophagus cohort. *Gastrointest Endosc*. 2016;84(1):40-6.e7.
- Korhonen P, Heintjes EM, Williams R, Hoti F, Christopher S, Majak M, et al. Pioglitazone use and risk of bladder cancer in patients with type 2 diabetes: retrospective cohort study using datasets from four European countries. *BMJ*. 2016;354:i3903.
- Kontopantelis E, Springate DA, Ashcroft DM, Valderas JM, van der Veer SN, Reeves D, et al. Associations between exemption and survival outcomes in the UK's primary care pay-for-performance programme: a

retrospective cohort study. *BMJ Qual Saf.* 2016;25(9):657-70.

- Knapen LM, van Dalem J, Keulemans YC, van Erp NP, Bazelier MT, De Bruin ML, et al. Use of incretin agents and risk of pancreatic cancer: a population-based cohort study. *Diabetes Obes Metab.* 2016;18(3):258-65.
- Knapen LM, de Jong RGPJ, Driessen JHM, Keulemans YC, van Erp NP, De Bruin ML, et al. The use of incretin agents and risk of acute and chronic pancreatitis: a population-based cohort study. *Diabetes, Obesity and Metabolism.* 2016.
- Klop C, van Staa TP, Cooper C, Harvey NC, de Vries F. The epidemiology of mortality after fracture in England: variation by age, sex, time, geographic location, and ethnicity. *Osteoporos Int.* 2016.
- Klop C, de Vries F, Bijlsma JWJ, Leufkens HGM, Welsing PMJ. Predicting the 10-year risk of hip and major osteoporotic fracture in rheumatoid arthritis and in the general population: an independent validation and update of UK FRAX without bone mineral density. *Annals of the Rheumatic Diseases.* 2016;75(12):2095-100.
- Klil-Drori AJ, Yin H, Tagalakis V, Aprikian A, Azoulay L. Androgen Deprivation Therapy for Prostate Cancer and the Risk of Venous Thromboembolism. *European Urology.* 2016;70(1):56-61.
- Klil-Drori AJ, Tascilar K, Yin H, Aprikian A, Bitton A, Azoulay L. Androgen Deprivation Therapy and the Incidence of Inflammatory Bowel Disease in Patients With Prostate Cancer. *Am J Epidemiol.* 2016;184(1):15-22.
- Kingswood JC, Nasuti P, Patel K, Myland M, Siva V, Gray E. The economic burden of tuberous sclerosis complex in UK patients with renal manifestations: a retrospective cohort study in the clinical practice research datalink (CPRD). *J Med Econ.* 2016;19(12):1116-26.
- Kingswood JC, Crawford P, Johnson SR, Sampson JR, Shepherd C, Demuth D, et al. The economic burden of tuberous sclerosis complex in the UK: A retrospective cohort study in the Clinical Practice Research Datalink. *J Med Econ.* 2016;19(11):1087-98.
- Kingswood C, Bolton P, Crawford P, Harland C, Johnson SR, Sampson JR, et al. The clinical profile of tuberous sclerosis complex (TSC) in the United Kingdom: A retrospective cohort study in the Clinical Practice Research Datalink (CPRD). *Eur J Paediatr Neurol.* 2016;20(2):296-308.
- Khunti K, Nikolajsen A, Thorsted BL, Andersen M, Davies MJ, Paul SK. Clinical inertia with regard to intensifying therapy in people with type 2 diabetes treated with basal insulin. *Diabetes Obes Metab.* 2016;18(4):401-9.
- Khosrow-Khavar F, Rej S, Yin H, Aprikian A, Azoulay L. Androgen Deprivation Therapy and the Risk of Dementia in Patients With Prostate Cancer. *Journal of Clinical Oncology.* 2016;JCO-2016.
- Karystianis G, Sheppard T, Dixon WG, Nenadic G. Modelling and extraction of variability in free-text medication prescriptions from an anonymised primary care electronic medical record research database. *BMC Med Inform Decis Mak.* 2016;16:18.
- Karasneh RA, Murray LJ, Hughes CM, Cardwell CR. Digoxin use after diagnosis of prostate cancer and survival: a population-based cohort study. *Pharmacoepidemiol Drug Saf.* 2016;25(9):1099-103.
- Juszczak D, Charlton J, McDermott L, Soames J, Sultana K, Ashworth M, et al. Electronically delivered, multicomponent intervention to reduce unnecessary antibiotic prescribing for respiratory infections in primary care: a cluster randomised trial using electronic health records-REDUCE Trial study original protocol. *BMJ Open.* 2016;6(8):e010892.

- Julious SA, Horspool MJ, Davis S, Bradburn M, Norman P, Shephard N, et al. PLEASANT: Preventing and Lessening Exacerbations of Asthma in School-age children Associated with a New Term – a cluster randomised controlled trial and economic evaluation. *Health Technology Assessment*. 2016;20(93):1-154.
- Judge A, Javaid MK, Leal J, Hawley S, Drew S, Sheard S, et al. Models of care for the delivery of secondary fracture prevention after hip fracture: a health service cost, clinical outcomes and cost-effectiveness study within a region of England. Southampton (UK): NIHR Journals Library; 2016 2016.
- Joshi R, Venkatesan S, Myles PR. A UK general practice population cohort study investigating the association between lipid lowering drugs and 30-day mortality following medically attended acute respiratory illness. *PeerJ*. 2016;4:e1902.
- Joseph RM, Movahedi M, Dixon WG, Symmons DPM. Smoking-Related Mortality in Patients With Early Rheumatoid Arthritis: A Retrospective Cohort Study Using the Clinical Practice Research Datalink. *Arthritis Care Res (Hoboken)*. 2016;68(11):1598-606.
- Johnson ME, Lefèvre C, Collings S-L, Evans D, Kloss S, Ridha E, et al. Early real-world evidence of persistence on oral anticoagulants for stroke prevention in non-valvular atrial fibrillation: a cohort study in UK primary care. *BMJ Open*. 2016;6(9):e011471.
- Jameson K, D'Oca K, Leigh P, Murray-Thomas T. Adherence to NICE guidance on glucagon-like peptide-1 receptor agonists among patients with type 2 diabetes mellitus: an evaluation using the Clinical Practice Research Datalink. *Current Medical Research and Opinion*. 2016;32(1):49-60.
- Iwagami M, Mansfield K, Quint J, Nitsch D, Tomlinson L. Diagnosis of acute kidney injury and its association with in-hospital mortality in patients with infective exacerbations of bronchiectasis: cohort study from a UK nationwide database. *BMC Pulm Med*. 2016;16:14.
- Imfeld P, Toovey S, Jick SS, Meier CR. Influenza infections and risk of Alzheimer's disease. *Brain Behav Immun*. 2016;57:187-92.
- Humes DJ, Walker AJ, Hunt BJ, Sultan AA, Ludvigsson JF, West J. Risk of symptomatic venous thromboembolism following emergency appendectomy in adults. *Br J Surg*. 2016;103(4):443-50.
- Hosking FJ, Carey IM, Shah SM, Harris T, DeWilde S, Beighton C, et al. Mortality Among Adults With Intellectual Disability in England: Comparisons With the General Population. *Am J Public Health*. 2016;106(8):1483-90.
- Hong J-L, McNeill AM, He J, Chen Y, Brodovicz KG. Identification of impaired fasting glucose, healthcare utilization and progression to diabetes in the UK using the Clinical Practice Research Datalink (CPRD). *Pharmacoepidemiol Drug Saf*. 2016.
- Hollingworth W, Biswas M, Maishman RL, Dayer MJ, McDonagh T, Purdy S, et al. The healthcare costs of heart failure during the last five years of life: A retrospective cohort study. *Int J Cardiol*. 2016;224:132-8.
- Holden SE, Jenkins-Jones S, Currie CJ. Association between Insulin Monotherapy versus Insulin plus Metformin and the Risk of All-Cause Mortality and Other Serious Outcomes: A Retrospective Cohort Study. *PloS One*. 2016;11(5):e0153594.
- Hobbs FDR, Bankhead C, Mukhtar T, Stevens S, Perera-Salazar R, Holt T, et al. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007-14. *Lancet*. 2016;387(10035):2323-30.

- Hicks BM, Yin H, Yu OHY, Pollak MN, Platt RW, Azoulay L. Glucagon-like peptide-1 analogues and risk of breast cancer in women with type 2 diabetes: population based cohort study using the UK Clinical Practice Research Datalink. *BMJ*. 2016;355:i5340.
- Hicks BM, Yin H, Bladou F, Ernst P, Azoulay L. Androgen deprivation therapy for prostate cancer and the risk of hospitalisation for community-acquired pneumonia. *Thorax*. 2016:thoraxjnl-2016-209512.
- Hesdorffer DC, Ishihara L, Webb DJ, Mynepalli L, Galwey NW, Hauser WA. Occurrence and Recurrence of Attempted Suicide Among People With Epilepsy. *JAMA Psychiatry*. 2016;73(1):80-6.
- Herrett E, Williamson E, van Staa T, Ranopa M, Free C, Chadborn T, et al. Text messaging reminders for influenza vaccine in primary care: a cluster randomised controlled trial (TXT4FLUJAB). *BMJ Open*. 2016;6(2):e010069.
- Hawley S, Leal J, Delmestri A, Prieto-Alhambra D, Arden NK, Cooper C, et al. Anti-Osteoporosis Medication Prescriptions and Incidence of Subsequent Fracture Among Primary Hip Fracture Patients in England and Wales: An Interrupted Time-Series Analysis. *J Bone Miner Res*. 2016.
- Hawley S, Cordtz R, Dreyer L, Edwards CJ, Arden NK, Delmestri A, et al. The impact of biologic therapy introduction on hip and knee replacement among rheumatoid arthritis patients: An interrupted time series analysis using the clinical practice research datalink. *Arthritis and Rheumatology*. 2016;68:3497-9.
- Hassan L, Senior J, Webb RT, Frisher M, Tully MP, While D, et al. Prevalence and appropriateness of psychotropic medication prescribing in a nationally representative cross-sectional survey of male and female prisoners in England. *BMC Psychiatry*. 2016;16(1):346.
- Happich M, Kirson NY, Desai U, King S, Birnbaum HG, Reed C, et al. Excess Costs Associated with Possible Misdiagnosis of Alzheimer's Disease Among Patients with Vascular Dementia in a UK CPRD Population. *J Alzheimers Dis*. 2016;53(1):171-83.
- Hancock E, Trueman D, Jhund P, McMurray JJ, Petrie M, Deschaseaux-Voinet C, et al. The cost-effectiveness of sacubitril/valsartan for the treatment of chronic heart failure with reduced ejection fraction: The use of model calibration to improve generalisability to a UK CHF population. *Value in Health*. 2016;19(7):A655-A6.
- Hancock E, Trueman D, Alexopoulos ST, Gielen V. The cost-effectiveness of sacubitril/valsartan for the treatment of chronic heart failure with reduced ejection fraction in alternative age groups. *Value in Health*. 2016;19(7):A655.
- Hagberg KW, Sahasrabudde VV, McGlynn KA, Jick SS. Does Angiotensin-Converting Enzyme Inhibitor and β -Blocker Use Reduce the Risk of Primary Liver Cancer? A Case-Control Study Using the U.K. Clinical Practice Research Datalink. *Pharmacotherapy*. 2016;36(2):187-95.
- Hagberg KW, Li L, Peng M, Shah K, Paris M, Jick S. Incidence rates of suicidal behaviors and treated depression in patients with and without psoriatic arthritis using the Clinical Practice Research Datalink. *Mod Rheumatol*. 2016;26(5):774-9.
- Hagberg KW, Li L, Peng M, Paris M, Shah K, Jick SS. Rates of Cancers and Opportunistic Infections in Patients With Psoriatic Arthritis Compared With Patients Without Psoriatic Arthritis. *J Clin Rheumatol*. 2016;22(5):241-7.
- Hagberg KW, Divan HA, Persson R, Nickel JC, Jick SS. Risk of erectile dysfunction associated with use of 5- α reductase inhibitors for benign prostatic hyperplasia or alopecia: population based studies using the Clinical

Practice Research Datalink. *BMJ*. 2016;354:i4823.

- Gulliford MC, Moore MV, Little P, Hay AD, Fox R, Prevost AT, et al. Safety of reduced antibiotic prescribing for self limiting respiratory tract infections in primary care: cohort study using electronic health records. *BMJ*. 2016;354:i3410.
- Gulliford MC, Booth HP, Reddy M, Charlton J, Fildes A, Prevost AT, et al. Effect of Contemporary Bariatric Surgical Procedures on Type 2 Diabetes Remission. A Population-Based Matched Cohort Study. *Obes Surg*. 2016;26(10):2308-15.
- Grimaldi-Bensouda L, Klungel O, Kurz X, de Groot MCH, Maciel Afonso AS, de Bruin ML, et al. Calcium channel blockers and cancer: a risk analysis using the UK Clinical Practice Research Datalink (CPRD). *BMJ Open*. 2016;6(1):e009147.
- Green AL, Fundament T. Deep brain stimulation compared to device-aided infusion therapies for parkinson's disease with early motor complications: A UK cost-effectiveness analysis. *Value in Health*. 2016;19(7):A697.
- Granerod J, Davies NWS, Ramanuj PP, Easton A, Brown DWG, Thomas SL. Increased rates of sequelae post-encephalitis in individuals attending primary care practices in the United Kingdom: a population-based retrospective cohort study. *J Neurol*. 2016.
- Glover G, Williams R, Heslop P, Oyinola J, Grey J. Mortality in people with intellectual disabilities in England. *J Intellect Disabil Res*. 2016.
- Gielen V, Gondos A, Alexopoulos S. Real-world hospitalisation and mortality of incident and prevalent patients with heart failure due to left ventricular systolic dysfunction in England. *Value in Health*. 2016;19(7):A640.
- Getz KD, Anderka MT, Werler MM, Jick SS. Maternal Pre-pregnancy Body Mass Index and Autism Spectrum Disorder among Offspring: A Population-Based Case-Control Study. *Paediatr Perinat Epidemiol*. 2016;30(5):479-87.
- Gamble J-M, Thomas JM, Twells LK, Midodzi WK, Majumdar SR. Comparative effectiveness of incretin-based therapies and the risk of death and cardiovascular events in 38,233 metformin monotherapy users. *Medicine*. 2016;95(26):e3995.
- Gamble J-M, Chibrikov E, Twells LK, Midodzi WK, Young SW, MacDonald D, et al. Association of insulin dosage with mortality or major adverse cardiovascular events: a retrospective cohort study. *The Lancet Diabetes & Endocrinology*. 2016.
- Gallagher AM, Williams T, Leufkens HGM, de Vries F. The Impact of the Choice of Data Source in Record Linkage Studies Estimating Mortality in Venous Thromboembolism. *PloS One*. 2016;11(2):e0148349.
- Fundament T, Eldridge PR, Green AL, Whone AL, Taylor RS, Williams AC, et al. Deep Brain Stimulation for Parkinson's Disease with Early Motor Complications: A UK Cost-Effectiveness Analysis. *PloS One*. 2016;11(7):e0159340.
- Frey N, Hügler T, Jick SS, Meier CR, Spöndlin J. Type II diabetes mellitus and incident osteoarthritis of the hand: a population-based case-control analysis. *Osteoarthritis and Cartilage*. 2016;24(9):1535-40.
- Freedman B, Martinez C, Katholing A, Rietbrock S. Residual Risk of Stroke and Death in Anticoagulant-Treated Patients With Atrial Fibrillation. *JAMA Cardiol*. 2016;1(3):366-8.

- Francis NA, Hood K, Lyons R, Butler CC. Understanding flucloxacillin prescribing trends and treatment non-response in UK primary care: a Clinical Practice Research Datalink (CPRD) study. *J Antimicrob Chemother.* 2016;71(7):2037-46.
- Forster AS, Burgess C, Dodhia H, Fuller F, Miller J, McDermott L, et al. Do health checks improve risk factor detection in primary care? Matched cohort study using electronic health records. *J Public Health (Oxf).* 2016;38(3):552-9.
- Forbes HJ, Bhaskaran K, Thomas SL, Smeeth L, Clayton T, Mansfield K, et al. Quantification of risk factors for postherpetic neuralgia in herpes zoster patients: A cohort study. *Neurology.* 2016;87(1):94-102.
- Fleming DM, Taylor RJ, Haguinet F, Schuck-Paim C, Logie J, Webb DJ, et al. Influenza-attributable burden in United Kingdom primary care. *Epidemiol Infect.* 2016;144(3):537-47.
- Flahavan EM, Jones ME, Mitchell L, Swain JL, Martinez JM, Ahn AH. Cluster headache treatment patterns and analgesic use in the clinical practice research datalink (CPRD). *Cephalalgia.* 2016;36:114-5.
- Fillion KB, Azoulay L, Platt RW, Dahl M, Dormuth CR, Clemens KK, et al. A Multicenter Observational Study of Incretin-based Drugs and Heart Failure. *N Engl J Med.* 2016;374(12):1145-54.
- Ferreira G, Lopman BA, Tam C, Harris J, Bollaerts K, Cattaert T, et al. Burden of medically attended acute gastroenteritis in England: A cohort study in CPRD. *Value in Health.* 2016;19(7):A409.
- Fernandes G, Alemayehu B, Liu J, Paul E, Das R. A UK observational cohort study of insulin treatment patterns and characterisation of insulin users in type 1 and type 2 diabetes mellitus. *Value in Health.* 2016;19(7):A684.
- Farmer AJ, Rodgers LR, Lonergan M, Shields B, Weedon MN, Donnelly L, et al. Adherence to Oral Glucose-Lowering Therapies and Associations With 1-Year HbA1c: A Retrospective Cohort Analysis in a Large Primary Care Database. *Diabetes Care.* 2016;39(2):258-63.
- Fairhurst C, Martin F, Watt I, Doran T, Bland M, Brackenbury WJ. Sodium channel-inhibiting drugs and cancer survival: protocol for a cohort study using the CPRD primary care database. *BMJ Open.* 2016;6(9):e011661.
- Faillie J-L, Yu OH, Yin H, Hillaire-Buys D, Barkun A, Azoulay L. Association of Bile Duct and Gallbladder Diseases With the Use of Incretin-Based Drugs in Patients With Type 2 Diabetes Mellitus. *JAMA Intern Med.* 2016;176(10):1474-81.
- Essers I, Stolwijk C, Boonen A, De Bruin ML, Bazelier MT, de Vries F, et al. Ankylosing spondylitis and risk of ischaemic heart disease: a population-based cohort study. *Annals of the Rheumatic Diseases.* 2016;75(1):203-9.
- Emdin CA, Rothwell PM, Salimi-Khorshidi G, Kiran A, Conrad N, Callender T, et al. Blood Pressure and Risk of Vascular Dementia: Evidence From a Primary Care Registry and a Cohort Study of Transient Ischemic Attack and Stroke. *Stroke.* 2016;47(6):1429-35.
- Eastwood SV, Mathur R, Atkinson M, Brophy S, Sudlow C, Flaig R, et al. Algorithms for the Capture and Adjudication of Prevalent and Incident Diabetes in UK Biobank. *PloS One.* 2016;11(9):e0162388.
- Driessen JHM, van den Bergh JPW, van Onzenoort HAW, Henry RMA, Leufkens HGM, de Vries F. Long-term use of dipeptidyl peptidase-4 inhibitors and risk of fracture; a retrospective population-based cohort study. *Diabetes, Obesity and Metabolism.* 2016.
- Drahos J, Li L, Jick SS, Cook MB. Metabolic syndrome in relation to Barrett's esophagus and esophageal

adenocarcinoma: Results from a large population-based case-control study in the Clinical Practice Research Datalink. *Cancer Epidemiol.* 2016;42:9-14.

- Doyle M, While D, Mok PLH, Windfuhr K, Ashcroft DM, Kontopantelis E, et al. Suicide risk in primary care patients diagnosed with a personality disorder: a nested case control study. *BMC Fam Pract.* 2016;17:106.
- Dos Santos G, Haguinet F, Cohet C, Webb D, Logie J, Ferreira GLC, et al. Risk of solid organ transplant rejection following vaccination with seasonal trivalent inactivated influenza vaccines in England: A self-controlled case-series. *Vaccine.* 2016;34(31):3598-606.
- DiSantostefano RL, Boudiaf N, Stempel DA, Barnes NC, Greening AP. The frequency of, and adherence to, single maintenance and reliever therapy instructions in asthma: a descriptive analysis. *NPJ Prim Care Respir Med.* 2016;26:16038.
- DiBello JR, Ioannou C, Rees J, Challacombe B, Maskell J, Choudhury N, et al. Prevalence of metabolic syndrome and its components among men with and without clinical benign prostatic hyperplasia: a large, cross-sectional, UK epidemiological study. *BJU international.* 2016;117(5):801-8.
- Delgado J, Masoli JAH, Bowman K, Strain WD, Kuchel GA, Walters K, et al. Outcomes of Treated Hypertension at Age 80 and Older: Cohort Analysis of 79,376 Individuals. *Journal of the American Geriatrics Society.* 2016.
- de Jong RGPJ, Gallagher AM, Herrett E, Masclee AAM, Janssen-Heijnen MLG, de Vries F. Comparability of the age and sex distribution of the UK Clinical Practice Research Datalink and the total Dutch population. *Pharmacoepidemiol Drug Saf.* 2016.
- Datta-Nemdharry P, Thomson A, Beynon J, Donegan K. Patterns of anti-diabetic medication use in patients with type 2 diabetes mellitus in England and Wales. *Pharmacoepidemiol Drug Saf.* 2016.
- Datta-Nemdharry P, Thomson A, Beynon J. Opportunities and Challenges in Developing a Cohort of Patients with Type 2 Diabetes Mellitus Using Electronic Primary Care Data. *PLOS ONE.* 2016;11(11):e0162236.
- D'Angelo S, Coggon D, Harris EC, Linaker C, Sayer AA, Gale CR, et al. Job dissatisfaction and the older worker: baseline findings from the Health and Employment After Fifty study. *Occup Environ Med.* 2016;73(8):512-9.
- Danese MD, Gleeson M, Kutikova L, Griffiths RI, Azough A, Khunti K, et al. Estimating the economic burden of cardiovascular events in patients receiving lipid-modifying therapy in the UK. *BMJ Open.* 2016;6(8):e011805.
- Danese M, Gleeson M, Kutikova L, Griffiths R, Azough A. For estimating costs in patients experiencing cardiovascular events using utilization data from the United Kingdom. *Value in Health.* 2016;19(7):A645-A6.
- Curtis EM, van der Velde R, Moon RJ, van den Bergh JPW, Geusens P, de Vries F, et al. Epidemiology of fractures in the United Kingdom 1988-2012: Variation with age, sex, geography, ethnicity and socioeconomic status. *Bone.* 2016;87:19-26.
- Crooks CJ, Card TR, West J. The Use of a Bayesian Hierarchy to Develop and Validate a Co-Morbidity Score to Predict Mortality for Linked Primary and Secondary Care Data from the NHS in England. *PloS One.* 2016;11(10):e0165507.
- Crane MM, Webb DJ, Watson E, Cunliffe T, English J. Hand Eczema and Steroid-Refractory Chronic Hand Eczema in General Practice: prevalence, and initial treatment. *Br J Dermatol.* 2016.
- Costello R, Winthrop KL, Pye SR, Brown B, Dixon WG. Influenza and Pneumococcal Vaccination Uptake in Patients with Rheumatoid Arthritis Treated with Immunosuppressive Therapy in the UK: A Retrospective

Cohort Study Using Data from the Clinical Practice Research Datalink. *PLoS One*. 2016;11(4):e0153848.

- Cope AL, Chestnutt IG, Wood F, Francis NA. Dental consultations in UK general practice and antibiotic prescribing rates: a retrospective cohort study. *Br J Gen Pract*. 2016;66(646):e329-36.
- Cohet C, Haguinet F, Dos Santos G, Webb D, Logie J, Lc Ferreira G, et al. Effect of the adjuvanted (AS03) A/H1N1 2009 pandemic influenza vaccine on the risk of rejection in solid organ transplant recipients in England: a self-controlled case series. *BMJ Open*. 2016;6(1):e009264.
- Cohen AT, Katholing A, Rietbrock S, Bamber L, Martinez C. Epidemiology of first and recurrent venous thromboembolism in patients with active cancer. A population-based cohort study. *Thromb Haemost*. 2016;117(1).
- Clarke AT, Johnson PCD, Hall GC, Ford I, Mills PR. High Dose Atorvastatin Associated with Increased Risk of Significant Hepatotoxicity in Comparison to Simvastatin in UK GPRD Cohort. *PLoS One*. 2016;11(3):e0151587.
- Chui CSL, Chan EW, Wong AYS, Root A, Douglas IJ, Wong ICK. Association between oral fluoroquinolones and seizures: A self-controlled case series study. *Neurology*. 2016;86(18):1708-15.
- Chu TPC, Shah A, Walker D, Coleman MP. Where are the opportunities for an earlier diagnosis of primary intracranial tumours in children and young adults? *European Journal of Paediatric Neurology*. 2016.
- Chu TPC, Moran GW, Card TR. The pattern of underlying cause of death in patients with inflammatory bowel disease in England: a record linkage study. *J Crohns Colitis*. 2016.
- Charlton RA, Pierini A, Klungsøyr K, Neville AJ, Jordan S, de Jong-van den Berg LTW, et al. Asthma medication prescribing before, during and after pregnancy: a study in seven European regions. *BMJ Open*. 2016;6(1):e009237.
- Chang KC-M, Lee JT, Vamos EP, Soljak M, Johnston D, Khunti K, et al. Impact of the National Health Service Health Check on cardiovascular disease risk: a difference-in-differences matching analysis. *CMAJ*. 2016;188(10):E228-38.
- Carr MJ, Ashcroft DM, Kontopantelis E, While D, Awenat Y, Cooper J, et al. Clinical management following self-harm in a UK-wide primary care cohort. *J Affect Disord*. 2016;197:182-8.
- Carr MJ, Ashcroft DM, Kontopantelis E, Awenat Y, Cooper J, Chew-Graham C, et al. The epidemiology of self-harm in a UK-wide primary care patient cohort, 2001-2013. *BMC Psychiatry*. 2016;16:53.
- Carey IM, Shah SM, Hosking FJ, DeWilde S, Harris T, Beighton C, et al. Health characteristics and consultation patterns of people with intellectual disability: a cross-sectional database study in English general practice. *Br J Gen Pract*. 2016;66(645):e264-70.
- Carey IM, Hosking FJ, Harris T, DeWilde S, Beighton C, Shah SM, et al. Do health checks for adults with intellectual disability reduce emergency hospital admissions? Evaluation of a natural experiment. *J Epidemiol Community Health*. 2016.
- Carey IM, Anderson HR, Atkinson RW, Beevers S, Cook DG, Dajnak D, et al. Traffic pollution and the incidence of cardiorespiratory outcomes in an adult cohort in London. *Occup Environ Med*. 2016;73(12):849-56.
- Canavan C, West J, Card T. Calculating Total Health Service Utilisation and Costs from Routinely Collected Electronic Health Records Using the Example of Patients with Irritable Bowel Syndrome Before and After Their First Gastroenterology Appointment. *Pharmacoeconomics*. 2016;34(2):181-94.

- Brownrigg JRW, Hughes CO, Burleigh D, Karthikesalingam A, Patterson BO, Holt PJ, et al. Microvascular disease and risk of cardiovascular events among individuals with type 2 diabetes: a population-level cohort study. *Lancet Diabetes Endocrinol.* 2016;4(7):588-97.
- Brown A, Kirichek O, Balkwill A, Reeves G, Beral V, Sudlow C, et al. Comparison of dementia recorded in routinely collected hospital admission data in England with dementia recorded in primary care. *Emerging Themes in Epidemiology.* 2016;13(1).
- Brodie MJ, Chung S, Wade A, Quelen C, Guiraud-Diawara A, François C, et al. Clobazam and clonazepam use in epilepsy: Results from a UK database incident user cohort study. *Epilepsy Res.* 2016;123:68-74.
- Brauer R, Ruigómez A, Klungel O, Reynolds R, Feudjo Tepie M, Smeeth L, et al. The risk of acute liver injury among users of antibiotic medications: a comparison of case-only studies. *Pharmacoepidemiol Drug Saf.* 2016;25 Suppl 1:39-46.
- Brassard P, Wu JW, Ernst P, Dell'Aniello S, Smiechowski B, Suissa S. The effect of statins on influenza-like illness morbidity and mortality. *Pharmacoepidemiol Drug Saf.* 2016.
- Bours S, de Vries F, van den Bergh JPW, Lalmohamed A, van Staa TP, Leufkens HGM, et al. Risk of vertebral and non-vertebral fractures in patients with sarcoidosis: a population-based cohort. *Osteoporos Int.* 2016;27(4):1603-10.
- Bouras G, Markar SR, Burns EM, Mackenzie HA, Bottle A, Athanasiou T, et al. Linked Hospital and Primary Care Database Analysis of the Incidence and Impact of Psychiatric Morbidity Following Gastrointestinal Cancer Surgery in England. *Ann Surg.* 2016;264(1):93-9.
- Bornand D, Toovey S, Jick SS, Meier CR. The risk of new onset depression in association with influenza--A population-based observational study. *Brain Behav Immun.* 2016;53:131-7.
- Booth HP, Prevost AT, Gulliford MC. Severity of obesity and management of hypertension, hypercholesterolaemia and smoking in primary care: population-based cohort study. *J Hum Hypertens.* 2016;30(1):40-5.
- Booth HP, Khan O, Fildes A, Prevost AT, Reddy M, Charlton J, et al. Changing Epidemiology of Bariatric Surgery in the UK: Cohort Study Using Primary Care Electronic Health Records. *Obes Surg.* 2016;26(8):1900-5.
- Boef AGC, Souverein PC, Vandenbroucke JP, van Hylckama Vlieg A, de Boer A, le Cessie S, et al. Instrumental variable analysis as a complementary analysis in studies of adverse effects: venous thromboembolism and second-generation versus third-generation oral contraceptives. *Pharmacoepidemiol Drug Saf.* 2016;25(3):317-24.
- Bloechliger M, Ceschi A, Rüegg S, Kupferschmidt H, Kraehenbuehl S, Jick SS, et al. Risk of Seizures Associated with Antidepressant Use in Patients with Depressive Disorder: Follow-up Study with a Nested Case-Control Analysis Using the Clinical Practice Research Datalink. *Drug Safety.* 2016;39(4):307-21.
- Bloechliger M, Ceschi A, Rüegg S, Jick SS, Meier CR, Bodmer M. Lifestyle factors, psychiatric and neurologic comorbidities, and drug use associated with incident seizures among adult patients with depression: a population-based nested case-control study. *Eur J Epidemiol.* 2016.
- Ble A, Hughes PM, Delgado J, Masoli JA, Bowman K, Zirk-Sadowski J, et al. Safety and Effectiveness of Statins for Prevention of Recurrent Myocardial Infarction in 12 156 Typical Older Patients: A Quasi-Experimental Study. *J Gerontol A Biol Sci Med Sci.* 2016.

- Blak BT, Rigney U, Sternhufvud C, Davis J, Hammar N. Weight change and healthcare resource use in English patients with type 2 diabetes mellitus initiating a new diabetes medication class. *Int J Clin Pract*. 2016;70(1):45-55.
- Berni E, Scott LA, Jenkins-Jones S, De Voogd H, Rocha MS, Butler CC, et al. Non-Response to Antibiotic Treatment in Adolescents for Four Common Infections in UK Primary Care 1991-2012: A Retrospective, Longitudinal Study. *Antibiotics (Basel)*. 2016;5(3).
- Berni E, Butler CC, Jenkins-Jones S, de Voogd H, Ouwens M, Morgan CL, et al. Comparative estimated effectiveness of antibiotic classes as initial and secondary treatments of respiratory tract infections: longitudinal analysis of routine data from UK primary care 1991-2012. *Current Medical Research and Opinion*. 2016;32(6):1023-32.
- Bedson J, Chen Y, Hayward RA, Ashworth J, Walters K, Dunn KM, et al. Trends in long-term opioid prescribing in primary care patients with musculoskeletal conditions: an observational database study. *Pain*. 2016;157(7):1525-31.
- Beau-Lejdstrom R, Douglas I, Evans SJW, Smeeth L. Latest trends in ADHD drug prescribing patterns in children in the UK: prevalence, incidence and persistence. *BMJ Open*. 2016;6(6):e010508.
- Barker I, Lloyd T, Steventon A. Effect of a national requirement to introduce named accountable general practitioners for patients aged 75 or older in England: regression discontinuity analysis of general practice utilisation and continuity of care. *BMJ Open*. 2016;6(9):e011422.
- Baker R, Tata LJ, Kendrick D, Orton E. Identification of incident poisoning, fracture and burn events using linked primary care, secondary care and mortality data from England: implications for research and surveillance. *Inj Prev*. 2016;22(1):59-67.
- Baker R, Tata LJ, Kendrick D, Burch T, Kennedy M, Orton E. Differing patterns in thermal injury incidence and hospitalisations among 0-4 year old children from England. *Burns*. 2016;42(7):1609-16.
- Baker R, Orton E, Tata LJ, Kendrick D. Epidemiology of poisonings, fractures and burns among 0-24 year olds in England using linked health and mortality data. *Eur J Public Health*. 2016.
- Azoulay L, Soldera S, Yin H, Bouganim N. Use of Calcium Channel Blockers and Risk of Breast Cancer: A Population-based Cohort Study. *Epidemiology*. 2016;27(4):594-601.
- Asaria M, Walker S, Palmer S, Gale CP, Shah AD, Abrams KR, et al. Using electronic health records to predict costs and outcomes in stable coronary artery disease. *Heart*. 2016;102(10):755-62.
- Armstrong D, Ashworth M, Dregan A, White P. The relationship between prior antimicrobial prescription and meningitis: a case-control study. *Br J Gen Pract*. 2016;66(645):e228-33.
- Ankarfeldt MZ, Thorsted BL, Groenwold RH, Adalsteinsson E, Sanni A, Klungel O. Assessment of channelling bias among initiators of glucose lowering drugs-a cohort study. *Value in Health*. 2016;19(7):A667.
- Alexandre L, Clark AB, Bhutta HY, Chan SSM, Lewis MPN, Hart AR. Association Between Statin Use After Diagnosis of Esophageal Cancer and Survival: A Population-Based Cohort Study. *Gastroenterology*. 2016;150(4):854-65.e1; quiz e16-17.
- Alexander M, Petri H, Ding Y, Wandel C, Khwaja O, Foskett N. Morbidity and medication in a large population of individuals with Down syndrome compared to the general population. *Dev Med Child Neurol*. 2016;58(3):246-54.

- Alexander M, Ding Y, Foskett N, Petri H, Wandel C, Khwaja O. Population prevalence of Down's syndrome in the United Kingdom. *J Intellect Disabil Res.* 2016;60(9):874-8.
- Afonso A, Schmiedl S, Becker C, Tcherny-Lessenot S, Primatesa P, Plana E, et al. A methodological comparison of two European primary care databases and replication in a US claims database: inhaled long-acting beta-2-agonists and the risk of acute myocardial infarction. *Eur J Clin Pharmacol.* 2016;72(9):1105-16.
- Abdul Sultan A, West J, Ban L, Humes D, Tata LJ, Fleming KM, et al. Adverse Pregnancy Outcomes Among Women with Inflammatory Bowel Disease: A Population-Based Study from England. *Inflamm Bowel Dis.* 2016;22(7):1621-30.
- Zingone F, Sultan AA, Humes DJ, West J. Risk of acute appendicitis in and around pregnancy: a population-based cohort study from England. *Ann Surg.* 2015;261(2):332-7.
- Yu OHY, Yin H, Azoulay L. The combination of DPP-4 inhibitors versus sulfonylureas with metformin after failure of first-line treatment in the risk for major cardiovascular events and death. *Can J Diabetes.* 2015;39(5):383-9.
- Yu OHY, Filion KB, Azoulay L, Patenaude V, Majdan A, Suissa S. Incretin-based drugs and the risk of congestive heart failure. *Diabetes Care.* 2015;38(2):277-84.
- Yu J, Tang J, Li Z, Sajjan S, O'Regan C, Modi A, et al. History of cardiovascular events and cardiovascular risk factors among patients initiating strontium ranelate for treatment of osteoporosis. *Int J Womens Health.* 2015;7:913-8.
- Yska JP, van Roon EN, de Boer A, Leufkens HGM, Wilffert B, de Heide LJM, et al. Remission of Type 2 Diabetes Mellitus in Patients After Different Types of Bariatric Surgery: A Population-Based Cohort Study in the United Kingdom. *JAMA Surg.* 2015;150(12):1126-33.
- Wvan der Zanden R, de Vries F, Lalmohamed A, Driessen JHM, de Boer A, Rohde G, et al. Use of Dipeptidyl-Peptidase-4 Inhibitors and the Risk of Pneumonia: A Population-Based Cohort Study. *PLoS One.* 2015;10(10):e0139367.
- Wilson JC, Toovey S, Jick SS, Meier CR. Previously diagnosed influenza infections and the risk of developing epilepsy. *Epidemiol Infect.* 2015;143(11):2408-15.
- Wijnands JMA, van Durme CMPG, Driessen JHM, Boonen A, Klop C, Leufkens B, et al. Individuals With Type 2 Diabetes Mellitus Are at an Increased Risk of Gout But This Is Not Due to Diabetes: A Population-Based Cohort Study. *Medicine.* 2015;94(32):e1358.
- Weng SF, Kai J, Andrew Neil H, Humphries SE, Qureshi N. Improving identification of familial hypercholesterolaemia in primary care: derivation and validation of the familial hypercholesterolaemia case ascertainment tool (FAMCAT). *Atherosclerosis.* 2015;238(2):336-43.
- Warnier MJ, Rutten FH, Souverein PC, de Boer A, Hoes AW, De Bruin ML. Are ECG monitoring recommendations before prescription of QT-prolonging drugs applied in daily practice? The example of haloperidol. *Pharmacoepidemiol Drug Saf.* 2015;24(7):701-8.
- Vissers PAJ, Cardwell CR, van de Poll-Franse LV, Young IS, Pouter F, Murray LJ. The association between glucose-lowering drug use and mortality among breast cancer patients with type 2 diabetes. *Breast Cancer Res Treat.* 2015;150(2):427-37.
- Vinogradova Y, Coupland C, Hippisley-Cox J. Use of combined oral contraceptives and risk of venous

thromboembolism: nested case-control studies using the QResearch and CPRD databases. *BMJ*. 2015;350:h2135.

- Vinogradova Y, Coupland C, Brindle P, Hippisley-Cox J. Patients who discontinued statin treatment: a protocol for cohort study using primary care data. *BMJ Open*. 2015;5(10):e008701.
- Venkatesan S, Myles PR, McCann G, Kousoulis AA, Hashmi M, Belatri R, et al. Development of processes allowing near real-time refinement and validation of triage tools during the early stage of an outbreak in readiness for surge: the FLU-CATs Study. *Health Technol Assess*. 2015;19(89):1-132.
- van Jaarsveld CHM, Gulliford MC. Childhood obesity trends from primary care electronic health records in England between 1994 and 2013: population-based cohort study. *Arch Dis Child*. 2015;100(3):214-9.
- van den Ham HA, Klungel OH, Singer DE, Leufkens HGM, van Staa TP. Comparative Performance of ATRIA, CHADS2, and CHA2DS2-VASc Risk Scores Predicting Stroke in Patients With Atrial Fibrillation: Results From a National Primary Care Database. *J Am Coll Cardiol*. 2015;66(17):1851-9.
- Turner SW, Richardson K, Burden A, Thomas M, Murray C, Price D. Initial step-up treatment changes in asthmatic children already prescribed inhaled corticosteroids: a historical cohort study. *NPJ Prim Care Respir Med*. 2015;25:15041.
- Tuccori M, Wu JW, Yin H, Majdan A, Azoulay L. The Use of Glyburide Compared With Other Sulfonylureas and the Risk of Cancer in Patients With Type 2 Diabetes. *Diabetes Care*. 2015;38(11):2083-9.
- Strongman H, D'Oca K, Langerman H, Das R. Comparison of diabetes-associated secondary healthcare utilization between alternative oral antihyperglycaemic dual therapy combinations with metformin in patients with type 2 diabetes: an observational cohort study. *Diabetes Obes Metab*. 2015;17(6):573-80.
- Stolwijk C, Essers I, van Tubergen A, Boonen A, Bazelier MT, De Bruin ML, et al. The epidemiology of extra-articular manifestations in ankylosing spondylitis: a population-based matched cohort study. *Annals of the Rheumatic Diseases*. 2015;74(7):1373-8.
- Stocks SJ, Kontopantelis E, Akbarov A, Rodgers S, Avery AJ, Ashcroft DM. Examining variations in prescribing safety in UK general practice: cross sectional study using the Clinical Practice Research Datalink. *BMJ*. 2015;351:h5501.
- Springate DA, Ashcroft DM, Kontopantelis E, Doran T, Ryan R, Reeves D. Can analyses of electronic patient records be independently and externally validated? Study 2--the effect of β -adrenoceptor blocker therapy on cancer survival: a retrospective cohort study. *BMJ Open*. 2015;5(4):e007299.
- Spoenclin J, Meier C, Jick SS, Meier CR. Oral and inhaled glucocorticoid use and risk of Achilles or biceps tendon rupture: a population-based case-control study. *Ann Med*. 2015;47(6):492-8.
- Shephard EA, Neal RD, Rose PW, Walter FM, Hamilton WT. Quantifying the risk of non-Hodgkin lymphoma in symptomatic primary care patients aged ≥ 40 years: a large case-control study using electronic records. *Br J Gen Pract*. 2015;65(634):e281-8.
- Shephard EA, Neal RD, Rose P, Walter FM, Litt EJ, Hamilton WT. Quantifying the risk of multiple myeloma from symptoms reported in primary care patients: a large case-control study using electronic records. *Br J Gen Pract*. 2015;65(631):e106-13.
- Sammon CJ, Miller A, Mahtani KR, Holt TA, McHugh NJ, Luqmani RA, et al. Missing laboratory test data in electronic general practice records: analysis of rheumatoid factor recording in the clinical practice research

datalink. *Pharmacoepidemiol Drug Saf.* 2015;24(5):504-9.

- Sammon CJ, Charlton RA, Snowball J, Weil JG. The incidence of childhood and adolescent seizures in the UK from 1999 to 2011: A retrospective cohort study using the Clinical Practice Research Datalink. *Vaccine.* 2015;33(51):7364-9.
- Saine ME, Carbonari DM, Newcomb CW, Nezamzadeh MS, Haynes K, Roy JA, et al. Determinants of saxagliptin use among patients with type 2 diabetes mellitus treated with oral anti-diabetic drugs. *BMC Pharmacol Toxicol.* 2015;16:8.
- Robson JC, Kiran A, Maskell J, Hutchings A, Arden N, Dasgupta B, et al. The relative risk of aortic aneurysm in patients with giant cell arteritis compared with the general population of the UK. *Annals of the Rheumatic Diseases.* 2015;74(1):129-35.
- Ridd MJ, Ferreira DLS, Montgomery AA, Salisbury C, Hamilton W. Patient-doctor continuity and diagnosis of cancer: electronic medical records study in general practice. *Br J Gen Pract.* 2015;65(634):e305-11.
- Renoux C, Lix LM, Patenaude V, Bresee LC, Paterson JM, Lafrance J-P, et al. Serotonin-Norepinephrine Reuptake Inhibitors and the Risk of AKI: A Cohort Study of Eight Administrative Databases and Meta-Analysis. *Clin J Am Soc Nephrol.* 2015;10(10):1716-22.
- Renoux C, Dell'Aniello S, Saarela O, Filion KB, Boivin J-F. Antiepileptic drugs and the risk of ischaemic stroke and myocardial infarction: a population-based cohort study. *BMJ Open.* 2015;5(8):e008365.
- Reinau D, Surber C, Jick SS, Meier CR. Nonsteroidal anti-inflammatory drugs and the risk of nonmelanoma skin cancer. *Int J Cancer.* 2015;137(1):144-53.
- Reilly S, Olier I, Planner C, Doran T, Reeves D, Ashcroft DM, et al. Inequalities in physical comorbidity: a longitudinal comparative cohort study of people with severe mental illness in the UK. *BMJ Open.* 2015;5(12):e009010.
- Redaniel MT, Martin RM, Ridd MJ, Wade J, Jeffreys M. Diagnostic intervals and its association with breast, prostate, lung and colorectal cancer survival in England: historical cohort study using the Clinical Practice Research Datalink. *PLoS One.* 2015;10(5):e0126608.
- Qizilbash N, Gregson J, Johnson ME, Pearce N, Douglas I, Wing K, et al. BMI and risk of dementia in two million people over two decades: a retrospective cohort study. *Lancet Diabetes Endocrinol.* 2015;3(6):431-6.
- Puneekar YS, Wurst K, Shukla A. Resource Use and Costs up to Two Years Post Diagnosis Among Newly Diagnosed COPD Patients in the UK Primary Care Setting: A Retrospective Cohort Study. *COPD.* 2015;12(3):267-75.
- Puneekar YS, Landis SH, Wurst K, Le H. Characteristics, disease burden and costs of COPD patients in the two years following initiation of long-acting bronchodilators in UK primary care. *Respir Res.* 2015;16:141.
- Prieto-Alhambra D, Javaid MK, Judge A, Maskell J, Cooper C, Arden NK, et al. Hormone replacement therapy and mid-term implant survival following knee or hip arthroplasty for osteoarthritis: a population-based cohort study. *Annals of the Rheumatic Diseases.* 2015;74(3):557-63.
- Pockett RD, Watkins J, McEwan P, Meier G. Burden of Illness in UK Subjects with Reported Respiratory Infections Vaccinated or Unvaccinated against Influenza: A Retrospective Observational Study. *PLoS One.* 2015;10(8):e0134928.

- Pfeil AM, Imfeld P, Pettengell R, Jick SS, Szucs TD, Meier CR, et al. Trends in incidence and medical resource utilisation in patients with chronic lymphocytic leukaemia: insights from the UK Clinical Practice Research Datalink (CPRD). *Ann Hematol.* 2015;94(3):421-9.
- Petri H, Nevitt A, Sarsour K, Napalkov P, Collinson N. Incidence of giant cell arteritis and characteristics of patients: data-driven analysis of comorbidities. *Arthritis Care Res (Hoboken).* 2015;67(3):390-5.
- Petherick ES, Pickett KE, Cullum NA. Can different primary care databases produce comparable estimates of burden of disease: results of a study exploring venous leg ulceration. *Fam Pract.* 2015;32(4):374-80.
- Perera R, McFadden E, McLellan J, Lung T, Clarke P, Pérez T, et al. Optimal strategies for monitoring lipid levels in patients at risk or with cardiovascular disease: a systematic review with statistical and cost-effectiveness modelling. *Health Technol Assess.* 2015;19(100):1-401, vii-viii.
- Peeters PJHL, Bazelier MT, Leufkens HGM, de Vries F, De Bruin ML. The risk of colorectal cancer in patients with type 2 diabetes: associations with treatment stage and obesity. *Diabetes Care.* 2015;38(3):495-502.
- Paul SK, Klein K, Thorsted BL, Wolden ML, Khunti K. Delay in treatment intensification increases the risks of cardiovascular events in patients with type 2 diabetes. *Cardiovasc Diabetol.* 2015;14:100.
- Parisi R, Rutter MK, Lunt M, Young HS, Symmons DPM, Griffiths CEM, et al. Psoriasis and the Risk of Major Cardiovascular Events: Cohort Study Using the Clinical Practice Research Datalink. *J Invest Dermatol.* 2015;135(9):2189-97.
- Palmer KT, Walker-Bone K, Harris EC, Linaker C, D'Angelo S, Sayer AA, et al. Health and Employment after Fifty (HEAF): a new prospective cohort study. *BMC Public Health.* 2015;15:1071.
- Palmer KT, D'Angelo S, Harris EC, Linaker C, Coggon D. Sensory impairments, problems of balance and accidental injury at work: a case-control study. *Occup Environ Med.* 2015;72(3):195-9.
- Palmaro A, Dupouy J, Lapeyre-Mestre M. Benzodiazepines and risk of death: Results from two large cohort studies in France and UK. *Eur Neuropsychopharmacol.* 2015;25(10):1566-77.
- O'Rorke MA, Murray LJ, Hughes CM, Cantwell MM, Cardwell CR. The effect of warfarin therapy on breast, colorectal, lung, and prostate cancer survival: a population-based cohort study using the Clinical Practice Research Datalink. *Cancer Causes Control.* 2015;26(3):355-66.
- Olsson M, Schnecke V, Cabrera C, Skrtic S, Lind M. Contemporary Risk Estimates of Three HbA1c Variables for Myocardial Infarction in 101,799 Patients Following Diagnosis of Type 2 Diabetes. *Diabetes Care.* 2015;38(8):1481-6.
- Nyeland ME, Ploug UJ, Richards A, Garcia Alvarez L, Demuth D, Muthutantri A, et al. Evaluation of the effectiveness of liraglutide and sitagliptin in type 2 diabetes: a retrospective study in UK primary care. *Int J Clin Pract.* 2015;69(3):281-91.
- Numbere B, Fleming KM, Walker A, Card TR. Adrenergic blockers and the risk for common solid cancers: a case-control study. *Eur J Cancer Prev.* 2015.
- Nielen JT, van den Bemt BJJ, Boonen A, Dagnelie PC, Emans PJ, Veldhorst N, et al. Identification of antithrombotic drugs related to total joint replacement using anonymised free-text notes: a search strategy in the Clinical Practice Research Datalink. *BMJ Open.* 2015;5(11):e009017.
- Nakafero G, Sanders RD, Nguyen-Van-Tam JS, Myles PR. Association between benzodiazepine use and

exacerbations and mortality in patients with asthma: a matched case-control and survival analysis using the United Kingdom Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf.* 2015;24(8):793-802.

- Muller S, Hider SL, Raza K, Stack RJ, Hayward RA, Mallen CD. An algorithm to identify rheumatoid arthritis in primary care: a Clinical Practice Research Datalink study. *BMJ Open.* 2015;5(12):e009309.
- Mukherjee S, Culliford D, Arden N, Edwards C. What is the risk of having a total hip or knee replacement for patients with lupus? *Lupus.* 2015;24(2):198-202.
- Mt-Isa S, Tomlin S, Sutcliffe A, Underwood M, Williamson P, Croft NM, et al. Prokinetics prescribing in paediatrics: evidence on cisapride, domperidone, and metoclopramide. *J Pediatr Gastroenterol Nutr.* 2015;60(4):508-14.
- Millett ERC, De Stavola BL, Quint JK, Smeeth L, Thomas SL. Risk factors for hospital admission in the 28 days following a community-acquired pneumonia diagnosis in older adults, and their contribution to increasing hospitalisation rates over time: a cohort study. *BMJ Open.* 2015;5(12):e008737.
- Miller A, Nightingale AL, Sammon CJ, Mahtani KR, Holt TA, McHugh NJ, et al. Estimating the diagnostic accuracy of rheumatoid factor in UK primary care: a study using the Clinical Practice Research Datalink. *Rheumatology (Oxford).* 2015;54(10):1882-9.
- Melzer D, Tavakoly B, Winder RE, Masoli JAH, Henley WE, Ble A, et al. Much more medicine for the oldest old: trends in UK electronic clinical records. *Age Ageing.* 2015;44(1):46-53.
- Mehta HB, Mehta V, Tsai C-L, Chen H, Aparasu RR, Johnson ML. Development and Validation of the RxDx-Dementia Risk Index to Predict Dementia in Patients with Type 2 Diabetes and Hypertension. *J Alzheimers Dis.* 2015;49(2):423-32.
- McGlynn KA, Hagberg K, Chen J, Graubard BI, London WT, Jick S, et al. Statin use and risk of primary liver cancer in the Clinical Practice Research Datalink. *J Natl Cancer Inst.* 2015;107(4).
- McFadden E, Stevens R, Glasziou P, Perera R. Implications of lower risk thresholds for statin treatment in primary prevention: analysis of CPRD and simulation modelling of annual cholesterol monitoring. *Prev Med.* 2015;70:14-6.
- McDonald HI, Thomas SL, Millett ERC, Nitsch D. CKD and the risk of acute, community-acquired infections among older people with diabetes mellitus: a retrospective cohort study using electronic health records. *Am J Kidney Dis.* 2015;66(1):60-8.
- Mc Menamin ÚC, Cardwell CR, Hughes CM, Murray LM. Low-dose aspirin and survival from lung cancer: a population-based cohort study. *BMC cancer.* 2015;15:911.
- Martinez C, Wallenhorst C, McFerran D, Hall DA. Incidence rates of clinically significant tinnitus: 10-year trend from a cohort study in England. *Ear Hear.* 2015;36(3):e69-75.
- Martinez C, Katholing A, Wallenhorst C, Granziera S, Cohen AT, Freedman SB. Increasing incidence of non-valvular atrial fibrillation in the UK from 2001 to 2013. *Heart.* 2015;101(21):1748-54.
- Macedo AF, Bell J, McCarron C, Conroy R, Richardson J, Scowcroft A, et al. Determinants of oral anticoagulation control in new warfarin patients: analysis using data from Clinical Practice Research Datalink. *Thromb Res.* 2015;136(2):250-60.
- Liang SF, Taweel A, Miles S, Kovalchuk Y, Spiridou A, Barratt B, et al. Semi automated transformation to OWL

formatted files as an approach to data integration. A feasibility study using environmental, disease register and primary care clinical data. *Methods Inf Med*. 2015;54(1):32-40.

- Li L, Hagberg KW, Peng M, Shah K, Paris M, Jick S. Rates of Cardiovascular Disease and Major Adverse Cardiovascular Events in Patients With Psoriatic Arthritis Compared to Patients Without Psoriatic Arthritis. *J Clin Rheumatol*. 2015;21(8):405-10.
- Levin D, Bell S, Sund R, Hartikainen SA, Tuomilehto J, Pukkala E, et al. Pioglitazone and bladder cancer risk: a multipopulation pooled, cumulative exposure analysis. *Diabetologia*. 2015;58(3):493-504.
- Lapi F, Azoulay L, Kezouh A, Benisty J, Matok I, Mugelli A, et al. The use of antidepressants and the risk of chronic atrial fibrillation. *J Clin Pharmacol*. 2015;55(4):423-30.
- Lamy FX, Chollet J, Clay E, Brignone M, Rive B, Saragoussi D. Pharmacotherapeutic strategies for patients treated for depression in UK primary care: a database analysis. *Current Medical Research and Opinion*. 2015;31(4):795-807.
- Kuo C-F, Grainge MJ, Mallen C, Zhang W, Doherty M. Rising burden of gout in the UK but continuing suboptimal management: a nationwide population study. *Annals of the Rheumatic Diseases*. 2015;74(4):661-7.
- Koshy E, Watt H, Curcin V, Bottle A, Sharland M, Saxena S. Tonsillectomy among children with low baseline acute throat infection consultation rates in UK general practices: a cohort study. *BMJ Open*. 2015;5(2):e006686.
- Kontopantelis E, Springate DA, Reeves D, Ashcroft DM, Rutter MK, Rutter M, et al. Glucose, blood pressure and cholesterol levels and their relationships to clinical outcomes in type 2 diabetes: a retrospective cohort study. *Diabetologia*. 2015;58(3):505-18.
- Kontopantelis E, Olier I, Planner C, Reeves D, Ashcroft DM, Gask L, et al. Primary care consultation rates among people with and without severe mental illness: a UK cohort study using the Clinical Practice Research Datalink. *BMJ Open*. 2015;5(12):e008650.
- Klop C, Gibson-Smith D, Elders PJM, Welsing PMJ, Leufkens HGM, Harvey NC, et al. Anti-osteoporosis drug prescribing after hip fracture in the UK: 2000-2010. *Osteoporos Int*. 2015;26(7):1919-28.
- Khunti K, Davies M, Majeed A, Thorsted BL, Wolden ML, Paul SK. Hypoglycemia and risk of cardiovascular disease and all-cause mortality in insulin-treated people with type 1 and type 2 diabetes: a cohort study. *Diabetes Care*. 2015;38(2):316-22.
- Kendrick T, Stuart B, Newell C, Geraghty AWA, Moore M. Did NICE guidelines and the Quality Outcomes Framework change GP antidepressant prescribing in England? Observational study with time trend analyses 2003-2013. *J Affect Disord*. 2015;186:171-7.
- Karasneh RA, Murray LJ, Mc Menamin ÚC, Hughes CM, Cardwell CR. Digoxin use after diagnosis of breast cancer and survival: a population-based cohort study. *Breast Cancer Res Treat*. 2015;151(3):661-9.
- Karasneh RA, Murray LJ, Hughes CM, Cardwell CR. Digoxin use after diagnosis of colorectal cancer and survival: a population-based cohort study. *Cancer Epidemiol Biomarkers Prev*. 2015;24(11):1804-7.
- Kang EM, Pinheiro SP, Hammad TA, Abou-Ali A. Evaluating the validity of clinical codes to identify cataract and glaucoma in the UK Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf*. 2015;24(1):38-44.

- Judge A, Wallace G, Prieto-Alhambra D, Arden NK, Edwards CJ. Can the publication of guidelines change the management of early rheumatoid arthritis? An interrupted time series analysis from the United Kingdom. *Rheumatology (Oxford)*. 2015;54(12):2244-8.
- Jick SS, Li L, Falcone GJ, Vassilev ZP, Wallander MA. Epidemiology of multiple sclerosis: results from a large observational study in the UK. *J Neurol*. 2015;262(9):2033-41.
- Jick S, Li L, Gastanaga VM, Liede A. Prevalence of hypercalcemia of malignancy among cancer patients in the UK: analysis of the Clinical Practice Research Datalink database. *Cancer Epidemiol*. 2015;39(6):901-7.
- Jeffreys M, Redaniel MT, Martin RM. The effect of pre-diagnostic vitamin D supplementation on cancer survival in women: a cohort study within the UK Clinical Practice Research Datalink. *BMC cancer*. 2015;15:670.
- Imfeld P, Bodmer M, Jick SS, Meier CR. Benzodiazepine Use and Risk of Developing Alzheimer's Disease or Vascular Dementia: A Case-Control Analysis. *Drug Safety*. 2015;38(10):909-19.
- Ilo D, Raluy-Callado M, Graham-Clarke P, Sadasivan R, Birt J, Donaldson R, et al. Patient characteristics and treatment patterns for patients with benign prostatic hyperplasia, erectile dysfunction or co-occurring benign prostatic hyperplasia and erectile dysfunction in general practices in the UK: a retrospective observational study. *Int J Clin Pract*. 2015;69(8):853-62.
- Humes DJ, Walker AJ, Blackwell J, Hunt BJ, West J. Variation in the risk of venous thromboembolism following colectomy. *Br J Surg*. 2015;102(13):1629-38.
- Humes DJ, Nordenskjöld A, Walker AJ, West J, Ludvigsson JF. Risk of venous thromboembolism in children after general surgery. *J Pediatr Surg*. 2015;50(11):1870-3.
- Horspool MJ, Julious SA, Mooney C, May R, Sully B, Smithson WH. Preventing and Lessening Exacerbations of Asthma in School-aged children Associated with a New Term (PLEASANT): Recruiting Primary Care Research Sites-the PLEASANT experience. *NPJ Prim Care Respir Med*. 2015;25:15066.
- Hopkins C, Andrews P, Holy CE. Does time to endoscopic sinus surgery impact outcomes in chronic rhinosinusitis? Retrospective analysis using the UK clinical practice research data. *Rhinology*. 2015;53(1):18-24.
- Holden SE, Jenkins-Jones S, Morgan CL, Scherthaner G, Currie CJ. Glucose-lowering with exogenous insulin monotherapy in type 2 diabetes: dose association with all-cause mortality, cardiovascular events and cancer. *Diabetes Obes Metab*. 2015;17(4):350-62.
- Hire AJ, Ashcroft DM, Springate DA, Steinke DT. ADHD in the United Kingdom: Regional and Socioeconomic Variations in Incidence Rates Amongst Children and Adolescents (2004-2013). *J Atten Disord*. 2015.
- Hippisley-Cox J, Coupland C. Development and validation of risk prediction equations to estimate future risk of heart failure in patients with diabetes: a prospective cohort study. *BMJ Open*. 2015;5(9):e008503.
- Hippisley-Cox J, Coupland C. Development and validation of risk prediction equations to estimate future risk of blindness and lower limb amputation in patients with diabetes: cohort study. *BMJ*. 2015;351:h5441.
- Hicks BM, Murray LJ, Hughes C, Cardwell CR. Post-diagnostic oral bisphosphonate use and colorectal cancer mortality: a population-based cohort study within the UK Clinical Practice Research Datalink. *Br J Cancer*. 2015;113(1):123-6.
- Hicks BM, Murray LJ, Hughes C, Cardwell CR. Clopidogrel use and cancer-specific mortality: a population-

based cohort study of colorectal, breast and prostate cancer patients. *Pharmacoepidemiol Drug Saf.* 2015;24(8):830-40.

- Herrett E, Gallagher AM, Bhaskaran K, Forbes H, Mathur R, van Staa T, et al. Data Resource Profile: Clinical Practice Research Datalink (CPRD). *Int J Epidemiol.* 2015;44(3):827-36.
- Hazra NC, Dregan A, Jackson S, Gulliford MC. Differences in Health at Age 100 According to Sex: Population-Based Cohort Study of Centenarians Using Electronic Health Records. *Journal of the American Geriatrics Society.* 2015;63(7):1331-7.
- Hazra N, Dregan A, Charlton J, Gulliford MC, D'Cruz DP. Incidence and mortality of relapsing polychondritis in the UK: a population-based cohort study. *Rheumatology (Oxford).* 2015;54(12):2181-7.
- Hayes JF, Bhaskaran K, Batterham R, Smeeth L, Douglas I. The effect of sibutramine prescribing in routine clinical practice on cardiovascular outcomes: a cohort study in the United Kingdom. *Int J Obes (Lond).* 2015;39(9):1359-64.
- Hamada S, Gulliford MC. Antidiabetic and cardiovascular drug utilisation in patients diagnosed with type 2 diabetes mellitus over the age of 80 years: a population-based cohort study. *Age Ageing.* 2015;44(4):566-73.
- Gordon JP, Evans M, Puelles J, McEwan PC. Factors Predictive of Weight Gain and Implications for Modeling in Type 2 Diabetes Patients Initiating Metformin and Sulfonylurea Combination Therapy. *Diabetes Ther.* 2015;6(4):495-507.
- Goossens ME, Zeegers MP, Bazelier MT, De Bruin ML, Buntinx F, de Vries F. Risk of bladder cancer in patients with diabetes: a retrospective cohort study. *BMJ Open.* 2015;5(6):e007470.
- Goossens ME, Buntinx F, Zeegers MP, Driessen JHM, De Bruin ML, De Vries F. Influence of metformin intake on the risk of bladder cancer in type 2 diabetes patients. *British Journal of Clinical Pharmacology.* 2015;80(6):1464-72.
- Goh KL, Bhaskaran K, Minassian C, Evans SJW, Smeeth L, Douglas IJ. Angiotensin receptor blockers and risk of dementia: cohort study in UK Clinical Practice Research Datalink. *British Journal of Clinical Pharmacology.* 2015;79(2):337-50.
- Gallagher N, Cardwell C, Hughes C, O'Reilly D. Increase in the pharmacological management of Type 2 diabetes with pay-for-performance in primary care in the UK. *Diabet Med.* 2015;32(1):62-8.
- Fournier J-P, Yin H, Nessim SJ, Montastruc J-L, Azoulay L. Tramadol for noncancer pain and the risk of hyponatremia. *Am J Med.* 2015;128(4):418-25.e5.
- Fournier J-P, Azoulay L, Yin H, Montastruc J-L, Suissa S. Tramadol use and the risk of hospitalization for hypoglycemia in patients with noncancer pain. *JAMA Intern Med.* 2015;175(2):186-93.
- Forster AS, Dodhia H, Booth H, Dregan A, Fuller F, Miller J, et al. Estimating the yield of NHS Health Checks in England: a population-based cohort study. *J Public Health (Oxf).* 2015;37(2):234-40.
- Fleming DM, Taylor RJ, Lustig RL, Schuck-Paim C, Haguinet F, Webb DJ, et al. Modelling estimates of the burden of Respiratory Syncytial virus infection in adults and the elderly in the United Kingdom. *BMC Infect Dis.* 2015;15:443.
- Fildes A, Charlton J, Rudisill C, Littlejohns P, Prevost AT, Gulliford MC. Probability of an Obese Person Attaining Normal Body Weight: Cohort Study Using Electronic Health Records. *Am J Public Health.*

2015;105(9):e54-9.

- Feudjo-Tepie M, Ferguson S, Roddam A, Taylor A, Bayly J, Critchlow C. Comorbidities, bone-sparing agent prescription history and their determinants among postmenopausal women in UK primary care settings: a retrospective database study. *Arch Osteoporos*. 2015;10:41.
- Ferguson S, Feudjo Tepie M, Taylor A, Roddam A, Critchlow C, Iqbal M, et al. The impact of persistence with bisphosphonates on health resource utilization and fracture risk in the UK: a study of patient records from the UK Clinical Practice Research Datalink. *J Eval Clin Pract*. 2015.
- Faillie JL, Filion KB, Patenaude V, Ernst P, Azoulay L. Dipeptidyl peptidase-4 inhibitors and the risk of community-acquired pneumonia in patients with type 2 diabetes. *Diabetes Obes Metab*. 2015;17(4):379-85.
- Edson-Heredia E, Zhu B, Lefevre C, Wang M, Barrett A, Bushe CJ, et al. Prevalence and incidence rates of cardiovascular, autoimmune, and other diseases in patients with psoriatic or psoriatic arthritis: a retrospective study using Clinical Practice Research Datalink. *J Eur Acad Dermatol Venereol*. 2015;29(5):955-63.
- Driessen JHM, Henry RMA, van Onzenoort HAW, Lalmohamed A, Burden AM, Prieto-Alhambra D, et al. Bone fracture risk is not associated with the use of glucagon-like peptide-1 receptor agonists: a population-based cohort analysis. *Calcif Tissue Int*. 2015;97(2):104-12.
- Douglas IJ, Bhaskaran K, Batterham RL, Smeeth L. The effectiveness of pharmaceutical interventions for obesity: weight loss with orlistat and sibutramine in a United Kingdom population-based cohort. *British Journal of Clinical Pharmacology*. 2015;79(6):1020-7.
- Douglas IJ, Bhaskaran K, Batterham RL, Smeeth L. Bariatric Surgery in the United Kingdom: A Cohort Study of Weight Loss and Clinical Outcomes in Routine Clinical Care. *PLoS Med*. 2015;12(12):e1001925.
- de Lusignan S, Liaw S-T, Dedman D, Khunti K, Sadek K, Jones S. An algorithm to improve diagnostic accuracy in diabetes in computerised problem orientated medical records (POMR) compared with an established algorithm developed in episode orientated records (EOMR). *J Innov Health Inform*. 2015;22(2):255-64.
- Davies NM, Taylor G, Taylor AE, Thomas KH, Windmeijer F, Martin RM, et al. What are the effects of varenicline compared with nicotine replacement therapy on long-term smoking cessation and clinically important outcomes? Protocol for a prospective cohort study. *BMJ Open*. 2015;5(11):e009665.
- Culliford D, Maskell J, Judge A, Cooper C, Prieto-Alhambra D, Arden NK, et al. Future projections of total hip and knee arthroplasty in the UK: results from the UK Clinical Practice Research Datalink. *Osteoarthritis and Cartilage*. 2015;23(4):594-600.
- Crooks CJ, West J, Card TR. A comparison of the recording of comorbidity in primary and secondary care by using the Charlson Index to predict short-term and long-term survival in a routine linked data cohort. *BMJ Open*. 2015;5(6):e007974.
- Close H, Mason JM, Wilson DW, Hungin APS, Jones R, Rubin G. Risk of Ischaemic Heart Disease in Patients with Inflammatory Bowel Disease: Cohort Study Using the General Practice Research Database. *PloS One*. 2015;10(10):e0139745.
- Clarson LE, Hider SL, Belcher J, Heneghan C, Roddy E, Mallen CD. Increased risk of vascular disease associated with gout: a retrospective, matched cohort study in the UK clinical practice research datalink. *Annals of the Rheumatic Diseases*. 2015;74(4):642-7.
- Cid Ruzafa J, Paczkowski R, Boye KS, Di Tanna GL, Sheetz MJ, Donaldson R, et al. Estimated glomerular

filtration rate progression in UK primary care patients with type 2 diabetes and diabetic kidney disease: a retrospective cohort study. *Int J Clin Pract.* 2015;69(8):871-82.

- Chu TPC, Shah A, Walker D, Coleman MP. Pattern of symptoms and signs of primary intracranial tumours in children and young adults: a record linkage study. *Arch Dis Child.* 2015;100(12):1115-22.
- Chhaya V, Saxena S, Cecil E, Subramanian V, Curcin V, Majeed A, et al. Impact of timing and duration of thiopurine treatment on first perianal surgery in Crohn's disease: UK population-based study (1995-2009). *Inflamm Bowel Dis.* 2015;21(2):385-91.
- Charlton RA, Snowball JM, Nightingale AL, Davis KJ. Safety of Fluticasone Propionate Prescribed for Asthma During Pregnancy: A UK Population-Based Cohort Study. *J Allergy Clin Immunol Pract.* 2015;3(5):772-9.e3.
- Charlton R, Garne E, Wang H, Klungsøyr K, Jordan S, Neville A, et al. Antiepileptic drug prescribing before, during and after pregnancy: a study in seven European regions. *Pharmacoepidemiol Drug Saf.* 2015;24(11):1144-54.
- Chang KC-M, Soljak M, Lee JT, Woringer M, Johnston D, Khunti K, et al. Coverage of a national cardiovascular risk assessment and management programme (NHS Health Check): Retrospective database study. *Prev Med.* 2015;78:1-8.
- Cardwell CR, Mc Menamin Ú, Hughes CM, Murray LJ. Statin use and survival from lung cancer: a population-based cohort study. *Cancer Epidemiol Biomarkers Prev.* 2015;24(5):833-41.
- Cardwell CR, Hicks BM, Hughes C, Murray LJ. Statin use after diagnosis of breast cancer and survival: a population-based cohort study. *Epidemiology.* 2015;26(1):68-78.
- Carbonari DM, Saine ME, Newcomb CW, Blak B, Roy JA, Haynes K, et al. Use of demographic and pharmacy data to identify patients included within both the Clinical Practice Research Datalink (CPRD) and The Health Improvement Network (THIN). *Pharmacoepidemiol Drug Saf.* 2015;24(9):999-1003.
- Cao X, Ejzykowicz F, Ramey DR, Sajjan S, Ambegaonkar BM, Mavros P, et al. Impact of Switching From High-Efficacy Lipid-Lowering Therapies to Generic Simvastatin on LDL-C Levels and LDL-C Goal Attainment Among High-Risk Primary and Secondary Prevention Populations in the United Kingdom. *Clin Ther.* 2015;37(4):804-15.
- Bruderer SG, Bodmer M, Jick SS, Meier CR. Poorly controlled type 2 diabetes mellitus is associated with a decreased risk of incident gout: a population-based case-control study. *Annals of the Rheumatic Diseases.* 2015;74(9):1651-8.
- Bruderer SG, Bodmer M, Jick SS, Meier CR. Association of hormone therapy and incident gout: population-based case-control study. *Menopause.* 2015;22(12):1335-42.
- Brauer R, Smeeth L, Anaya-Izquierdo K, Timmis A, Denaxas SC, Farrington CP, et al. Antipsychotic drugs and risks of myocardial infarction: a self-controlled case series study. *Eur Heart J.* 2015;36(16):984-92.
- Brauer R, Bhaskaran K, Chaturvedi N, Dexter DT, Smeeth L, Douglas I. Glitazone Treatment and Incidence of Parkinson's Disease among People with Diabetes: A Retrospective Cohort Study. *PLoS Med.* 2015;12(7):e1001854.
- Bouras G, Burns EM, Howell A-M, Bottle A, Athanasiou T, Darzi A. Risk of Post-Discharge Venous Thromboembolism and Associated Mortality in General Surgery: A Population-Based Cohort Study Using Linked Hospital and Primary Care Data in England. *PLoS One.* 2015;10(12):e0145759.

- Booth HP, Prevost AT, Gulliford MC. Access to weight reduction interventions for overweight and obese patients in UK primary care: population-based cohort study. *BMJ Open*. 2015;5(1):e006642.
- Bloechliger M, Rüegg S, Jick SS, Meier CR, Bodmer M. Antipsychotic drug use and the risk of seizures: follow-up study with a nested case-control analysis. *CNS Drugs*. 2015;29(7):591-603.
- Ble A, Masoli JAH, Barry HE, Winder RE, Tavakoly B, Henley WE, et al. Any versus long-term prescribing of high risk medications in older people using 2012 Beers Criteria: results from three cross-sectional samples of primary care records for 2003/4, 2007/8 and 2011/12. *BMC Geriatr*. 2015;15:146.
- Black RJ, Joseph RM, Brown B, Movahedi M, Lunt M, Dixon WG. Half of U.K. patients with rheumatoid arthritis are prescribed oral glucocorticoid therapy in primary care: a retrospective drug utilisation study. *Arthritis Res Ther*. 2015;17:375.
- Been JV, Szatkowski L, van Staa T-P, Leufkens HG, van Schayck OC, Sheikh A, et al. Smoke-free legislation and the incidence of paediatric respiratory infections and wheezing/asthma: interrupted time series analyses in the four UK nations. *Scientific Reports*. 2015;5:15246.
- Becker C, Wilson JC, Jick SS, Meier CR. Non-steroidal anti-inflammatory drugs and the risk of head and neck cancer: A case-control analysis. *Int J Cancer*. 2015;137(10):2424-31.
- Becker C, Jick SS, Meier CR, Bodmer M. No evidence for a decreased risk of thyroid cancer in association with use of metformin or other antidiabetic drugs: a case-control study. *BMC cancer*. 2015;15:719.
- Baril L, Rosillon D, Willame C, Angelo MG, Zima J, van den Bosch JH, et al. Risk of spontaneous abortion and other pregnancy outcomes in 15-25 year old women exposed to human papillomavirus-16/18 AS04- adjuvanted vaccine in the United Kingdom. *Vaccine*. 2015;33(48):6884-91.
- Barakat MF, McDonald HI, Collier TJ, Smeeth L, Nitsch D, Quint JK. Acute kidney injury in stable COPD and at exacerbation. *Int J Chron Obstruct Pulmon Dis*. 2015;10:2067-77.
- Baker A, Chen L-C, Elliott RA, Godman B. The impact of the 'Better Care Better Value' prescribing policy on the utilisation of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers for treating hypertension in the UK primary care setting: longitudinal quasi-experimental design. *BMC Health Serv Res*. 2015;15:367.
- Azoulay L, Eberg M, Benayoun S, Pollak M. 5 α -Reductase Inhibitors and the Risk of Cancer-Related Mortality in Men With Prostate Cancer. *JAMA Oncol*. 2015;1(3):314-20.
- Atkinson RW, Carey IM, Kent AJ, van Staa TP, Anderson HR, Cook DG. Long-term exposure to outdoor air pollution and the incidence of chronic obstructive pulmonary disease in a national English cohort. *Occup Environ Med*. 2015;72(1):42-8.
- Assayag J, Pollak MN, Azoulay L. The use of aspirin and the risk of mortality in patients with prostate cancer. *J Urol*. 2015;193(4):1220-5.
- Asghar Z, Coupland C, Siriwardena N. Influenza vaccination and risk of stroke: Self-controlled case-series study. *Vaccine*. 2015;33(41):5458-63.
- Arana A, Johannes CB, McQuay LJ, Varas-Lorenzo C, Fife D, Rothman KJ. Risk of Out-of-Hospital Sudden Cardiac Death in Users of Domperidone, Proton Pump Inhibitors, or Metoclopramide: A Population-Based Nested Case-Control Study. *Drug Safety*. 2015;38(12):1187-99.

- Ali AK, Hartzema AG, Winterstein AG, Segal R, Lu X, Hendeles L. Application of multicategory exposure marginal structural models to investigate the association between long-acting beta-agonists and prescribing of oral corticosteroids for asthma exacerbations in the Clinical Practice Research Datalink. *Value Health*. 2015;18(2):260-70.
- Abdul Sultan A, Crooks CJ, Card T, Tata LJ, Fleming KM, West J. Causes of death in people with coeliac disease in England compared with the general population: a competing risk analysis. *Gut*. 2015;64(8):1220-6.
- Zin CS, Chen LC, Knaggs RD. Changes in trends and pattern of strong opioid prescribing in primary care. *Eur J Pain*. 2014;18(9):1343-51.
- Zhou EH, Gelperin K, Levenson MS, Rose M, Hsueh Y-H, Graham DJ. Risk of acute myocardial infarction, stroke, or death in patients initiating olmesartan or other angiotensin receptor blockers - a cohort study using the Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf*. 2014;23(4):340-7.
- Wurst KE, Shukla A, Muellerova H, Davis KJ. Respiratory pharmacotherapy use in patients newly diagnosed with chronic obstructive pulmonary disease in a primary care setting in the UK: a retrospective cohort study. *COPD*. 2014;11(5):521-30.
- Wurst KE, Puneekar YS, Shukla A. Treatment evolution after COPD diagnosis in the UK primary care setting. *PLoS One*. 2014;9(9):e105296.
- Wing K, Douglas I, Bhaskaran K, Klungel OH, Reynolds RF, Pirmohamed M, et al. Development of predictive genetic tests for improving the safety of new medicines: the utilization of routinely collected electronic health records. *Drug Discov Today*. 2014;19(4):361-6.
- Wilson AB, Neogi T, Prout M, Jick S. Relative risk of myelodysplastic syndromes in patients with autoimmune disorders in the General Practice Research Database. *Cancer Epidemiol*. 2014;38(5):544-9.
- West J, Fleming KM, Tata LJ, Card TR, Crooks CJ. Incidence and prevalence of celiac disease and dermatitis herpetiformis in the UK over two decades: population-based study. *Am J Gastroenterol*. 2014;109(5):757-68.
- Weich S, Pearce HL, Croft P, Singh S, Crome I, Bashford J, et al. Effect of anxiolytic and hypnotic drug prescriptions on mortality hazards: retrospective cohort study. *BMJ*. 2014;348:g1996.
- Wei L, Mackenzie IS, MacDonald TM, George J. Cardiovascular risk associated with sodium-containing medicines. *Expert Opin Drug Saf*. 2014;13(11):1515-23.
- Wallace G, Judge A, Prieto-Alhambra D, de Vries F, Arden NK, Cooper C. The effect of body mass index on the risk of post-operative complications during the 6 months following total hip replacement or total knee replacement surgery. *Osteoarthritis and Cartilage*. 2014;22(7):918-27.
- Walker S, Hyde C, Hamilton W. Risk of breast cancer in symptomatic women in primary care: a case-control study using electronic records. *Br J Gen Pract*. 2014;64(629):e788-93.
- Walker AJ, West J, Card TR, Humes DJ, Grainge MJ. Variation in the risk of venous thromboembolism in people with colorectal cancer: a population-based cohort study from England. *J Thromb Haemost*. 2014;12(5):641-9.
- Vinogradova Y, Coupland C, Hippisley-Cox J. Exposure to combined oral contraceptives and risk of venous thromboembolism: a protocol for nested case-control studies using the QResearch and the CPRD databases. *BMJ Open*. 2014;4(4):e004499.

- van Staa T-P, Gulliford M, Ng ESW, Goldacre B, Smeeth L. Prediction of cardiovascular risk using Framingham, ASSIGN and QRISK2: how well do they predict individual rather than population risk? *PLoS One*. 2014;9(10):e106455.
- van Staa T-P, Dyson L, McCann G, Padmanabhan S, Belatri R, Goldacre B, et al. The opportunities and challenges of pragmatic point-of-care randomised trials using routinely collected electronic records: evaluations of two exemplar trials. *Health Technol Assess*. 2014;18(43):1-146.
- van Staa TP, Carr DF, O'Meara H, McCann G, Pirmohamed M. Predictors and outcomes of increases in creatine phosphokinase concentrations or rhabdomyolysis risk during statin treatment. *British Journal of Clinical Pharmacology*. 2014;78(3):649-59.
- Tsilidis KK, Capothanassi D, Allen NE, Rizos EC, Lopez DS, van Veldhoven K, et al. Metformin does not affect cancer risk: a cohort study in the U.K. *Clinical Practice Research Datalink analyzed like an intention-to-treat trial*. *Diabetes Care*. 2014;37(9):2522-32.
- Toovey S, Nieforth K, Smith P, Schlagenhaut P, Adamcova M, Tatt I, et al. Comparative benefit of malaria chemoprophylaxis modelled in United Kingdom travellers. *Travel Med Infect Dis*. 2014;12(6 Pt B):726-32.
- Thomas G, Khunti K, Curcin V, Molokhia M, Millett C, Majeed A, et al. Obesity paradox in people newly diagnosed with type 2 diabetes with and without prior cardiovascular disease. *Diabetes Obes Metab*. 2014;16(4):317-25.
- Taylor PN, Minassian C, Rehman A, Iqbal A, Draman MS, Hamilton W, et al. TSH levels and risk of miscarriage in women on long-term levothyroxine: a community-based study. *J Clin Endocrinol Metab*. 2014;99(10):3895-902.
- Taylor PN, Iqbal A, Minassian C, Sayers A, Draman MS, Greenwood R, et al. Falling threshold for treatment of borderline elevated thyrotropin levels-balancing benefits and risks: evidence from a large community-based study. *JAMA Intern Med*. 2014;174(1):32-9.
- Tate AR, Beloff N, Al-Radwan B, Wickson J, Puri S, Williams T, et al. Exploiting the potential of large databases of electronic health records for research using rapid search algorithms and an intuitive query interface. *J Am Med Inform Assoc*. 2014;21(2):292-8.
- Spoenclin J, Voegel JJ, Jick SS, Meier CR. Antihypertensive drugs and the risk of incident rosacea. *Br J Dermatol*. 2014;171(1):130-6.
- Spoenclin J, Bichsel F, Voegel JJ, Jick SS, Meier CR. The association between psychiatric diseases, psychotropic drugs and the risk of incident rosacea. *Br J Dermatol*. 2014;170(4):878-83.
- Søndergaard J, Halling A. The PROTECCT-M study: a cohort study investigating associations between novel specific biomarkers, patient-related, healthcare system markers and the trajectory of COPD patients treated in primary care. *BMC Pulm Med*. 2014;14:88.
- Siriwardena AN, Asghar Z, Coupland CCA. Influenza and pneumococcal vaccination and risk of stroke or transient ischaemic attack-matched case control study. *Vaccine*. 2014;32(12):1354-61.
- Scowcroft ACE, Cowie MR. Atrial fibrillation: improvement in identification and stroke preventive therapy - data from the UK Clinical Practice Research Datalink, 2000-2012. *Int J Cardiol*. 2014;171(2):169-73.
- Schneider C, Adamcova M, Jick SS, Schlagenhaut P, Miller MK, Rhein H-G, et al. Use of anti-malarial drugs and the risk of developing eye disorders. *Travel Med Infect Dis*. 2014;12(1):40-7.

- Ruigómez A, Brauer R, Rodríguez LAG, Huerta C, Requena G, Gil M, et al. Ascertainment of acute liver injury in two European primary care databases. *Eur J Clin Pharmacol*. 2014;70(10):1227-35.
- Richy FF, Sabidó-Espin M, Guedes S, Corvino FA, Gottwald-Hostalek U. Incidence of lactic acidosis in patients with type 2 diabetes with and without renal impairment treated with metformin: a retrospective cohort study. *Diabetes Care*. 2014;37(8):2291-5.
- Reinau D, Surber C, Jick SS, Meier CR. Epidemiology of basal cell carcinoma in the United Kingdom: incidence, lifestyle factors, and comorbidities. *Br J Cancer*. 2014;111(1):203-6.
- Reeves D, Springate DA, Ashcroft DM, Ryan R, Doran T, Morris R, et al. Can analyses of electronic patient records be independently and externally validated? The effect of statins on the mortality of patients with ischaemic heart disease: a cohort study with nested case-control analysis. *BMJ Open*. 2014;4(4):e004952.
- Redaniel MT, Gardner MP, Martin RM, Jeffreys M. The association of vitamin D supplementation with the risk of cancer in postmenopausal women. *Cancer Causes Control*. 2014;25(2):267-71.
- Ratib S, West J, Crooks CJ, Fleming KM. Diagnosis of liver cirrhosis in England, a cohort study, 1998-2009: a comparison with cancer. *Am J Gastroenterol*. 2014;109(2):190-8.
- Ratib S, Fleming KM, Crooks CJ, Aithal GP, West J. 1 and 5 year survival estimates for people with cirrhosis of the liver in England, 1998-2009: a large population study. *J Hepatol*. 2014;60(2):282-9.
- Quint JK, Müllerova H, DiSantostefano RL, Forbes H, Eaton S, Hurst JR, et al. Validation of chronic obstructive pulmonary disease recording in the Clinical Practice Research Datalink (CPRD-GOLD). *BMJ Open*. 2014;4(7):e005540.
- Puneekar YS, Shukla A, Müllerova H. COPD management costs according to the frequency of COPD exacerbations in UK primary care. *Int J Chron Obstruct Pulmon Dis*. 2014;9:65-73.
- Price SJ, Shephard EA, Stapley SA, Barraclough K, Hamilton WT. Non-visible versus visible haematuria and bladder cancer risk: a study of electronic records in primary care. *Br J Gen Pract*. 2014;64(626):e584-9.
- Price D, Thomas V, von Ziegenweidt J, Gould S, Hutton C, King C. Switching patients from other inhaled corticosteroid devices to the Easyhaler®: historical, matched-cohort study of real-life asthma patients. *J Asthma Allergy*. 2014;7:31-51.
- Prasad V, Kendrick D, Sayal K, Thomas SL, West J. Injury among children and young adults with epilepsy. *Pediatrics*. 2014;133(5):827-35.
- Pouwels S, de Boer A, Leufkens HGM, Weber WEJ, Cooper C, van Staa TP, et al. Risk of fracture in patients with Guillain-Barré syndrome. *Osteoporos Int*. 2014;25(7):1845-51.
- Pouwels S, de Boer A, Leufkens HGM, Weber WEJ, Cooper C, van Onzenoort HaW, et al. Risk of fracture in patients with muscular dystrophies. *Osteoporos Int*. 2014;25(2):509-18.
- Pouwels S, de Boer A, Leufkens HGM, Weber WEJ, Cooper C, de Vries F. Risk of fracture in patients with Charcot-Marie-Tooth disease. *Muscle Nerve*. 2014;50(6):919-24.
- Peeters PJHL, Bazelier MT, Uitdehaag BMJ, Leufkens HGM, De Bruin ML, de Vries F. The risk of venous thromboembolism in patients with multiple sclerosis: the Clinical Practice Research Datalink. *J Thromb Haemost*. 2014;12(4):444-51.

- Pauriah M, Elder DHJ, Ogston S, Noman AYS, Majeed A, Wyatt JC, et al. High-potency statin and ezetimibe use and mortality in survivors of an acute myocardial infarction: a population-based study. *Heart*. 2014;100(11):867-72.
- Palmer KT, D'Angelo S, Harris EC, Linaker C, Coggon D. The role of mental health problems and common psychotropic drug treatments in accidental injury at work: a case-control study. *Occup Environ Med*. 2014;71(5):308-12.
- Palmer KT, D'Angelo S, Harris EC, Linaker C, Coggon D. Epilepsy, diabetes mellitus and accidental injury at work. *Occup Med (Lond)*. 2014;64(6):448-53.
- O'Meara H, Carr DF, Evelyn J, Hobbs M, McCann G, van Staa T, et al. Electronic health records for biological sample collection: feasibility study of statin-induced myopathy using the Clinical Practice Research Datalink. *British Journal of Clinical Pharmacology*. 2014;77(5):831-8.
- Neal RD, Din NU, Hamilton W, Ukoumunne OC, Carter B, Stapley S, et al. Comparison of cancer diagnostic intervals before and after implementation of NICE guidelines: analysis of data from the UK General Practice Research Database. *Br J Cancer*. 2014;110(3):584-92.
- Murray LJ, Cooper JA, Hughes CM, Powe DG, Cardwell CR. Post-diagnostic prescriptions for low-dose aspirin and breast cancer-specific survival: a nested case-control study in a breast cancer cohort from the UK Clinical Practice Research Datalink. *Breast Cancer Res*. 2014;16(2):R34.
- Mundy LM, Sampson T, Logie JW. Retapamulin prescriptions and monitored off-label use. *Paediatr Drugs*. 2014;16(4):331-6.
- Müllerová H, Shukla A, Hawkins A, Quint J. Risk factors for acute exacerbations of COPD in a primary care population: a retrospective observational cohort study. *BMJ Open*. 2014;4(12):e006171.
- Müllerová H, Lu C, Li H, Tabberer M. Prevalence and burden of breathlessness in patients with chronic obstructive pulmonary disease managed in primary care. *PloS One*. 2014;9(1):e85540.
- Muller S, Hider SL, Belcher J, Helliwell T, Mallen CD. Is cancer associated with polymyalgia rheumatica? A cohort study in the General Practice Research Database. *Annals of the Rheumatic Diseases*. 2014;73(10):1769-73.
- Morgan E, Patterson CC, Cardwell CR. General practice-recorded depression and antidepressant use in young people with newly diagnosed Type 1 diabetes: a cohort study using the Clinical Practice Research Datalink. *Diabet Med*. 2014;31(2):241-5.
- Morgan CL, Puelles J, Poole CD, Currie CJ. The effect of withdrawal of rosiglitazone on treatment pathways, diabetes control and patient outcomes: a retrospective cohort study. *J Diabetes Complicat*. 2014;28(3):360-4.
- Morgan CL, Mukherjee J, Jenkins-Jones S, Holden SE, Currie CJ. Combination therapy with metformin plus sulphonylureas versus metformin plus DPP-4 inhibitors: association with major adverse cardiovascular events and all-cause mortality. *Diabetes Obes Metab*. 2014;16(10):977-83.
- Morgan CL, Mukherjee J, Jenkins-Jones S, Holden SE, Currie CJ. Association between first-line monotherapy with sulphonylurea versus metformin and risk of all-cause mortality and cardiovascular events: a retrospective, observational study. *Diabetes Obes Metab*. 2014;16(10):957-62.
- Moore A, Harnden A, Mayon-White R. Recognising Kawasaki disease in UK primary care: a descriptive study using the Clinical Practice Research Datalink. *Br J Gen Pract*. 2014;64(625):e477-83.

- Menon S, Nightingale P, Trudgill N. Is hormone replacement therapy in post-menopausal women associated with a reduced risk of oesophageal cancer? *United European Gastroenterol J.* 2014;2(5):374-82.
- McGrogan A, Snowball J, de Vries CS. Pregnancy losses in women with Type 1 or Type 2 diabetes in the UK: an investigation using primary care records. *Diabet Med.* 2014;31(3):357-65.
- McDonald HI, Nitsch D, Millett ERC, Sinclair A, Thomas SL. New estimates of the burden of acute community-acquired infections among older people with diabetes mellitus: a retrospective cohort study using linked electronic health records. *Diabet Med.* 2014;31(5):606-14.
- McCourt C, Coleman HG, Murray LJ, Cantwell MM, Dolan O, Powe DG, et al. Beta-blocker usage after malignant melanoma diagnosis and survival: a population-based nested case-control study. *Br J Dermatol.* 2014;170(4):930-8.
- Matok I, Azoulay L, Yin H, Suissa S. Immortal time bias in observational studies of drug effects in pregnancy. *Birth Defects Res Part A Clin Mol Teratol.* 2014;100(9):658-62.
- Mathur R, Bhaskaran K, Chaturvedi N, Leon DA, vanStaa T, Grundy E, et al. Completeness and usability of ethnicity data in UK-based primary care and hospital databases. *J Public Health (Oxf).* 2014;36(4):684-92.
- Matcho A, Ryan P, Fife D, Reich C. Fidelity assessment of a clinical practice research datalink conversion to the OMOP common data model. *Drug Safety.* 2014;37(11):945-59.
- Martinez C, Katholing A, Freedman SB. Adverse prognosis of incidentally detected ambulatory atrial fibrillation. A cohort study. *Thromb Haemost.* 2014;112(2):276-86.
- Martinez C, Cohen AT, Bamber L, Rietbrock S. Epidemiology of first and recurrent venous thromboembolism: a population-based cohort study in patients without active cancer. *Thromb Haemost.* 2014;112(2):255-63.
- Margulis AV, Kang EM, Hammad TA. Patterns of prescription of antidepressants and antipsychotics across and within pregnancies in a population-based UK cohort. *Matern Child Health J.* 2014;18(7):1742-52.
- Makar GA, Holmes JH, Yang Y-X. Angiotensin-converting enzyme inhibitor therapy and colorectal cancer risk. *J Natl Cancer Inst.* 2014;106(2):djt374.
- Maguire A, Mitchell BD, Ruzafa JC. Antihyperglycaemic treatment patterns, observed glycaemic control and determinants of treatment change among patients with type 2 diabetes in the United Kingdom primary care: a retrospective cohort study. *BMC Endocr Disord.* 2014;14:73.
- Mackenzie IS, Morant SV, Bloomfield GA, MacDonald TM, O'Riordan J. Incidence and prevalence of multiple sclerosis in the UK 1990-2010: a descriptive study in the General Practice Research Database. *J Neurol Neurosurg Psychiatr.* 2014;85(1):76-84.
- Macedo AF, Douglas I, Smeeth L, Forbes H, Ebrahim S. Statins and the risk of type 2 diabetes mellitus: cohort study using the UK clinical practice research datalink. *BMC Cardiovasc Disord.* 2014;14:85.
- Lipscombe LL, Austin PC, Alessi-Severini S, Blackburn DF, Blais L, Bresee L, et al. Atypical antipsychotics and hyperglycemic emergencies: multicentre, retrospective cohort study of administrative data. *Schizophr Res.* 2014;154(1-3):54-60.
- Liang H, Vallarino C, Joseph G, Manne S, Perez A, Zhang S. Increased risk of subsequent myocardial infarction in patients with type 2 diabetes: a retrospective cohort study using the U.K. General Practice Research Database. *Diabetes Care.* 2014;37(5):1329-37.

- Li L, Setoguchi S, Jick S. Validity of non-fatal idiopathic serious ventricular arrhythmia in UK-based Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf.* 2014;23(9):999-1001.
- Li L, Roddam A, Ferguson S, Feudjo-Tepie M, Taylor A, Jick S. Switch patterns of osteoporosis medication and its impact on persistence among postmenopausal women in the U.K. *General Practice Research Database. Menopause.* 2014;21(10):1106-13.
- Li L, Ambegaonkar BM, Reckless JPD, Jick S. Association of a reduction in low-density lipoprotein cholesterol with incident cardiovascular and cerebrovascular events among people with type 2 diabetes mellitus. *Eur J Prev Cardiol.* 2014;21(7):855-65.
- Langan SM, Minassian C, Smeeth L, Thomas SL. Risk of stroke following herpes zoster: a self-controlled case-series study. *Clin Infect Dis.* 2014;58(11):1497-503.
- Lambrelli D, Burge R, Raluy-Callado M, Chen S-Y, Wu N, Schoenfeld MJ. Retrospective database study to assess the economic impact of hip fracture in the United Kingdom. *J Med Econ.* 2014;17(11):817-25.
- Kontopantelis E, Springate D, Reeves D, Ashcroft DM, Valderas JM, Doran T. Withdrawing performance indicators: retrospective analysis of general practice performance under UK Quality and Outcomes Framework. *BMJ.* 2014;348:g330.
- Klop C, Welsing PMJ, Cooper C, Harvey NC, Elders PJM, Bijlsma JWJ, et al. Mortality in British hip fracture patients, 2000-2010: a population-based retrospective cohort study. *Bone.* 2014;66:171-7.
- Khalid JM, Raluy-Callado M, Curtis BH, Boye KS, Maguire A, Reaney M. Rates and risk of hospitalisation among patients with type 2 diabetes: retrospective cohort study using the UK General Practice Research Database linked to English Hospital Episode Statistics. *Int J Clin Pract.* 2014;68(1):40-8.
- Jones RCM, Price D, Ryan D, Sims EJ, von Ziegenweidt J, Mascarenhas L, et al. Opportunities to diagnose chronic obstructive pulmonary disease in routine care in the UK: a retrospective study of a clinical cohort. *Lancet Respir Med.* 2014;2(4):267-76.
- Jick SS, Li L, Falcone GJ, Vassilev ZP, Wallander MA. Mortality of patients with multiple sclerosis: a cohort study in UK primary care. *J Neurol.* 2014;261(8):1508-17.
- Jameson K, Jick S, Hagberg KW, Ambegaonkar B, Giles A, O'Donoghue D. Prevalence and management of chronic kidney disease in primary care patients in the UK. *Int J Clin Pract.* 2014;68(9):1110-21.
- Ishiguro C, Wang X, Li L, Jick S. Antipsychotic drugs and risk of idiopathic venous thromboembolism: a nested case-control study using the CPRD. *Pharmacoepidemiol Drug Saf.* 2014;23(11):1168-75.
- Huiart L, Ferdynus C, Dell'Aniello S, Bakiri N, Giorgi R, Suissa S. Measuring persistence to hormonal therapy in patients with breast cancer: accounting for temporary treatment discontinuation. *Pharmacoepidemiol Drug Saf.* 2014;23(8):882-9.
- Holden SE, Gale EaM, Jenkins-Jones S, Currie CJ. How many people inject insulin? UK estimates from 1991 to 2010. *Diabetes Obes Metab.* 2014;16(6):553-9.
- Herrett E, van Staa T, Free C, Smeeth L. Text messaging reminders for influenza vaccine in primary care: protocol for a cluster randomised controlled trial (TXT4FLUJAB). *BMJ Open.* 2014;4(5):e004633.
- Hazra N, Gulliford M. Evaluating pancreatitis in primary care: a population-based cohort study. *Br J Gen Pract.* 2014;64(622):e295-301.

- Hassan L, Senior J, Frisher M, Edge D, Shaw J. A comparison of psychotropic medication prescribing patterns in East of England prisons and the general population. *Journal of Psychopharmacology (Oxford, England)*. 2014;28(4):357-62.
- Hancock AT, Mallen CD, Muller S, Belcher J, Roddy E, Helliwell T, et al. Risk of vascular events in patients with polymyalgia rheumatica. *CMAJ*. 2014;186(13):E495-501.
- Hagberg KW, McGlynn KA, Sahasrabudde VV, Jick S. Anti-diabetic medications and risk of primary liver cancer in persons with type II diabetes. *Br J Cancer*. 2014;111(9):1710-7.
- Gulliford MC, van Staa TP, McDermott L, McCann G, Charlton J, Dregan A, et al. Cluster randomized trials utilizing primary care electronic health records: methodological issues in design, conduct, and analysis (eCRT Study). *Trials*. 2014;15:220.
- Gulliford MC, van Staa T, Dregan A, McDermott L, McCann G, Ashworth M, et al. Electronic health records for intervention research: a cluster randomized trial to reduce antibiotic prescribing in primary care (eCRT study). *Annals of family medicine*. 2014;12(4):344-51.
- Gulliford MC, Dregan A, Moore MV, Ashworth M, Staa Tv, McCann G, et al. Continued high rates of antibiotic prescribing to adults with respiratory tract infection: survey of 568 UK general practices. *BMJ Open*. 2014;4(10):e006245.
- Gulliford MC, Charlton J, Bhattarai N, Charlton C, Rudisill C. Impact and cost-effectiveness of a universal strategy to promote physical activity in primary care: population-based cohort study and Markov model. *Eur J Health Econ*. 2014;15(4):341-51.
- Gulliford MC, Bhattarai N, Charlton J, Rudisill C. Cost-effectiveness of a universal strategy of brief dietary intervention for primary prevention in primary care: population-based cohort study and Markov model. *Cost Eff Resour Alloc*. 2014;12(1):4.
- Gulliford M, Charlton J, Bhattarai N, Rudisill C. Social and material deprivation and the cost-effectiveness of an intervention to promote physical activity: cohort study and Markov model. *J Public Health (Oxf)*. 2014;36(4):674-83.
- Gibson-Smith D, Klop C, Elders PJM, Welsing PMJ, van Schoor N, Leufkens HGM, et al. The risk of major and any (non-hip) fragility fracture after hip fracture in the United Kingdom: 2000-2010. *Osteoporos Int*. 2014;25(11):2555-63.
- Gao W, Gulliford M, Bennett MI, Murtagh FEM, Higginson IJ. Managing cancer pain at the end of life with multiple strong opioids: a population-based retrospective cohort study in primary care. *PloS One*. 2014;9(1):e79266.
- Gallagher AM, van Staa TP, Murray-Thomas T, Schoof N, Clemens A, Ackermann D, et al. Population-based cohort study of warfarin-treated patients with atrial fibrillation: incidence of cardiovascular and bleeding outcomes. *BMJ Open*. 2014;4(1):e003839.
- Gadoud A, Kane E, Macleod U, Ansell P, Oliver S, Johnson M. Palliative care among heart failure patients in primary care: a comparison to cancer patients using English family practice data. *PloS One*. 2014;9(11):e113188.
- Furlano RI, Bloechliger M, Jick H, Meier CR. Bone fractures in children with autistic spectrum disorder. *J Dev Behav Pediatr*. 2014;35(6):353-9.

- Fox BD, Azoulay L, Dell'Aniello S, Langleben D, Lapi F, Benisty J, et al. The use of antidepressants and the risk of idiopathic pulmonary arterial hypertension. *Can J Cardiol.* 2014;30(12):1633-9.
- Fournier J-P, Yin H, Yu OHY, Azoulay L. Metformin and low levels of thyroid-stimulating hormone in patients with type 2 diabetes mellitus. *CMAJ.* 2014;186(15):1138-45.
- Fournier J-P, Duijnhoven RG, Renoux C, Dell'Aniello S, Klungel OH, Suissa S. Concurrent use of statins and hormone therapy and risk of venous thromboembolism in postmenopausal women: a population-based case-control study. *Menopause.* 2014;21(9):1023-6.
- Forbes HJ, Bhaskaran K, Thomas SL, Smeeth L, Clayton T, Langan SM. Quantification of risk factors for herpes zoster: population based case-control study. *BMJ.* 2014;348:g2911.
- Filion KB, Chateau D, Targownik LE, Gershon A, Durand M, Tamim H, et al. Proton pump inhibitors and the risk of hospitalisation for community-acquired pneumonia: replicated cohort studies with meta-analysis. *Gut.* 2014;63(4):552-8.
- Faillie J-L, Azoulay L, Patenaude V, Hillaire-Buys D, Suissa S. Incretin based drugs and risk of acute pancreatitis in patients with type 2 diabetes: cohort study. *BMJ.* 2014;348:g2780.
- Eppenga WL, Lalmohamed A, Geerts AF, Derijks HJ, Wensing M, Egberts A, et al. Risk of lactic acidosis or elevated lactate concentrations in metformin users with renal impairment: a population-based cohort study. *Diabetes Care.* 2014;37(8):2218-24.
- Duijnhoven RG, Straus SMJM, Souverein PC, de Boer A, Bosch JLHR, Hoes AW, et al. Long-term use of 5 α -reductase inhibitors and the risk of male breast cancer. *Cancer Causes Control.* 2014;25(11):1577-82.
- Driessen JHM, van Onzenoort HAW, Henry RMA, Lalmohamed A, van den Bergh JP, Neef C, et al. Use of dipeptidyl peptidase-4 inhibitors for type 2 diabetes mellitus and risk of fracture. *Bone.* 2014;68:124-30.
- Dregan A, van Staa TP, McDermott L, McCann G, Ashworth M, Charlton J, et al. Point-of-care cluster randomized trial in stroke secondary prevention using electronic health records. *Stroke.* 2014;45(7):2066-71.
- Dregan A, Charlton J, Chowienczyk P, Gulliford MC. Chronic inflammatory disorders and risk of type 2 diabetes mellitus, coronary heart disease, and stroke: a population-based cohort study. *Circulation.* 2014;130(10):837-44.
- Donegan K, King B, Bryan P. Safety of pertussis vaccination in pregnant women in UK: observational study. *BMJ.* 2014;349:g4219.
- Dial S, Nessim SJ, Kezouh A, Benisty J, Suissa S. Antihypertensive agents acting on the renin-angiotensin system and the risk of sepsis. *British Journal of Clinical Pharmacology.* 2014;78(5):1151-8.
- Das R, Langerman H, D'Oca K, Strongman H. An Observational Cohort Study of Diabetes-Associated Secondary Health Care Utilisation in Patients With Type 2 Diabetes Prescribed Dual Combination Therapy With Oral Anti-Hyperglycaemic Agents in the UK. *Value Health.* 2014;17(7):A360.
- Currie CJ, Berni E, Jenkins-Jones S, Poole CD, Ouwens M, Driessen S, et al. Antibiotic treatment failure in four common infections in UK primary care 1991-2012: longitudinal analysis. *BMJ.* 2014;349:g5493.
- Couraud S, Dell'Aniello S, Bouganim N, Azoulay L. Cardiac glycosides and the risk of breast cancer in women with chronic heart failure and supraventricular arrhythmia. *Breast Cancer Res Treat.* 2014;146(3):619-26.

- Couraud S, Azoulay L, Dell'Aniello S, Suissa S. Cardiac glycosides use and the risk of lung cancer: a nested case-control study. *BMC cancer*. 2014;14:573.
- Cornish RP, Henderson J, Boyd AW, Granell R, Van Staa T, Macleod J. Validating childhood asthma in an epidemiological study using linked electronic patient records. *BMJ Open*. 2014;4(4):e005345.
- Cooper C, Fox KM, Borer JS. Ischaemic cardiac events and use of strontium ranelate in postmenopausal osteoporosis: a nested case-control study in the CPRD. *Osteoporos Int*. 2014;25(2):737-45.
- Close H, Reilly J, Mason JM, Kripalani M, Wilson D, Main J, et al. Renal failure in lithium-treated bipolar disorder: a retrospective cohort study. *PLoS One*. 2014;9(3):e90169.
- Chen L, Reed C, Happich M, Nyhuis A, Lenox-Smith A. Health care resource utilisation in primary care prior to and after a diagnosis of Alzheimer's disease: a retrospective, matched case-control study in the United Kingdom. *BMC Geriatr*. 2014;14:76.
- Charlton R, Snowball J, Sammon C, de Vries C. The Clinical Practice Research Datalink for drug safety in pregnancy research: an overview. *Therapie*. 2014;69(1):83-9.
- Cardwell CR, Mc Menamin ÚC, Hicks BM, Hughes C, Cantwell MM, Murray LJ. Drugs affecting the renin-angiotensin system and survival from cancer: a population based study of breast, colorectal and prostate cancer patient cohorts. *BMC Med*. 2014;12:28.
- Cardwell CR, Kunzmann AT, Cantwell MM, Hughes C, Baron JA, Powe DG, et al. Low-dose aspirin use after diagnosis of colorectal cancer does not increase survival: a case-control analysis of a population-based cohort. *Gastroenterology*. 2014;146(3):700-8.e2.
- Cardwell CR, Hicks BM, Hughes C, Murray LJ. Statin use after colorectal cancer diagnosis and survival: a population-based cohort study. *J Clin Oncol*. 2014;32(28):3177-83.
- Cardwell CR, Flahavan EM, Hughes CM, Coleman HG, O'Sullivan JM, Powe DG, et al. Low-dose aspirin and survival in men with prostate cancer: a study using the UK Clinical Practice Research Datalink. *Cancer Causes Control*. 2014;25(1):33-43.
- Card TR, Siffledeen J, Fleming KM. Are IBD patients more likely to have a prior diagnosis of irritable bowel syndrome? Report of a case-control study in the General Practice Research Database. *United European Gastroenterol J*. 2014;2(6):505-12.
- Bultink IEM, Harvey NC, Lalmohamed A, Cooper C, Lems WF, van Staa TP, et al. Elevated risk of clinical fractures and associated risk factors in patients with systemic lupus erythematosus versus matched controls: a population-based study in the United Kingdom. *Osteoporos Int*. 2014;25(4):1275-83.
- Bruderer SG, Bodmer M, Jick SS, Bader G, Schlienger RG, Meier CR. Incidence of and risk factors for severe hypoglycaemia in treated type 2 diabetes mellitus patients in the UK--a nested case-control analysis. *Diabetes Obes Metab*. 2014;16(9):801-11.
- Bruderer S, Bodmer M, Jick SS, Meier CR. Use of diuretics and risk of incident gout: a population-based case-control study. *Arthritis & Rheumatology (Hoboken, NJ)*. 2014;66(1):185-96.
- Bradley MC, Motterlini N, Padmanabhan S, Cahir C, Williams T, Fahey T, et al. Potentially inappropriate prescribing among older people in the United Kingdom. *BMC Geriatr*. 2014;14:72.
- Booth HP, Prevost AT, Gulliford MC. Impact of body mass index on prevalence of multimorbidity in primary

care: cohort study. *Fam Pract*. 2014;31(1):38-43.

- Booth H, Khan O, Prevost T, Reddy M, Dregan A, Charlton J, et al. Incidence of type 2 diabetes after bariatric surgery: population-based matched cohort study. *Lancet Diabetes Endocrinol*. 2014;2(12):963-8.
- Boggon R, Timmis A, Hemingway H, Raju S, Malvestiti FM, Van Staa TP. Smoking cessation interventions following acute coronary syndrome: a missed opportunity? *Eur J Prev Cardiol*. 2014;21(6):767-73.
- Bloechliger M, Schlagenhaut P, Toovey S, Schnetzler G, Tatt I, Tomianovic D, et al. Malaria chemoprophylaxis regimens: a descriptive drug utilization study. *Travel Med Infect Dis*. 2014;12(6 Pt B):718-25.
- Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet*. 2014;384(9945):755-65.
- Bensimon L, Yin H, Suissa S, Pollak MN, Azoulay L. Type 2 diabetes and the risk of mortality among patients with prostate cancer. *Cancer Causes Control*. 2014;25(3):329-38.
- Becker C, Jick SS, Meier CR, Bodmer M. Metformin and the risk of head and neck cancer: a case-control analysis. *Diabetes Obes Metab*. 2014;16(11):1148-54.
- Beattie CJ, Fulton RL, Higgins P, Padmanabhan S, McCallum L, Walters MR, et al. Allopurinol initiation and change in blood pressure in older adults with hypertension. *Hypertension*. 2014;64(5):1102-7.
- Bannister CA, Poole CD, Jenkins-Jones S, Morgan CL, Elwyn G, Spasic I, et al. External validation of the UKPDS risk engine in incident type 2 diabetes: a need for new type 2 diabetes-specific risk equations. *Diabetes Care*. 2014;37(2):537-45.
- Bannister CA, Holden SE, Jenkins-Jones S, Morgan CL, Halcox JP, Schernthaner G, et al. Can people with type 2 diabetes live longer than those without? A comparison of mortality in people initiated with metformin or sulphonylurea monotherapy and matched, non-diabetic controls. *Diabetes Obes Metab*. 2014;16(11):1165-73.
- Azoulay L, Dell'Aniello S, Simon TA, Renoux C, Suissa S. Initiation of warfarin in patients with atrial fibrillation: early effects on ischaemic strokes. *Eur Heart J*. 2014;35(28):1881-7.
- Amirthalingam G, Andrews N, Campbell H, Ribeiro S, Kara E, Donegan K, et al. Effectiveness of maternal pertussis vaccination in England: an observational study. *Lancet*. 2014;384(9953):1521-8.
- Alexandre L, Clark AB, Bhutta HY, Holt S, Lewis MPN, Hart AR. Statin use is associated with reduced risk of histologic subtypes of esophageal cancer: a nested case-control analysis. *Gastroenterology*. 2014;146(3):661-8.
- Abdul Sultan A, Tata LJ, Fleming KM, Crooks CJ, Ludvigsson JF, Dhalwani NN, et al. Pregnancy complications and adverse birth outcomes among women with celiac disease: a population-based study from England. *Am J Gastroenterol*. 2014;109(10):1653-61.
- Abbing-Karahagopian V, Kurz X, de Vries F, van Staa TP, Alvarez Y, Hesse U, et al. Bridging differences in outcomes of pharmacoepidemiological studies: design and first results of the PROTECT project. *Curr Clin Pharmacol*. 2014;9(2):130-8.
- Abbing-Karahagopian V, Huerta C, Souverein PC, de Abajo F, Leufkens HGM, Slattery J, et al. Antidepressant prescribing in five European countries: application of common definitions to assess the prevalence, clinical observations, and methodological implications. *Eur J Clin Pharmacol*. 2014;70(7):849-57.

- Wilson JC, O'Rourke MA, Cooper JA, Murray LJ, Hughes CM, Gormley GJ, et al. Non-steroidal anti-inflammatory drug use and cervical cancer risk: a case-control study using the Clinical Practice Research Datalink. *Cancer Epidemiol.* 2013;37(6):897-904.
- Wei L, MacDonald TM, Mackenzie IS. Pioglitazone and bladder cancer: a propensity score matched cohort study. *British Journal of Clinical Pharmacology.* 2013;75(1):254-9.
- Walker S, Hyde C, Hamilton W. Risk of uterine cancer in symptomatic women in primary care: case-control study using electronic records. *Br J Gen Pract.* 2013;63(614):e643-8.
- Walker AJ, Card TR, West J, Crooks C, Grainge MJ. Incidence of venous thromboembolism in patients with cancer - a cohort study using linked United Kingdom databases. *Eur J Cancer.* 2013;49(6):1404-13.
- Vinogradova Y, Coupland C, Hippisley-Cox J. Exposure to bisphosphonates and risk of common non-gastrointestinal cancers: series of nested case-control studies using two primary-care databases. *Br J Cancer.* 2013;109(3):795-806.
- Vinogradova Y, Coupland C, Hippisley-Cox J. Exposure to bisphosphonates and risk of gastrointestinal cancers: series of nested case-control studies with QResearch and CPRD data. *BMJ.* 2013;346:f114.
- Vasilakis-Scaramozza C, Aschengrau A, Cabral HJ, Jick SS. Antihypertensive drugs and the risk of congenital anomalies. *Pharmacotherapy.* 2013;33(5):476-82.
- Vasilakis-Scaramozza C, Aschengrau A, Cabral HJ, Jick SS. Asthma drugs and the risk of congenital anomalies. *Pharmacotherapy.* 2013;33(4):363-8.
- Vasilakis-Scaramozza C, Aschengrau A, Cabral H, Jick SS. Antidepressant use during early pregnancy and the risk of congenital anomalies. *Pharmacotherapy.* 2013;33(7):693-700.
- Varyani FK, West J, Card TR. Primary biliary cirrhosis does not increase the risk of UTIs following diagnosis compared to other chronic liver diseases? *Liver Int.* 2013;33(3):384-8.
- van Staa T-P, Smeeth L, Ng ESW, Goldacre B, Gulliford M. The efficiency of cardiovascular risk assessment: do the right patients get statin treatment? *Heart.* 2013;99(21):1597-602.
- Van Den Ham HA, Klungel OH, Leufkens HGM, Van Staa TP. The patterns of anticoagulation control and the risk of stroke, bleeding and mortality in patients with non-valvular atrial fibrillation. *J Thromb Haemost.* 2013;11(1):107-15.
- Tsang C, Bottle A, Majeed A, Aylin P. Adverse events recorded in English primary care: observational study using the General Practice Research Database. *Br J Gen Pract.* 2013;63(613):e534-42.
- Tsang C, Bottle A, Majeed A, Aylin P. Cancer diagnosed by emergency admission in England: an observational study using the general practice research database. *BMC Health Serv Res.* 2013;13:308.
- Thomas KH, Martin RM, Davies NM, Metcalfe C, Windmeijer F, Gunnell D. Smoking cessation treatment and risk of depression, suicide, and self harm in the Clinical Practice Research Datalink: prospective cohort study. *BMJ.* 2013;347:f5704.
- Thomas KH, Davies N, Metcalfe C, Windmeijer F, Martin RM, Gunnell D. Validation of suicide and self-harm records in the Clinical Practice Research Datalink. *British Journal of Clinical Pharmacology.* 2013;76(1):145-57.
- Taylor KS, Heneghan CJ, Farmer AJ, Fuller AM, Adler AI, Aronson JK, et al. All-cause and cardiovascular

mortality in middle-aged people with type 2 diabetes compared with people without diabetes in a large U.K. primary care database. *Diabetes Care*. 2013;36(8):2366-71.

- Taylor B, Jick H, Maclaughlin D. Prevalence and incidence rates of autism in the UK: time trend from 2004-2010 in children aged 8 years. *BMJ Open*. 2013;3(10):e003219.
- Spoenclin J, Voegel JJ, Jick SS, Meier CR. Migraine, triptans, and the risk of developing rosacea: a population-based study within the United Kingdom. *J Am Acad Dermatol*. 2013;69(3):399-406.
- Solaymani-Dodaran M, Card TR, West J. Cause-specific mortality of people with Barrett's esophagus compared with the general population: a population-based cohort study. *Gastroenterology*. 2013;144(7):1375-83, 83.e1.
- Smiechowski BB, Azoulay L, Yin H, Pollak MN, Suissa S. The use of metformin and the incidence of lung cancer in patients with type 2 diabetes. *Diabetes Care*. 2013;36(1):124-9.
- Smiechowski B, Azoulay L, Yin H, Pollak MN, Suissa S. The use of metformin and colorectal cancer incidence in patients with type II diabetes mellitus. *Cancer Epidemiol Biomarkers Prev*. 2013;22(10):1877-83.
- Skow A, Douglas I, Smeeth L. The association between Parkinson's disease and anti-epilepsy drug carbamazepine: a case-control study using the UK General Practice Research Database. *British Journal of Clinical Pharmacology*. 2013;76(5):816-22.
- Shephard E, Neal R, Rose P, Walter F, Hamilton WT. Clinical features of kidney cancer in primary care: a case-control study using primary care records. *Br J Gen Pract*. 2013;63(609):e250-5.
- Scowcroft ACE, Lee S, Mant J. Thromboprophylaxis of elderly patients with AF in the UK: an analysis using the General Practice Research Database (GPRD) 2000-2009. *Heart*. 2013;99(2):127-32.
- Schneider C, Adamcova M, Jick SS, Schlagenhauf P, Miller MK, Rhein H-G, et al. Antimalarial chemoprophylaxis and the risk of neuropsychiatric disorders. *Travel Med Infect Dis*. 2013;11(2):71-80.
- Sammon CJ, McGrogan A, Snowball J, de Vries CS. Pandemic influenza vaccination during pregnancy: an investigation of vaccine uptake during the 2009/10 pandemic vaccination campaign in Great Britain. *Hum Vaccin Immunother*. 2013;9(4):917-23.
- Refoios Camejo R, McGrath C, Miraldo M, Rutten F. The determinants of cost-effectiveness potential: an historical perspective on lipid-lowering therapies. *Pharmacoeconomics*. 2013;31(5):445-54.
- Reed C, Hong J, Novick D, Lenox-Smith A, Happich M. Health care costs before and after diagnosis of depression in patients with unexplained pain: a retrospective cohort study using the United Kingdom General Practice Research Database. *Clinicoecon Outcomes Res*. 2013;5:37-47.
- Quint JK, Herrett E, Bhaskaran K, Timmis A, Hemingway H, Wedzicha JA, et al. Effect of β blockers on mortality after myocardial infarction in adults with COPD: population based cohort study of UK electronic healthcare records. *BMJ*. 2013;347:f6650.
- Qiu H, Rhoads GG, Berlin JA, Marcella SW, Demissie K. Initial metformin or sulphonylurea exposure and cancer occurrence among patients with type 2 diabetes mellitus. *Diabetes Obes Metab*. 2013;15(4):349-57.
- Pouwels S, de Boer A, Javaid MK, Hilton-Jones D, Verschuur J, Cooper C, et al. Fracture rate in patients with myasthenia gravis: the general practice research database. *Osteoporos Int*. 2013;24(2):467-76.

- Pouwels S, Bazelier MT, de Boer A, Weber WEJ, Neef CK, Cooper C, et al. Five-year fracture risk estimation in patients with Parkinson's disease. *Bone*. 2013;56(2):266-70.
- Pouwels S, Bazelier MT, de Boer A, Weber WEJ, Neef C, Cooper C, et al. Risk of fracture in patients with Parkinson's disease. *Osteoporos Int*. 2013;24(8):2283-90.
- Poole CD, Halcox JP, Jenkins-Jones S, Carr ESM, Schifflers MG, Ray KK, et al. Omega-3 Fatty acids and mortality outcome in patients with and without type 2 diabetes after myocardial infarction: a retrospective, matched-cohort study. *Clin Ther*. 2013;35(1):40-51.
- Padwal RS, Klarenbach SW, Wang X, Sharma AM, Karmali S, Birch DW, et al. A simple prediction rule for all-cause mortality in a cohort eligible for bariatric surgery. *JAMA Surg*. 2013;148(12):1109-15.
- Norton S, Koduri G, Nikiphorou E, Dixey J, Williams P, Young A. A study of baseline prevalence and cumulative incidence of comorbidity and extra-articular manifestations in RA and their impact on outcome. *Rheumatology (Oxford)*. 2013;52(1):99-110.
- Nicholson A, Ford E, Davies KA, Smith HE, Rait G, Tate AR, et al. Optimising use of electronic health records to describe the presentation of rheumatoid arthritis in primary care: a strategy for developing code lists. *PLoS One*. 2013;8(2):e54878.
- Nicholas JM, Ridsdale L, Richardson MP, Grieve AP, Gulliford MC. Fracture risk with use of liver enzyme inducing antiepileptic drugs in people with active epilepsy: cohort study using the general practice research database. *Seizure*. 2013;22(1):37-42.
- Murray ML, Insuk S, Banaschewski T, Neubert AC, McCarthy S, Buitelaar JK, et al. An inventory of European data sources for the long-term safety evaluation of methylphenidate. *Eur Child Adolesc Psychiatry*. 2013;22(10):605-18.
- Morgan CL, Owens DR, Aubonnet P, Carr ESM, Jenkins-Jones S, Poole CD, et al. Primary prevention of diabetic retinopathy with fibrates: a retrospective, matched cohort study. *BMJ Open*. 2013;3(12):e004025.
- Morales DR, Donnan PT, Daly F, Staa TV, Sullivan FM. Impact of clinical trial findings on Bell's palsy management in general practice in the UK 2001-2012: interrupted time series regression analysis. *BMJ Open*. 2013;3(7).
- Minassian C, Thomas SL, Williams DJ, Campbell O, Smeeth L. Acute maternal infection and risk of pre-eclampsia: a population-based case-control study. *PLoS One*. 2013;8(9):e73047.
- Micali N, Hagberg KW, Petersen I, Treasure JL. The incidence of eating disorders in the UK in 2000-2009: findings from the General Practice Research Database. *BMJ Open*. 2013;3(5).
- Martinez C, Jones RW, Rietbrock S. Trends in the prevalence of antipsychotic drug use among patients with Alzheimer's disease and other dementias including those treated with antidementia drugs in the community in the UK: a cohort study. *BMJ Open*. 2013;3(1).
- Margulis AV, Abou-Ali A, Strazzeri MM, Ding Y, Kuyateh F, Frimpong EY, et al. Use of selective serotonin reuptake inhibitors in pregnancy and cardiac malformations: a propensity-score matched cohort in CPRD. *Pharmacoepidemiol Drug Saf*. 2013;22(9):942-51.
- Lopez Bernal JA, Upton MN, Henderson AJ, Dedman D, McCarthy A, Davey Smith G, et al. Lower respiratory tract infection in the first year of life is associated with worse lung function in adult life: prospective results from the Barry Caerphilly Growth study. *Ann Epidemiol*. 2013;23(7):422-7.

- Li L, Setoguchi S, Cabral H, Jick S. Opioid use for noncancer pain and risk of fracture in adults: a nested case-control study using the general practice research database. *Am J Epidemiol.* 2013;178(4):559-69.
- Li L, Setoguchi S, Cabral H, Jick S. Opioid use for noncancer pain and risk of myocardial infarction amongst adults. *J Intern Med.* 2013;273(5):511-26.
- Li L, Setoguchi S, Cabral H, Jick S. Opioids and risk of type 2 diabetes in adults with non-cancer pain. *Pain Physician.* 2013;16(1):77-88.
- Lapi F, Azoulay L, Yin H, Nessim SJ, Suissa S. Concurrent use of diuretics, angiotensin converting enzyme inhibitors, and angiotensin receptor blockers with non-steroidal anti-inflammatory drugs and risk of acute kidney injury: nested case-control study. *BMJ.* 2013;346:e8525.
- Lapi F, Azoulay L, Niazi MT, Yin H, Benayoun S, Suissa S. Androgen deprivation therapy and risk of acute kidney injury in patients with prostate cancer. *JAMA.* 2013;310(3):289-96.
- Lanza LL, McQuay LJ, Rothman KJ, Bone HG, Kaunitz AM, Harel Z, et al. Use of depot medroxyprogesterone acetate contraception and incidence of bone fracture. *Obstet Gynecol.* 2013;121(3):593-600.
- Lalmohamed A, MacGregor AJ, de Vries F, Leufkens HGM, van Staa TP. Patterns of risk of cancer in patients with metal-on-metal hip replacements versus other bearing surface types: a record linkage study between a prospective joint registry and general practice electronic health records in England. *PloS One.* 2013;8(7):e65891.
- Kontopantelis E, Reeves D, Valderas JM, Campbell S, Doran T. Recorded quality of primary care for patients with diabetes in England before and after the introduction of a financial incentive scheme: a longitudinal observational study. *BMJ Qual Saf.* 2013;22(1):53-64.
- Khunti K, Wolden ML, Thorsted BL, Andersen M, Davies MJ. Clinical inertia in people with type 2 diabetes: a retrospective cohort study of more than 80,000 people. *Diabetes Care.* 2013;36(11):3411-7.
- Kehoe PG, Davies NM, Martin RM, Ben-Shlomo Y. Associations of angiotensin targeting antihypertensive drugs with mortality and hospitalization in primary care patients with dementia. *J Alzheimers Dis.* 2013;33(4):999-1008.
- Kalra PR, Morley C, Barnes S, Menown I, Kassianos G, Padmanabhan S, et al. Discontinuation of beta-blockers in cardiovascular disease: UK primary care cohort study. *Int J Cardiol.* 2013;167(6):2695-9.
- Kadam UT, Blagojevic M, Belcher J. Statin use and clinical osteoarthritis in the general population: a longitudinal study. *J Gen Intern Med.* 2013;28(7):943-9.
- Jordan KP, Hayward RA, Blagojevic-Bucknall M, Croft P. Incidence of prostate, breast, lung and colorectal cancer following new consultation for musculoskeletal pain: a cohort study among UK primary care patients. *Int J Cancer.* 2013;133(3):713-20.
- Jones ME, Campbell G, Patel D, Brunner E, Shatapathy CC, Murray-Thomas T, et al. Risk of Mortality (including Sudden Cardiac Death) and Major Cardiovascular Events in Users of Olanzapine and Other Antipsychotics: A Study with the General Practice Research Database. *Cardiovasc Psychiatry Neurol.* 2013;2013:647476.
- Jick SS, Hagberg KW. The risk of adverse outcomes in association with use of testosterone products: a cohort study using the UK-based general practice research database. *British Journal of Clinical Pharmacology.* 2013;75(1):260-70.

- Jameson K, Amber V, D'Oca K, Mills D, Giles A, Ambegaonkar B. Impact of lipid-lowering therapy on the prevalence of dyslipidaemia in patients at high-risk of cardiovascular events in UK primary care - a retrospective database study. *Int J Clin Pract.* 2013;67(12):1228-37.
- Iyer PG, Borah BJ, Heien HC, Das A, Cooper GS, Chak A. Association of Barrett's esophagus with type II Diabetes Mellitus: results from a large population-based case-control study. *Clin Gastroenterol Hepatol.* 2013;11(9):1108-14.e5.
- Imfeld P, Brauchli Pernus YB, Jick SS, Meier CR. Epidemiology, co-morbidities, and medication use of patients with Alzheimer's disease or vascular dementia in the UK. *J Alzheimers Dis.* 2013;35(3):565-73.
- Imfeld P, Bodmer M, Schuerch M, Jick SS, Meier CR. Seizures in patients with Alzheimer's disease or vascular dementia: a population-based nested case-control analysis. *Epilepsia.* 2013;54(4):700-7.
- Imfeld P, Bodmer M, Schuerch M, Jick SS, Meier CR. Risk of incident stroke in patients with Alzheimer disease or vascular dementia. *Neurology.* 2013;81(10):910-9.
- Humes DJ, Radcliffe RS, Camm C, West J. Population-based study of presentation and adverse outcomes after femoral hernia surgery. *Br J Surg.* 2013;100(13):1827-32.
- Howell-Jones R, Soldan K, Wetten S, Mesher D, Williams T, Gill ON, et al. Declining genital Warts in young women in England associated with HPV 16/18 vaccination: an ecological study. *J Infect Dis.* 2013;208(9):1397-403.
- Hong J-L, Meier CR, Sandler RS, Jick SS, Stürmer T. Risk of colorectal cancer after initiation of orlistat: matched cohort study. *BMJ.* 2013;347:f5039.
- Hong J, Reed C, Novick D, Happich M. Costs associated with treatment of chronic low back pain: an analysis of the UK General Practice Research Database. *Spine.* 2013;38(1):75-82.
- Holden SH, Barnett AH, Peters JR, Jenkins-Jones S, Poole CD, Morgan CL, et al. The incidence of type 2 diabetes in the United Kingdom from 1991 to 2010. *Diabetes Obes Metab.* 2013;15(9):844-52.
- Holden SE, Jenkins-Jones S, Poole CD, Morgan CL, Coghill D, Currie CJ. The prevalence and incidence, resource use and financial costs of treating people with attention deficit/hyperactivity disorder (ADHD) in the United Kingdom (1998 to 2010). *Child Adolesc Psychiatry Ment Health.* 2013;7(1):34.
- Hippisley-Cox J, Coupland C. Predicting risk of emergency admission to hospital using primary care data: derivation and validation of QAdmissions score. *BMJ Open.* 2013;3(8):e003482.
- Hicks BM, Murray LJ, Powe DG, Hughes CM, Cardwell CR. β -Blocker usage and colorectal cancer mortality: a nested case-control study in the UK Clinical Practice Research Datalink cohort. *Ann Oncol.* 2013;24(12):3100-6.
- Herrett E, Shah AD, Boggon R, Denaxas S, Smeeth L, van Staa T, et al. Completeness and diagnostic validity of recording acute myocardial infarction events in primary care, hospital care, disease registry, and national mortality records: cohort study. *BMJ.* 2013;346:f2350.
- Herrett E, George J, Denaxas S, Bhaskaran K, Timmis A, Hemingway H, et al. Type and timing of heralding in ST-elevation and non-ST-elevation myocardial infarction: an analysis of prospectively collected electronic healthcare records linked to the national registry of acute coronary syndromes. *Eur Heart J Acute Cardiovasc Care.* 2013;2(3):235-45.

- Hawkins NM, Scholes S, Bajekal M, Love H, O'Flaherty M, Raine R, et al. The UK National Health Service: delivering equitable treatment across the spectrum of coronary disease. *Circ Cardiovasc Qual Outcomes*. 2013;6(2):208-16.
- Hammad TA, Margulis AV, Ding Y, Strazzeri MM, Epperly H. Determining the predictive value of Read codes to identify congenital cardiac malformations in the UK Clinical Practice Research Datalink. *Pharmacoepidemiol Drug Saf*. 2013;22(11):1233-8.
- Hall GC, Morant SV, Carroll D, Gabriel ZL, McQuay HJ. An observational descriptive study of the epidemiology and treatment of neuropathic pain in a UK general population. *BMC Fam Pract*. 2013;14:28.
- Hak E, Shea KM, Jick SS. Association of infant pneumococcal vaccination with pneumococcal pneumonia among mothers: a nested case-control study using the GPRD. *Vaccine*. 2013;31(12):1590-6.
- Hak E, Mulder B, Schuiling-Veninga CCM, de Vries TW, Jick SS. Use of acid-suppressive drugs in pregnancy and the risk of childhood asthma: bidirectional crossover study using the general practice research database. *Drug Safety*. 2013;36(11):1097-104.
- Hak E, de Vries TW, Hoekstra PJ, Jick SS. Association of childhood attention-deficit/hyperactivity disorder with atopic diseases and skin infections? A matched case-control study using the General Practice Research Database. *Ann Allergy Asthma Immunol*. 2013;111(2):102-6.e2.
- Hagberg KW, Taylor A, Hernandez RK, Jick S. Incidence of bone metastases in breast cancer patients in the United Kingdom: results of a multi-database linkage study using the general practice research database. *Cancer Epidemiol*. 2013;37(3):240-6.
- George J, Majeed W, Mackenzie IS, Macdonald TM, Wei L. Association between cardiovascular events and sodium-containing effervescent, dispersible, and soluble drugs: nested case-control study. *BMJ*. 2013;347:f6954.
- García Rodríguez LA, Johansson S, Cea Soriano L. Use of clopidogrel and proton pump inhibitors after a serious acute coronary event: risk of coronary events and peptic ulcer bleeding. *Thromb Haemost*. 2013;110(5):1014-24.
- Frisher M, Martino OI, Bashford J, Crome I, Croft P. Incidence of psychoses among drug dependent patients in primary care with no psychiatric history: a retrospective observational matched-cohort study. *The European Journal of Psychiatry*. 2013;27:240-7.
- Evans SJW, Douglas I, Rawlins MD, Wexler NS, Tabrizi SJ, Smeeth L. Prevalence of adult Huntington's disease in the UK based on diagnoses recorded in general practice records. *J Neurol Neurosurg Psychiatr*. 2013;84(10):1156-60.
- Erviti J, Alonso A, Oliva B, Gorricho J, López A, Timoner J, et al. Oral bisphosphonates are associated with increased risk of subtrochanteric and diaphyseal fractures in elderly women: a nested case-control study. *BMJ Open*. 2013;3(1).
- Erviti J, Alonso A, Gorricho J, López A. Oral bisphosphonates may not decrease hip fracture risk in elderly Spanish women: a nested case-control study. *BMJ Open*. 2013;3(2).
- Douglas JJ, Langham J, Bhaskaran K, Brauer R, Smeeth L. Orlistat and the risk of acute liver injury: self controlled case series study in UK Clinical Practice Research Datalink. *BMJ*. 2013;346:f1936.
- Douglas I, Evans S, Rawlins MD, Smeeth L, Tabrizi SJ, Wexler NS. Juvenile Huntington's disease: a population-

based study using the General Practice Research Database. *BMJ Open*. 2013;3(4).

- Donegan K, Beau-Lejdstrom R, King B, Seabroke S, Thomson A, Bryan P. Bivalent human papillomavirus vaccine and the risk of fatigue syndromes in girls in the UK. *Vaccine*. 2013;31(43):4961-7.
- Dommett RM, Redaniel T, Stevens MCG, Martin RM, Hamilton W. Risk of childhood cancer with symptoms in primary care: a population-based case-control study. *Br J Gen Pract*. 2013;63(606):e22-9.
- Dommett RM, Redaniel MT, Stevens MCG, Hamilton W, Martin RM. Features of cancer in teenagers and young adults in primary care: a population-based nested case-control study. *Br J Cancer*. 2013;108(11):2329-33.
- Dodd CN, Romio SA, Black S, Vellozzi C, Andrews N, Sturkenboom M, et al. International collaboration to assess the risk of Guillain Barré Syndrome following Influenza A (H1N1) 2009 monovalent vaccines. *Vaccine*. 2013;31(40):4448-58.
- Dawson J, Fulton R, McInnes GT, Morton R, Morrison D, Padmanabhan S, et al. Acetaminophen use and change in blood pressure in a hypertensive population. *Journal of Hypertension*. 2013;31(7):1485-90; discussion 90.
- Davies NM, Smith GD, Windmeijer F, Martin RM. COX-2 selective nonsteroidal anti-inflammatory drugs and risk of gastrointestinal tract complications and myocardial infarction: an instrumental variable analysis. *Epidemiology*. 2013;24(3):352-62.
- Davies NM, Gunnell D, Thomas KH, Metcalfe C, Windmeijer F, Martin RM. Physicians' prescribing preferences were a potential instrument for patients' actual prescriptions of antidepressants. *J Clin Epidemiol*. 2013;66(12):1386-96.
- Currie CJ, Poole CD, Evans M, Peters JR, Morgan CL. Mortality and other important diabetes-related outcomes with insulin vs other antihyperglycemic therapies in type 2 diabetes. *J Clin Endocrinol Metab*. 2013;98(2):668-77.
- Culliford D, Maskell J, Judge A, Arden NK, group CS. A population-based survival analysis describing the association of body mass index on time to revision for total hip and knee replacements: results from the UK general practice research database. *BMJ Open*. 2013;3(11):e003614.
- Crowther MJ, Lambert PC, Abrams KR. Adjusting for measurement error in baseline prognostic biomarkers included in a time-to-event analysis: a joint modelling approach. *BMC Med Res Methodol*. 2013;13:146.
- Crooks CJ, West J, Card TR. Comorbidities affect risk of nonvariceal upper gastrointestinal bleeding. *Gastroenterology*. 2013;144(7):1384-93, 93.e1-2; quiz e18-19.
- Crooks CJ, Card TR, West J. Excess long-term mortality following non-variceal upper gastrointestinal bleeding: a population-based cohort study. *PLoS Med*. 2013;10(4):e1001437.
- Cotton SJ, Belcher J, Rose P, K Jagadeesan S, Neal RD. The risk of a subsequent cancer diagnosis after herpes zoster infection: primary care database study. *Br J Cancer*. 2013;108(3):721-6.
- Cornish RP, Boyd A, Van Staa T, Salisbury C, Macleod J. Socio-economic position and childhood multimorbidity: a study using linkage between the Avon Longitudinal Study of Parents and Children and the General Practice Research Database. *Int J Equity Health*. 2013;12:66.
- Charlton RA, Snowball JM, Bloomfield K, de Vries CS. Colorectal cancer risk reduction following macrogol

exposure: a cohort and nested case control study in the UK. *PloS One*. 2013;8(12):e83203.

- Charlton RA, Hutchison A, Davis KJ, de Vries CS. Asthma management in pregnancy. *PloS One*. 2013;8(4):e60247.
- Charlton J, Rudisill C, Bhattarai N, Gulliford M. Impact of deprivation on occurrence, outcomes and health care costs of people with multiple morbidity. *J Health Serv Res Policy*. 2013;18(4):215-23.
- Chapman CR, Bradshaw DH. Only modest long-term opioid dose escalation occurs over time in chronic nonmalignant pain management. *J Pain Palliat Care Pharmacother*. 2013;27(4):370-7.
- Carr DF, O'Meara H, Jorgensen AL, Campbell J, Hobbs M, McCann G, et al. SLCO1B1 genetic variant associated with statin-induced myopathy: a proof-of-concept study using the clinical practice research datalink. *Clin Pharmacol Ther*. 2013;94(6):695-701.
- Carey IM, Atkinson RW, Kent AJ, van Staa T, Cook DG, Anderson HR. Mortality associations with long-term exposure to outdoor air pollution in a national English cohort. *Am J Respir Crit Care Med*. 2013;187(11):1226-33.
- Cardwell CR, Coleman HG, Murray LJ, Entschladen F, Powe DG. Beta-blocker usage and breast cancer survival: a nested case-control study within a UK clinical practice research datalink cohort. *Int J Epidemiol*. 2013;42(6):1852-61.
- Card TR, Siffledeen J, West J, Fleming KM. An excess of prior irritable bowel syndrome diagnoses or treatments in Celiac disease: evidence of diagnostic delay. *Scand J Gastroenterol*. 2013;48(7):801-7.
- Busby J, Schroeder K, Woltersdorf W, Sterne JAC, Ben-Shlomo Y, Hay A, et al. Temporal growth and geographic variation in the use of laboratory tests by NHS general practices: using routine data to identify research priorities. *Br J Gen Pract*. 2013;63(609):e256-66.
- Brilleman SL, Salisbury C. Comparing measures of multimorbidity to predict outcomes in primary care: a cross sectional study. *Fam Pract*. 2013;30(2):172-8.
- Brilleman SL, Purdy S, Salisbury C, Windmeijer F, Gravelle H, Hollinghurst S. Implications of comorbidity for primary care costs in the UK: a retrospective observational study. *Br J Gen Pract*. 2013;63(609):e274-82.
- Briggs PE, Praet CA, Humphreys SC, Zhao C. Impact of UK Medical Eligibility Criteria implementation on prescribing of combined hormonal contraceptives. *J Fam Plann Reprod Health Care*. 2013;39(3):190-6.
- Booth HP, Prevost AT, Gulliford MC. Epidemiology of clinical body mass index recording in an obese population in primary care: a cohort study. *J Public Health (Oxf)*. 2013;35(1):67-74.
- Booth HP, Prevost AT, Gulliford MC. Validity of smoking prevalence estimates from primary care electronic health records compared with national population survey data for England, 2007 to 2011. *Pharmacoepidemiol Drug Saf*. 2013;22(12):1357-61.
- Boggon R, van Staa TP, Chapman M, Gallagher AM, Hammad TA, Richards MA. Cancer recording and mortality in the General Practice Research Database and linked cancer registries. *Pharmacoepidemiol Drug Saf*. 2013;22(2):168-75.
- Boggon R, Hubbard R, Smeeth L, Gulliford M, Cassell J, Eaton S, et al. Variability of antibiotic prescribing in patients with chronic obstructive pulmonary disease exacerbations: a cohort study. *BMC Pulm Med*. 2013;13:32.

- Bhaskaran K, Forbes HJ, Douglas I, Leon DA, Smeeth L. Representativeness and optimal use of body mass index (BMI) in the UK Clinical Practice Research Datalink (CPRD). *BMJ Open*. 2013;3(9):e003389.
- Becker C, Meier CR, Jick SS, Bodmer M. Case-control analysis on metformin and cancer of the esophagus. *Cancer Causes Control*. 2013;24(10):1763-70.
- Becker C, Jick SS, Meier CR, Bodmer M. Metformin and the risk of endometrial cancer: a case-control analysis. *Gynecol Oncol*. 2013;129(3):565-9.
- Bazelier MT, de Vries F, Vestergaard P, Herings RMC, Gallagher AM, Leufkens HGM, et al. Risk of fracture with thiazolidinediones: an individual patient data meta-analysis. *Front Endocrinol (Lausanne)*. 2013;4:11.
- Bannon FJ, O'Rourke MA, Murray LJ, Hughes CM, Gavin AT, Fleming SJ, et al. Non-steroidal anti-inflammatory drug use and brain tumour risk: a case-control study within the Clinical Practice Research Datalink. *Cancer Causes Control*. 2013;24(11):2027-34.
- Azoulay L, Dell'Aniello S, Simon T, Renoux C, Suissa S. The concurrent use of antithrombotic therapies and the risk of bleeding in patients with atrial fibrillation. *Thromb Haemost*. 2013;109(3):431-9.
- Atkinson RW, Carey IM, Kent AJ, van Staa TP, Anderson HR, Cook DG. Long-term exposure to outdoor air pollution and incidence of cardiovascular diseases. *Epidemiology*. 2013;24(1):44-53.
- Assayag J, Yin H, Benayoun S, Pollak MN, Suissa S, Azoulay L. Androgen deprivation therapy and the risk of colorectal cancer in patients with prostate cancer. *Cancer Causes Control*. 2013;24(5):839-45.
- Alexandropoulou K, van Vlymen J, Reid F, Poullis A, Kang J-Y. Temporal trends of Barrett's oesophagus and gastro-oesophageal reflux and related oesophageal cancer over a 10-year period in England and Wales and associated proton pump inhibitor and H2RA prescriptions: a GPRD study. *Eur J Gastroenterol Hepatol*. 2013;25(1):15-21.
- Abdul Sultan A, West J, Tata LJ, Fleming KM, Nelson-Piercy C, Grainge MJ. Risk of first venous thromboembolism in pregnant women in hospital: population based cohort study from England. *BMJ*. 2013;347:f6099.
- Abdul Sultan A, Tata LJ, Grainge MJ, West J. The incidence of first venous thromboembolism in and around pregnancy using linked primary and secondary care data: a population based cohort study from England and comparative meta-analysis. *PloS One*. 2013;8(7):e70310.
- Zenner D, Kruijshaar ME, Andrews N, Abubakar I. Risk of tuberculosis in pregnancy: a national, primary care-based cohort and self-controlled case series study. *Am J Respir Crit Care Med*. 2012;185(7):779-84.
- Wright E, Schofield PT, Seed P, Molokhia M. Bisphosphonates and risk of upper gastrointestinal cancer--a case control study using the General Practice Research Database (GPRD). *PloS One*. 2012;7(10):e47616.
- Wei L, MacDonald T, Shimi S. Association between appendectomy in females and subsequent pregnancy rate: a cohort study. *Fertil Steril*. 2012;98(2):401-5.
- Webb RT, Kontopantelis E, Doran T, Qin P, Creed F, Kapur N. Risk of self-harm in physically ill patients in UK primary care. *J Psychosom Res*. 2012;73(2):92-7.
- Webb RT, Kontopantelis E, Doran T, Qin P, Creed F, Kapur N. Suicide risk in primary care patients with major physical diseases: a case-control study. *Arch Gen Psychiatry*. 2012;69(3):256-64.

- Warren-Gash C, Hayward AC, Hemingway H, Denaxas S, Thomas SL, Timmis AD, et al. Influenza infection and risk of acute myocardial infarction in England and Wales: a CALIBER self-controlled case series study. *J Infect Dis.* 2012;206(11):1652-9.
- Wang Z, Shah AD, Tate AR, Denaxas S, Shawe-Taylor J, Hemingway H. Extracting diagnoses and investigation results from unstructured text in electronic health records by semi-supervised machine learning. *PLoS One.* 2012;7(1):e30412.
- Walker AJ, Grainge MJ, Card TR. Aspirin and other non-steroidal anti-inflammatory drug use and colorectal cancer survival: a cohort study. *Br J Cancer.* 2012;107(9):1602-7.
- Walker AJ, Grainge M, Bates TE, Card TR. Survival of glioma and colorectal cancer patients using tricyclic antidepressants post-diagnosis. *Cancer Causes Control.* 2012;23(12):1959-64.
- Violato M, Gray A, Papanicolas I, Ouellet M. Resource use and costs associated with coeliac disease before and after diagnosis in 3,646 cases: results of a UK primary care database analysis. *PLoS One.* 2012;7(7):e41308.
- van Staa TP, Patel D, Gallagher AM, de Bruin ML. Glucose-lowering agents and the patterns of risk for cancer: a study with the General Practice Research Database and secondary care data. *Diabetologia.* 2012;55(3):654-65.
- Vamos EP, Harris M, Millett C, Pape UJ, Khunti K, Curcin V, et al. Association of systolic and diastolic blood pressure and all cause mortality in people with newly diagnosed type 2 diabetes: retrospective cohort study. *BMJ.* 2012;345:e5567.
- Toovey S, Prinssen EP, Rayner CR, Thakrar BT, Dutkowski R, Koerner A, et al. Post-marketing assessment of neuropsychiatric adverse events in influenza patients treated with oseltamivir: an updated review. *Adv Ther.* 2012;29(10):826-48.
- Taylor A, Stapley S, Hamilton W. Jaundice in primary care: a cohort study of adults aged >45 years using electronic medical records. *Fam Pract.* 2012;29(4):416-20.
- Suissa D, Delaney JAC, Dial S, Brassard P. Non-steroidal anti-inflammatory drugs and the risk of *Clostridium difficile*-associated disease. *British Journal of Clinical Pharmacology.* 2012;74(2):370-5.
- Stapley S, Peters TJ, Neal RD, Rose PW, Walter FM, Hamilton W. The risk of pancreatic cancer in symptomatic patients in primary care: a large case-control study using electronic records. *Br J Cancer.* 2012;106(12):1940-4.
- Staa T-Pv, Goldacre B, Gulliford M, Cassell J, Pirmohamed M, Taweel A, et al. Pragmatic randomised trials using routine electronic health records: putting them to the test. *BMJ.* 2012;344:e55.
- Spoenclin J, Voegel JJ, Jick SS, Meier CR. A study on the epidemiology of rosacea in the U.K. *Br J Dermatol.* 2012;167(3):598-605.
- Singh G, Bell GS, Driever PH, Sander JW. Cancer risk in people with epilepsy using valproate-sodium. *Acta Neurol Scand.* 2012;125(4):234-40.
- Simkiss DE, Spencer NJ, Stallard N, Thorogood M. Health service use in families where children enter public care: a nested case control study using the General Practice Research Database. *BMC Health Serv Res.* 2012;12:65.

- Shi N, Durden E, Torres A, Cao Z, Happich M. Predictors of Treatment with Duloxetine or Venlafaxine XR among Adult Patients Treated for Depression in Primary Care Practices in the United Kingdom. *Depress Res Treat.* 2012;2012:815363.
- Shi N, Cao Z, Durden E, Schacht A, Torres A, Happich M. Healthcare utilization among patients with depression before and after initiating duloxetine in the United Kingdom. *J Med Econ.* 2012;15(4):672-80.
- Shephard EA, Stapley S, Neal RD, Rose P, Walter FM, Hamilton WT. Clinical features of bladder cancer in primary care. *Br J Gen Pract.* 2012;62(602):e598-604.
- Shah AD, Martinez C, Hemingway H. The freetext matching algorithm: a computer program to extract diagnoses and causes of death from unstructured text in electronic health records. *BMC Med Inform Decis Mak.* 2012;12:88.
- Saragoussi D, Chollet J, Bineau S, Chalem Y, Milea D. Antidepressant switching patterns in the treatment of major depressive disorder: a General Practice Research Database (GPRD) Study. *Int J Clin Pract.* 2012;66(11):1079-87.
- Sammon CJ, McGrogan A, Snowball J, de Vries CS. Factors associated with uptake of seasonal and pandemic influenza vaccine among clinical risk groups in the UK: an analysis using the General Practice Research Database. *Vaccine.* 2012;30(14):2483-9.
- Renoux C, Dell'Aniello S, Brophy JM, Suissa S. Dopamine agonist use and the risk of heart failure. *Pharmacoepidemiol Drug Saf.* 2012;21(1):34-41.
- Redaniel MTM, Jeffreys M, May MT, Ben-Shlomo Y, Martin RM. Associations of type 2 diabetes and diabetes treatment with breast cancer risk and mortality: a population-based cohort study among British women. *Cancer Causes Control.* 2012;23(11):1785-95.
- Platt RW, Delaney JAC, Suissa S. The positivity assumption and marginal structural models: the example of warfarin use and risk of bleeding. *Eur J Epidemiol.* 2012;27(2):77-83.
- Perry DC, Bruce CE, Pope D, Dangerfield P, Platt MJ, Hall AJ. Legg-Calvé-Perthes disease in the UK: geographic and temporal trends in incidence reflecting differences in degree of deprivation in childhood. *Arthritis Rheum.* 2012;64(5):1673-9.
- Perry DC, Bruce CE, Pope D, Dangerfield P, Platt MJ, Hall AJ. Comorbidities in Perthes' disease: a case control study using the General Practice Research database. *J Bone Joint Surg Br.* 2012;94(12):1684-9.
- Nicholas JM, Ridsdale L, Richardson MP, Ashworth M, Gulliford MC. Trends in antiepileptic drug utilisation in UK primary care 1993-2008: cohort study using the General Practice Research Database. *Seizure.* 2012;21(6):466-70.
- Müllerova H, Chigbo C, Hagan GW, Woodhead MA, Miravittles M, Davis KJ, et al. The natural history of community-acquired pneumonia in COPD patients: a population database analysis. *Respir Med.* 2012;106(8):1124-33.
- Morgan CL, Poole CD, Evans M, Barnett AH, Jenkins-Jones S, Currie CJ. What next after metformin? A retrospective evaluation of the outcome of second-line, glucose-lowering therapies in people with type 2 diabetes. *J Clin Endocrinol Metab.* 2012;97(12):4605-12.
- Morgan CL, Jenkins-Jones S, Currie CJ, Rees DA. Evaluation of adverse outcome in young women with polycystic ovary syndrome versus matched, reference controls: a retrospective, observational study. *J Clin*

Endocrinol Metab. 2012;97(9):3251-60.

- Mackenzie IS, Macdonald TM, Thompson A, Morant S, Wei L. Spironolactone and risk of incident breast cancer in women older than 55 years: retrospective, matched cohort study. *BMJ*. 2012;345:e4447.
- Lo Re V, Haynes K, Ming EE, Wood Ives J, Horne LN, Fortier K, et al. Safety of saxagliptin: rationale for and design of a series of postmarketing observational studies. *Pharmacoepidemiol Drug Saf*. 2012;21(11):1202-15.
- Li L, Roddam A, Gitlin M, Taylor A, Shepherd S, Shearer A, et al. Persistence with osteoporosis medications among postmenopausal women in the UK General Practice Research Database. *Menopause*. 2012;19(1):33-40.
- Li H, Hodgson E, Watson L, Shukla A, Nelson JJ. Comorbidities and Concomitant Medication Use in Men with Prostate Cancer or High Levels of PSA Compared to Matched Controls: A GPRD Analysis. *J Cancer Epidemiol*. 2012;2012:291704.
- Leonard CE, Freeman CP, Newcomb CW, Reese PP, Herlim M, Bilker WB, et al. Proton pump inhibitors and traditional nonsteroidal anti-inflammatory drugs and the risk of acute interstitial nephritis and acute kidney injury. *Pharmacoepidemiol Drug Saf*. 2012;21(11):1155-72.
- Lee S, Monz BU, Clemens A, Brueckmann M, Lip GYH. Representativeness of the dabigatran, apixaban and rivaroxaban clinical trial populations to real-world atrial fibrillation patients in the United Kingdom: a cross-sectional analysis using the General Practice Research Database. *BMJ Open*. 2012;2(6).
- Le Ray I, Dell'Aniello S, Bonnetain F, Azoulay L, Suissa S. Local estrogen therapy and risk of breast cancer recurrence among hormone-treated patients: a nested case-control study. *Breast Cancer Res Treat*. 2012;135(2):603-9.
- Langan SM, Groves RW, Card TR, Gulliford MC. Incidence, mortality, and disease associations of pyoderma gangrenosum in the United Kingdom: a retrospective cohort study. *J Invest Dermatol*. 2012;132(9):2166-70.
- Lalmohamed A, de Vries F, Bazelier MT, Cooper A, van Staa T-P, Cooper C, et al. Risk of fracture after bariatric surgery in the United Kingdom: population based, retrospective cohort study. *BMJ*. 2012;345:e5085.
- Lalmohamed A, Bazelier MT, Van Staa TP, Uitdehaag BMJ, Leufkens HGM, De Boer A, et al. Causes of death in patients with multiple sclerosis and matched referent subjects: a population-based cohort study. *Eur J Neurol*. 2012;19(7):1007-14.
- Khashan AS, Quigley EMM, McNamee R, McCarthy FP, Shanahan F, Kenny LC. Increased risk of miscarriage and ectopic pregnancy among women with irritable bowel syndrome. *Clin Gastroenterol Hepatol*. 2012;10(8):902-9.
- Jick S, Pennap D. Drospirenone- and levonorgestrel-containing oral contraceptives and the risk of gallbladder disease. *Contraception*. 2012;86(3):220-3.
- Jick H, Wilson A, Wiggins P, Chamberlain DP. Comparison of prescription drug costs in the United States and the United Kingdom, part 2: proton pump inhibitors. *Pharmacotherapy*. 2012;32(6):489-92.
- Irizarry MC, Webb DJ, Boudiaf N, Logie J, Habel LA, Udaltsova N, et al. Risk of cancer in patients exposed to gabapentin in two electronic medical record systems. *Pharmacoepidemiol Drug Saf*. 2012;21(2):214-25.
- Imfeld P, Bodmer M, Jick SS, Meier CR. Metformin, other antidiabetic drugs, and risk of Alzheimer's disease: a

population-based case-control study. *Journal of the American Geriatrics Society*. 2012;60(5):916-21.

- Humes DJ, West J. Role of acute diverticulitis in the development of complicated colonic diverticular disease and 1-year mortality after diagnosis in the UK: population-based cohort study. *Gut*. 2012;61(1):95-100.
- Hodgkinson JA, Taylor CJ, Hobbs FDR. Treatment pathways for patients with atrial fibrillation. *Int J Clin Pract*. 2012;66(1):44-52.
- Hirst CJ, Cabrera C, Kirby M. Epidemiology of castration resistant prostate cancer: a longitudinal analysis using a UK primary care database. *Cancer Epidemiol*. 2012;36(6):e349-53.
- Hirji I, Guo Z, Andersson SW, Hammar N, Gomez-Caminero A. Incidence of urinary tract infection among patients with type 2 diabetes in the UK General Practice Research Database (GPRD). *J Diabetes Complicat*. 2012;26(6):513-6.
- Hirji I, Andersson SW, Guo Z, Hammar N, Gomez-Caminero A. Incidence of genital infection among patients with type 2 diabetes in the UK General Practice Research Database. *J Diabetes Complicat*. 2012;26(6):501-5.
- Higginson IJ, Gao W. Opioid prescribing for cancer pain during the last 3 months of life: associated factors and 9-year trends in a nationwide United Kingdom cohort study. *J Clin Oncol*. 2012;30(35):4373-9.
- Hesdorffer DC, Ishihara L, Mynepalli L, Webb DJ, Weil J, Hauser WA. Epilepsy, suicidality, and psychiatric disorders: a bidirectional association. *Ann Neurol*. 2012;72(2):184-91.
- Hernandez RK, Do TP, Critchlow CW, Dent RE, Jick SS. Patient-related risk factors for fracture-healing complications in the United Kingdom General Practice Research Database. *Acta Orthop*. 2012;83(6):653-60.
- Hawkins NM, Scholes S, Bajekal M, Love H, O'Flaherty M, Raine R, et al. Community care in England: reducing socioeconomic inequalities in heart failure. *Circulation*. 2012;126(9):1050-7.
- Green J, Czanner G, Reeves G, Watson J, Wise L, Roddam A, et al. Menopausal hormone therapy and risk of gastrointestinal cancer: nested case-control study within a prospective cohort, and meta-analysis. *Int J Cancer*. 2012;130(10):2387-96.
- Girman CJ, Kou TD, Brodovicz K, Alexander CM, O'Neill EA, Engel S, et al. Risk of acute renal failure in patients with Type 2 diabetes mellitus. *Diabet Med*. 2012;29(5):614-21.
- Gallagher AM, de Vries F, Plumb JM, Haß B, Clemens A, van Staa T-P. Quality of INR control and outcomes following venous thromboembolism. *Clin Appl Thromb Hemost*. 2012;18(4):370-8.
- Forbes HJ, Thomas SL, Smeeth L, Langan SM. Prescription of antiviral therapy after herpes zoster in general practice: who receives therapy? *Br J Gen Pract*. 2012;62(605):e808-14.
- Fleming KM, Aithal GP, Card TR, West J. All-cause mortality in people with cirrhosis compared with the general population: a population-based cohort study. *Liver Int*. 2012;32(1):79-84.
- Ernst P, Renoux C, Dell'Aniello S, Suissa S. Pramipexole use and the risk of pneumonia. *BMC neurology*. 2012;12:113.
- Edwards CJ, Campbell J, van Staa T, Arden NK. Regional and temporal variation in the treatment of rheumatoid arthritis across the UK: a descriptive register-based cohort study. *BMJ Open*. 2012;2(6).
- Dregan A, van Staa T, McDermott L, McCann G, Ashworth M, Charlton J, et al. Cluster randomized trial in the general practice research database: 2. Secondary prevention after first stroke (eCRT study): study protocol for

a randomized controlled trial. *Trials*. 2012;13:181.

- Dregan A, Moller H, Murray-Thomas T, Gulliford MC. Validity of cancer diagnosis in a primary care database compared with linked cancer registrations in England. Population-based cohort study. *Cancer Epidemiol*. 2012;36(5):425-9.
- Douglas JJ, Evans SJW, Hingorani AD, Grosso AM, Timmis A, Hemingway H, et al. Clopidogrel and interaction with proton pump inhibitors: comparison between cohort and within person study designs. *BMJ*. 2012;345:e4388.
- Dommert RM, Redaniel MT, Stevens MCG, Hamilton W, Martin RM. Features of childhood cancer in primary care: a population-based nested case-control study. *Br J Cancer*. 2012;106(5):982-7.
- Denaxas SC, George J, Herrett E, Shah AD, Kalra D, Hingorani AD, et al. Data resource profile: cardiovascular disease research using linked bespoke studies and electronic health records (CALIBER). *Int J Epidemiol*. 2012;41(6):1625-38.
- de Moor C, Reardon G, McLaughlin J, Maiese EM, Navaratnam P. A retrospective comparison of acute rhinosinusitis outcomes in patients prescribed antibiotics, mometasone furoate nasal spray, or both. *Am J Rhinol Allergy*. 2012;26(4):308-14.
- de Jong HJ, Klungel OH, van Dijk L, Vandebriel RJ, Leufkens HGM, van der Laan JW, et al. Use of statins is associated with an increased risk of rheumatoid arthritis. *Annals of the Rheumatic Diseases*. 2012;71(5):648-54.
- Culliford DJ, Maskell J, Kiran A, Judge A, Javaid MK, Cooper C, et al. The lifetime risk of total hip and knee arthroplasty: results from the UK general practice research database. *Osteoarthritis and Cartilage*. 2012;20(6):519-24.
- Crooks CJ, Card TR, West J. Defining upper gastrointestinal bleeding from linked primary and secondary care data and the effect on occurrence and 28 day mortality. *BMC Health Serv Res*. 2012;12:392.
- Close H, Mason JM, Wilson D, Hungin APS. Hormone replacement therapy is associated with gastro-oesophageal reflux disease: a retrospective cohort study. *BMC Gastroenterol*. 2012;12:56.
- Charlton R, Snowball J, Bloomfield K, de Vries C. Colorectal cancer incidence on the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2012;21(7):775-83.
- Cardwell CR, Abnet CC, Veal P, Hughes CM, Cantwell MM, Murray LJ. Exposure to oral bisphosphonates and risk of cancer. *Int J Cancer*. 2012;131(5):E717-25.
- Cai B, Xu W, Bortnichak E, Watson DJ. An algorithm to identify medical practices common to both the General Practice Research Database and The Health Improvement Network database. *Pharmacoepidemiol Drug Saf*. 2012;21(7):770-4.
- Brodovicz KG, Kou TD, Alexander CM, O'Neill EA, Engel SS, Girman CJ, et al. Impact of diabetes duration and chronic pancreatitis on the association between type 2 diabetes and pancreatic cancer risk. *Diabetes Obes Metab*. 2012;14(12):1123-8.
- Bradley MC, Murray LJ, Cantwell MM, Hughes CM. Proton pump inhibitors and histamine-2-receptor antagonists and pancreatic cancer risk: a nested case-control study. *Br J Cancer*. 2012;106(1):233-9.
- Boggon R, Lip GYH, Gallagher AM, van Staa TP. Resource utilization and outcomes in patients with atrial

fibrillation: a case control study. *Appl Health Econ Health Policy*. 2012;10(4):249-59.

- Boggon. Current prescribing of statins and persistence to statins following ACS in the UK: a MINAP/GPRD study. *British Journal of Cardiology*. 2012;19(1).
- Bodmer M, Becker C, Meier C, Jick SS, Meier CR. Use of metformin is not associated with a decreased risk of colorectal cancer: a case-control analysis. *Cancer Epidemiol Biomarkers Prev*. 2012;21(2):280-6.
- Bodmer M, Becker C, Meier C, Jick SS, Meier CR. Use of antidiabetic agents and the risk of pancreatic cancer: a case-control analysis. *Am J Gastroenterol*. 2012;107(4):620-6.
- Bodmer M, Becker C, Jick SS, Meier CR. Metformin does not alter the risk of lung cancer: a case-control analysis. *Lung Cancer*. 2012;78(2):133-7.
- Blumentals WA, Nevitt A, Peng MM, Toovey S. Body mass index and the incidence of influenza-associated pneumonia in a UK primary care cohort. *Influenza Other Respir Viruses*. 2012;6(1):28-36.
- Bhattarai N, Charlton J, Rudisill C, Gulliford MC. Coding, recording and incidence of different forms of coronary heart disease in primary care. *PloS One*. 2012;7(1):e29776.
- Bhaskaran K, Douglas I, Evans S, van Staa T, Smeeth L. Angiotensin receptor blockers and risk of cancer: cohort study among people receiving antihypertensive drugs in UK General Practice Research Database. *BMJ*. 2012;344:e2697.
- Berger A, Sadosky A, Dukes E, Edelsberg J, Oster G. Clinical characteristics and patterns of healthcare utilization in patients with painful neuropathic disorders in UK general practice: a retrospective cohort study. *BMC neurology*. 2012;12:8.
- Bazelier MT, van Staa T-P, Uitdehaag BMJ, Cooper C, Leufkens HGM, Vestergaard P, et al. A simple score for estimating the long-term risk of fracture in patients with multiple sclerosis. *Neurology*. 2012;79(9):922-8.
- Bazelier MT, Mueller-Schotte S, Leufkens HGM, Uitdehaag BMJ, van Staa T, de Vries F. Risk of cataract and glaucoma in patients with multiple sclerosis. *Mult Scler*. 2012;18(5):628-38.
- Bartholomeeusen S, Van Zundert J, Truyers C, Buntinx F, Paulus D. Higher incidence of common diagnoses in patients with low back pain in primary care. *Pain Pract*. 2012;12(1):1-6.
- Bakhai A, Rigney U, Hollis S, Emmas C. Co-administration of statins with cytochrome P450 3A4 inhibitors in a UK primary care population. *Pharmacoepidemiol Drug Saf*. 2012;21(5):485-93.
- Azoulay L, Yin H, Filion KB, Assayag J, Majdan A, Pollak MN, et al. The use of pioglitazone and the risk of bladder cancer in people with type 2 diabetes: nested case-control study. *BMJ*. 2012;344:e3645.
- Azoulay L, Dell'Aniello S, Simon TA, Langleben D, Renoux C, Suissa S. A net clinical benefit analysis of warfarin and aspirin on stroke in patients with atrial fibrillation: a nested case-control study. *BMC Cardiovasc Disord*. 2012;12:49.
- Azoulay L, Assimes TL, Yin H, Bartels DB, Schiffrin EL, Suissa S. Long-term use of angiotensin receptor blockers and the risk of cancer. *PloS One*. 2012;7(12):e50893.
- Amar RK, Jick SS, Rosenberg D, Maher TM, Meier CR. Drug-/radiation-induced interstitial lung disease in the United Kingdom general population: incidence, all-cause mortality and characteristics at diagnosis. *Respirology*. 2012;17(5):861-8.

- Amar RK, Jick SS, Rosenberg D, Maher TM, Meier CR. Incidence of the pneumoconioses in the United Kingdom general population between 1997 and 2008. *Respiration*. 2012;84(3):200-6.
- Watson L, Wilson BP, Alsop J, Kumar S. Weight and glycaemic control in type 2 diabetes: what is the outcome of insulin initiation? *Diabetes Obes Metab*. 2011;13(9):823-31.
- Walters P, Schofield P, Howard L, Ashworth M, Tylee A. The relationship between asthma and depression in primary care patients: a historical cohort and nested case control study. *PLoS One*. 2011;6(6):e20750.
- Walker AM. Identification of esophageal cancer in the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2011;20(11):1159-67.
- Walker AJ, West J, Grainge MJ, Card TR. Angiotensin converting enzyme inhibitors and hepatocellular carcinoma incidence in the General Practice Research Database. *Cancer Causes Control*. 2011;22(12):1743-7.
- Walker AJ, Card T, Bates TE, Muir K. Tricyclic antidepressants and the incidence of certain cancers: a study using the GPRD. *Br J Cancer*. 2011;104(1):193-7.
- Varyani FK, West J, Card TR. An increased risk of urinary tract infection precedes development of primary biliary cirrhosis. *BMC Gastroenterol*. 2011;11:95.
- Van Staa TP, Setakis E, Di Tanna GL, Lane DA, Lip GYH. A comparison of risk stratification schemes for stroke in 79,884 atrial fibrillation patients in general practice. *J Thromb Haemost*. 2011;9(1):39-48.
- van Soest EM, Valkhoff VE, Mazzaglia G, Schade R, Molokhia M, Goldstein JL, et al. Suboptimal gastroprotective coverage of NSAID use and the risk of upper gastrointestinal bleeding and ulcers: an observational study using three European databases. *Gut*. 2011;60(12):1650-9.
- Vamos EP, Pape UJ, Bottle A, Hamilton FL, Curcin V, Ng A, et al. Association of practice size and pay-for-performance incentives with the quality of diabetes management in primary care. *CMAJ*. 2011;183(12):E809-16.
- Toschke AM, Gulliford MC, Wolfe CDA, Rudd AG, Heuschmann PU. Antihypertensive treatment after first stroke in primary care: results from the General Practitioner Research Database. *Journal of Hypertension*. 2011;29(1):154-60.
- Toovey S, Jick SS, Meier CR. Parkinson's disease or Parkinson symptoms following seasonal influenza. *Influenza Other Respir Viruses*. 2011;5(5):328-33.
- Suissa S, Azoulay L, Dell'Aniello S, Evans M, Vora J, Pollak M. Long-term effects of insulin glargine on the risk of breast cancer. *Diabetologia*. 2011;54(9):2254-62.
- Suh D-C, Lee D-H, McGuire M, Kim C-M. Impact of rosiglitazone therapy on the lipid profile, glycemic control, and medication costs among type 2 diabetes patients. *Current Medical Research and Opinion*. 2011;27(8):1623-33.
- Stowe J, Andrews N, Bryan P, Seabroke S, Miller E. Risk of convulsions in children after monovalent H1N1 (2009) and trivalent influenza vaccines: a database study. *Vaccine*. 2011;29(51):9467-72.
- Stapley S, Hamilton W. Gynaecological symptoms reported by young women: examining the potential for earlier diagnosis of cervical cancer. *Fam Pract*. 2011;28(6):592-8.
- Shah AD, Nicholas O, Timmis AD, Feder G, Abrams KR, Chen R, et al. Threshold haemoglobin levels and the

prognosis of stable coronary disease: two new cohorts and a systematic review and meta-analysis. *PLoS Med.* 2011;8(5):e1000439.

- Shafe ACE, Lee S, Dalrymple JSO, Whorwell PJ. The LUCK study: Laxative Usage in patients with GP-diagnosed Constipation in the UK, within the general population and in pregnancy. An epidemiological study using the General Practice Research Database (GPRD). *Therap Adv Gastroenterol.* 2011;4(6):343-63.
- Schneider-Lindner V, Quach C, Hanley JA, Suissa S. Secular trends of antibacterial prescribing in UK paediatric primary care. *J Antimicrob Chemother.* 2011;66(2):424-33.
- Schneider-Lindner V, Quach C, Hanley JA, Suissa S. Antibacterial drugs and the risk of community-associated methicillin-resistant *Staphylococcus aureus* in children. *Arch Pediatr Adolesc Med.* 2011;165(12):1107-14.
- Salisbury C, Johnson L, Purdy S, Valderas JM, Montgomery AA. Epidemiology and impact of multimorbidity in primary care: a retrospective cohort study. *Br J Gen Pract.* 2011;61(582):e12-21.
- Rothenbacher D, Primatesta P, Ferreira A, Cea-Soriano L, Rodríguez LAG. Frequency and risk factors of gout flares in a large population-based cohort of incident gout. *Rheumatology (Oxford).* 2011;50(5):973-81.
- Ridsdale L, Charlton J, Ashworth M, Richardson MP, Gulliford MC. Epilepsy mortality and risk factors for death in epilepsy: a population-based study. *Br J Gen Pract.* 2011;61(586):e271-8.
- Rani FA, Byrne P, Cranswick N, Murray ML, Wong ICK. Mortality in children and adolescents prescribed antipsychotic medication: a retrospective cohort study using the UK general practice research database. *Drug Safety.* 2011;34(9):773-81.
- Punekar YS, Ahmad A, Saleh HA. Estimating the effect of nasal steroid treatment on repeat polypectomies: survival time analysis using the General Practice Research Database. *Rhinology.* 2011;49(2):190-4.
- Prieto-Alhambra D, Javaid MK, Maskell J, Judge A, Nevitt M, Cooper C, et al. Changes in hip fracture rate before and after total knee replacement due to osteoarthritis: a population-based cohort study. *Annals of the Rheumatic Diseases.* 2011;70(1):134-8.
- Prieto-Alhambra D, Javaid MK, Judge A, Murray D, Carr A, Cooper C, et al. Association between bisphosphonate use and implant survival after primary total arthroplasty of the knee or hip: population based retrospective cohort study. *BMJ.* 2011;343:d7222.
- Prieto-Alhambra D, Javaid MK, Judge A, Maskell J, Kiran A, de Vries F, et al. Fracture risk before and after total hip replacement in patients with osteoarthritis: potential benefits of bisphosphonate use. *Arthritis Rheum.* 2011;63(4):992-1001.
- Price D, Haughney J, Sims E, Ali M, von Ziegenweidt J, Hillyer EV, et al. Effectiveness of inhaler types for real-world asthma management: retrospective observational study using the GPRD. *J Asthma Allergy.* 2011;4:37-47.
- Patch C, Charlton J, Roderick PJ, Gulliford MC. Use of antihypertensive medications and mortality of patients with autosomal dominant polycystic kidney disease: a population-based study. *Am J Kidney Dis.* 2011;57(6):856-62.
- Parkin L, Sharples K, Hernandez RK, Jick SS. Risk of venous thromboembolism in users of oral contraceptives containing drospirenone or levonorgestrel: nested case-control study based on UK General Practice Research Database. *BMJ.* 2011;342:d2139.

- Parkin L, Hagberg KW, Jick H. Comprehensive comparison of drug prescribing in the United States and United Kingdom. *Pharmacotherapy*. 2011;31(7):623-9.
- Osterman MT, Yang Y-X, Brensinger C, Forde KA, Lichtenstein GR, Lewis JD. No increased risk of myocardial infarction among patients with ulcerative colitis or Crohn's disease. *Clin Gastroenterol Hepatol*. 2011;9(10):875-80.
- Miller DP, Watkins SE, Sampson T, Davis KJ. Long-term use of fluticasone propionate/salmeterol fixed-dose combination and incidence of cataracts and glaucoma among chronic obstructive pulmonary disease patients in the UK General Practice Research Database. *Int J Chron Obstruct Pulmon Dis*. 2011;6:467-76.
- Mehta NN, Yu Y, Pinnelas R, Krishnamoorthy P, Shin DB, Troxel AB, et al. Attributable risk estimate of severe psoriasis on major cardiovascular events. *Am J Med*. 2011;124(8):775.e1-6.
- Martín-Merino E, Johansson S, Morris T, García Rodríguez LA. Androgen deprivation therapy and the risk of coronary heart disease and heart failure in patients with prostate cancer: a nested case-control study in UK primary care. *Drug Safety*. 2011;34(11):1061-77.
- Luqmani R, Suppiah R, Edwards CJ, Phillip R, Maskell J, Culliford D, et al. Mortality in Wegener's granulomatosis: a bimodal pattern. *Rheumatology (Oxford)*. 2011;50(4):697-702.
- Laredo L, Vargas E, Blasco AJ, Aguilar MD, Moreno A, Portolés A. Risk of cerebrovascular accident associated with use of antipsychotics: population-based case-control study. *Journal of the American Geriatrics Society*. 2011;59(7):1182-7.
- Langan SM, Groves RW, West J. The relationship between neurological disease and bullous pemphigoid: a population-based case-control study. *J Invest Dermatol*. 2011;131(3):631-6.
- Lafuma A, Salmon JF, Robert J, Berdeaux G. Treatment persistence and cost-effectiveness of latanoprost/latanoprost-timolol, bimatoprost/bimatoprost-timolol, and travoprost/travoprost-timolol in glaucoma: an analysis based on the United Kingdom general practitioner research database. *Clin Ophthalmol*. 2011;5:361-7.
- Khan NF, Watson E, Rose PW. Primary care consultation behaviours of long-term, adult survivors of cancer in the UK. *Br J Gen Pract*. 2011;61(584):197-9.
- Khan NF, Mant D, Carpenter L, Forman D, Rose PW. Long-term health outcomes in a British cohort of breast, colorectal and prostate cancer survivors: a database study. *Br J Cancer*. 2011;105 Suppl 1:S29-37.
- Kelly F, Armstrong B, Atkinson R, Anderson HR, Barratt B, Beevers S, et al. The London low emission zone baseline study. *Res Rep Health Eff Inst*. 2011(163):3-79.
- Julious SA, Campbell MJ, Bianchi SM, Murray-Thomas T. Seasonality of medical contacts in school-aged children with asthma: association with school holidays. *Public Health*. 2011;125(11):769-76.
- Jordan RE, Lancashire RJ, Adab P. An evaluation of Birmingham Own Health telephone care management service among patients with poorly controlled diabetes. A retrospective comparison with the General Practice Research Database. *BMC Public Health*. 2011;11:707.
- Imkampe AK, Gulliford MC. Trends in Type 1 diabetes incidence in the UK in 0- to 14-year-olds and in 15- to 34-year-olds, 1991-2008. *Diabet Med*. 2011;28(7):811-4.
- Humes DJ, Fleming KM, Spiller RC, West J. Concurrent drug use and the risk of perforated colonic diverticular

disease: a population-based case-control study. *Gut*. 2011;60(2):219-24.

- Huiart L, Dell'Aniello S, Suissa S. Use of tamoxifen and aromatase inhibitors in a large population-based cohort of women with breast cancer. *Br J Cancer*. 2011;104(10):1558-63.
- Huerta C, Rivero E, Montoro MA, García-Rodríguez LA. Risk factors for intestinal ischaemia among patients registered in a UK primary care database: a nested case-control study. *Aliment Pharmacol Ther*. 2011;33(8):969-78.
- Hodgkinson JA, Taylor CJ, Hobbs FDR. Predictors of incident atrial fibrillation and influence of medications: a retrospective case-control study. *Br J Gen Pract*. 2011;61(587):e353-61.
- Herrett EL, Thomas SL, Smeeth L. Validity of diagnoses in the general practice research database. *Br J Gen Pract*. 2011;61(588):438-9.
- Hammersley V, Flint R, Pinnock H, Sheikh A. Developing and testing search strategies to identify patients with active seasonal allergic rhinitis in general practice. *Prim Care Respir J*. 2011;20(1):71-4.
- Gulliford MC, van Staa T, McDermott L, Dregan A, McCann G, Ashworth M, et al. Cluster randomised trial in the General Practice Research Database: 1. Electronic decision support to reduce antibiotic prescribing in primary care (eCRT study). *Trials*. 2011;12:115.
- Gao W, Gulliford M, Higginson IJ. Prescription patterns of analgesics in the last 3 months of life: a retrospective analysis of 10202 lung cancer patients. *Br J Cancer*. 2011;104(11):1704-10.
- Gallagher AM, Smeeth L, Seabroke S, Leufkens HGM, van Staa TP. Risk of death and cardiovascular outcomes with thiazolidinediones: a study with the general practice research database and secondary care data. *PLoS One*. 2011;6(12):e28157.
- Gallagher AM, Setakis E, Plumb JM, Clemens A, van Staa TP. Risks of stroke and mortality associated with suboptimal anticoagulation in atrial fibrillation patients. *Thromb Haemost*. 2011;106(5):968-77.
- French CE, Hughes G, Nicholson A, Yung M, Ross JD, Williams T, et al. Estimation of the rate of pelvic inflammatory disease diagnoses: trends in England, 2000-2008. *Sex Transm Dis*. 2011;38(3):158-62.
- Filion KB, Joseph L, Boivin J-F, Suissa S, Brophy JM. Thiazolidinediones and the risk of incident congestive heart failure among patients with type 2 diabetes mellitus. *Pharmacoepidemiol Drug Saf*. 2011;20(8):785-96.
- Ernst P, Dell'Aniello S, Mikaeloff Y, Suissa S. Risk of herpes zoster in patients prescribed inhaled corticosteroids: a cohort study. *BMC Pulm Med*. 2011;11:59.
- Dregan A, Toschke MA, Wolfe CD, Rudd A, Ashworth M, Gulliford MC, et al. Utility of electronic patient records in primary care for stroke secondary prevention trials. *BMC Public Health*. 2011;11:86.
- Dregan A, Grieve A, van Staa T, Gulliford MC, e CRTRT. Potential application of item-response theory to interpretation of medical codes in electronic patient records. *BMC Med Res Methodol*. 2011;11:168.
- Douglas I, Smeeth L, Irvine D. The use of antidepressants and the risk of haemorrhagic stroke: a nested case control study. *British Journal of Clinical Pharmacology*. 2011;71(1):116-20.
- Dixon WG, Solomon DH. Bisphosphonates and esophageal cancer--a pathway through the confusion. *Nat Rev Rheumatol*. 2011;7(6):369-72.
- DiSantostefano RL, Davis KJ. Prescription patterns in asthma patients initiating salmeterol in UK general

practice: a retrospective cohort study using the General Practice Research Database (GPRD). *Drug Safety*. 2011;34(6):511-20.

- Desai S, Wetten S, Woodhall SC, Peters L, Hughes G, Soldan K. Genital warts and cost of care in England. *Sex Transm Infect*. 2011;87(6):464-8.
- De Natale R, Le Pen C, Berdeaux G. Efficiency of glaucoma drug regulation in 5 European countries: a 1995-2006 longitudinal prescription analysis. *J Glaucoma*. 2011;20(4):234-9.
- Davies NM, Kehoe PG, Ben-Shlomo Y, Martin RM. Associations of anti-hypertensive treatments with Alzheimer's disease, vascular dementia, and other dementias. *J Alzheimers Dis*. 2011;26(4):699-708.
- Chi C-C, Mayon-White RT, Wojnarowska FT. Safety of topical corticosteroids in pregnancy: a population-based cohort study. *J Invest Dermatol*. 2011;131(4):884-91.
- Chi CC, Kirtschig G, Aberer W, Gabbud JP, Lipozenčić J, Kárpáti S, et al. Evidence-based (S3) guideline on topical corticosteroids in pregnancy. *Br J Dermatol*. 2011;165(5):943-52.
- Chen Y-C, Wu J-C, Haschler I, Majeed A, Chen T-J, Wetter T. Academic impact of a public electronic health database: bibliometric analysis of studies using the general practice research database. *PLoS One*. 2011;6(6):e21404.
- Charlton RA, Weil JG, Cunnington MC, Ray S, de Vries CS. Comparing the General Practice Research Database and the UK Epilepsy and Pregnancy Register as tools for postmarketing teratogen surveillance: anticonvulsants and the risk of major congenital malformations. *Drug Safety*. 2011;34(2):157-71.
- Byford S, Barrett B, Despiéglé N, Wade A. Impact of treatment success on health service use and cost in depression: longitudinal database analysis. *Pharmacoeconomics*. 2011;29(2):157-70.
- Brauchli YB, Jick SS, Meier CR. Statin use and risk of first-time psoriasis diagnosis. *J Am Acad Dermatol*. 2011;65(1):77-83.
- Bolland BJR, Culliford DJ, Maskell J, Latham JM, Dunlop DG, Arden NK. The effect of hip and knee arthroplasty on oral anti-inflammatory use and the relationship to body mass index: results from the UK general practice research database. *Osteoarthritis and Cartilage*. 2011;19(1):29-36.
- Boggon R, van Staa TP, Timmis A, Hemingway H, Ray KK, Begg A, et al. Clopidogrel discontinuation after acute coronary syndromes: frequency, predictors and associations with death and myocardial infarction--a hospital registry-primary care linked cohort (MINAP-GPRD). *Eur Heart J*. 2011;32(19):2376-86.
- Bodmer M, Brauchli YB, Jick SS, Meier CR. Diabetes mellitus and the risk of cholecystectomy. *Dig Liver Dis*. 2011;43(9):742-7.
- Bodmer M, Becker C, Meier C, Jick SS, Meier CR. Use of metformin and the risk of ovarian cancer: a case-control analysis. *Gynecol Oncol*. 2011;123(2):200-4.
- Bennett D, Hodgson ME, Shukla A, Logie JW. Prevalence of diagnosed adult immune thrombocytopenia in the United Kingdom. *Adv Ther*. 2011;28(12):1096-104.
- Becker C, Jick SS, Meier CR. NSAID use and risk of Parkinson disease: a population-based case-control study. *Eur J Neurol*. 2011;18(11):1336-42.
- Becker C, Brobert GP, Johansson S, Jick SS, Meier CR. Risk of incident depression in patients with Parkinson

disease in the UK. *Eur J Neurol*. 2011;18(3):448-53.

- Bazelier MT, van Staa T, Uitdehaag BMJ, Cooper C, Leufkens HGM, Vestergaard P, et al. The risk of fracture in patients with multiple sclerosis: the UK general practice research database. *J Bone Miner Res*. 2011;26(9):2271-9.
- Azoulay L, Yin H, Renoux C, Suissa S. The use of atypical antipsychotics and the risk of breast cancer. *Breast Cancer Res Treat*. 2011;129(2):541-8.
- Azoulay L, Yin H, Benayoun S, Renoux C, Boivin J-F, Suissa S. Androgen-deprivation therapy and the risk of stroke in patients with prostate cancer. *European Urology*. 2011;60(6):1244-50.
- Azoulay L, Dell'Aniello S, Huiart L, du Fort GG, Suissa S. Concurrent use of tamoxifen with CYP2D6 inhibitors and the risk of breast cancer recurrence. *Breast Cancer Res Treat*. 2011;126(3):695-703.
- Azoulay L, Dell'Aniello S, Gagnon B, Pollak M, Suissa S. Metformin and the incidence of prostate cancer in patients with type 2 diabetes. *Cancer Epidemiol Biomarkers Prev*. 2011;20(2):337-44.
- Ackers R, Besag FMC, Hughes E, Squier W, Murray ML, Wong ICK. Mortality rates and causes of death in children with epilepsy prescribed antiepileptic drugs: a retrospective cohort study using the UK General Practice Research Database. *Drug Safety*. 2011;34(5):403-13.
- Yong M, Schoonen WM, Li L, Kanas G, Coalson J, Mowat F, et al. Epidemiology of paediatric immune thrombocytopenia in the General Practice Research Database. *Br J Haematol*. 2010;149(6):855-64.
- Wade AG, Saragoussi D, Despiéglé N, François C, Guelfucci F, Toumi M. Healthcare expenditure in severely depressed patients treated with escitalopram, generic SSRIs or venlafaxine in the UK. *Current Medical Research and Opinion*. 2010;26(5):1161-70.
- Thomas M, Murray-Thomas T, Fan T, Williams T, Taylor S. Prescribing patterns of asthma controller therapy for children in UK primary care: a cross-sectional observational study. *BMC Pulm Med*. 2010;10:29.
- Taylor CJ, Hodgkinson J, Hobbs FDR. Rhythm control agents and adverse events in patients with atrial fibrillation. *Int J Clin Pract*. 2010;64(8):1069-75.
- Tate AR, Nicholson A, Cassell JA. Are GPs under-investigating older patients presenting with symptoms of ovarian cancer? Observational study using General Practice Research Database. *Br J Cancer*. 2010;102(6):947-51.
- Siriwardena AN, Gwini SM, Coupland CAC. Influenza vaccination, pneumococcal vaccination and risk of acute myocardial infarction: matched case-control study. *CMAJ*. 2010;182(15):1617-23.
- Schneider C, Jick SS, Bothner U, Meier CR. COPD and the risk of depression. *Chest*. 2010;137(2):341-7.
- Schneider C, Jick SS, Bothner U, Meier CR. Reflux disease, gastrointestinal ulcer or weight loss in patients with COPD. *COPD*. 2010;7(3):172-8.
- Schneider C, Bothner U, Jick SS, Meier CR. Chronic obstructive pulmonary disease and the risk of cardiovascular diseases. *Eur J Epidemiol*. 2010;25(4):253-60.
- Schaer BA, Schneider C, Jick SS, Conen D, Osswald S, Meier CR. Risk for incident atrial fibrillation in patients who receive antihypertensive drugs: a nested case-control study. *Ann Intern Med*. 2010;152(2):78-84.
- Sazonov V, Beetsch J, Phatak H, Wentworth C, Evans M. Association between dyslipidemia and vascular

events in patients treated with statins: report from the UK General Practice Research Database. *Atherosclerosis*. 2010;208(1):210-6.

- Sarpatwari A, Bennett D, Logie JW, Shukla A, Beach KJ, Newland AC, et al. Thromboembolic events among adult patients with primary immune thrombocytopenia in the United Kingdom General Practice Research Database. *Haematologica*. 2010;95(7):1167-75.
- Rodríguez LAG, Wallander M-A, Martín-Merino E, Johansson S. Heart failure, myocardial infarction, lung cancer and death in COPD patients: a UK primary care study. *Respir Med*. 2010;104(11):1691-9.
- Renoux C, Dell'Aniello S, Suissa S. Hormone replacement therapy and the risk of venous thromboembolism: a population-based study. *J Thromb Haemost*. 2010;8(5):979-86.
- Renoux C, Dell'aniello S, Garbe E, Suissa S. Transdermal and oral hormone replacement therapy and the risk of stroke: a nested case-control study. *BMJ*. 2010;340:c2519.
- Nicholson A, Rait G, Murray-Thomas T, Hughes G, Mercer CH, Cassell J. Management of epididymo-orchitis in primary care: results from a large UK primary care database. *Br J Gen Pract*. 2010;60(579):e407-22.
- Nicholson A, Rait G, Murray-Thomas T, Hughes G, Mercer CH, Cassell J. Management of first-episode pelvic inflammatory disease in primary care: results from a large UK primary care database. *Br J Gen Pract*. 2010;60(579):e395-406.
- Ng E, McGrail KM, Johnson JA. Hospitalization risk in a type 2 diabetes cohort. *Health Rep*. 2010;21(3):29-35.
- Miller DP, Watkins SE, Sampson T, Davis KJ. Long-term use of fluticasone propionate/salmeterol fixed-dose combination and incidence of nonvertebral fractures among patients with COPD in the UK General Practice Research Database. *Phys Sportsmed*. 2010;38(4):19-27.
- Merriman NA, Putt ME, Metz DC, Yang Y-X. Hip fracture risk in patients with a diagnosis of pernicious anemia. *Gastroenterology*. 2010;138(4):1330-7.
- Meier C, Brauchli YB, Jick SS, Kraenzlin ME, Meier CR. Use of depot medroxyprogesterone acetate and fracture risk. *J Clin Endocrinol Metab*. 2010;95(11):4909-16.
- Mehta NN, Azfar RS, Shin DB, Neimann AL, Troxel AB, Gelfand JM. Patients with severe psoriasis are at increased risk of cardiovascular mortality: cohort study using the General Practice Research Database. *Eur Heart J*. 2010;31(8):1000-6.
- McDowell SE, Coleman JJ, Evans SJW, Gill PS, Ferner RE. Laboratory monitoring and adverse patient outcomes with antihypertensive therapy in primary care. *Pharmacoepidemiol Drug Saf*. 2010;19(5):482-9.
- Martinez C, Assimes TL, Mines D, Dell'aniello S, Suissa S. Use of venlafaxine compared with other antidepressants and the risk of sudden cardiac death or near death: a nested case-control study. *BMJ*. 2010;340:c249.
- MacDonald MR, Eurich DT, Majumdar SR, Lewsey JD, Bhagra S, Jhund PS, et al. Treatment of type 2 diabetes and outcomes in patients with heart failure: a nested case-control study from the U.K. General Practice Research Database. *Diabetes Care*. 2010;33(6):1213-8.
- Kurd SK, Troxel AB, Crits-Christoph P, Gelfand JM. The risk of depression, anxiety, and suicidality in patients with psoriasis: a population-based cohort study. *Arch Dermatol*. 2010;146(8):891-5.

- Kiri VA, Soriano J, Visick G, Fabbri L. Recent trends in lung cancer and its association with COPD: an analysis using the UK GP Research Database. *Prim Care Respir J.* 2010;19(1):57-61.
- Khan NF, Ward AM, Watson E, Rose PW. Consulting and prescribing behaviour for anxiety and depression in long-term survivors of cancer in the UK. *Eur J Cancer.* 2010;46(18):3339-44.
- Khan NF, Perera R, Harper S, Rose PW. Adaptation and validation of the Charlson Index for Read/OXMIS coded databases. *BMC Fam Pract.* 2010;11:1.
- Khan NF, Mant D, Rose PW. Quality of care for chronic diseases in a British cohort of long-term cancer survivors. *Annals of family medicine.* 2010;8(5):418-24.
- Khan NF, Harrison SE, Rose PW. Validity of diagnostic coding within the General Practice Research Database: a systematic review. *Br J Gen Pract.* 2010;60(572):e128-36.
- Jordan KP, Croft P. Mortality and cancer in patients with new musculoskeletal episodes: a cohort study. *Br J Gen Pract.* 2010;60(572):e105-11.
- Johansson S, Wallander M-A, de Abajo FJ, García Rodríguez LA. Prospective drug safety monitoring using the UK primary-care General Practice Research Database: theoretical framework, feasibility analysis and extrapolation to future scenarios. *Drug Safety.* 2010;33(3):223-32.
- Jick H, Hagberg KW. Effectiveness of influenza vaccination in the United kingdom, 1996-2007. *Pharmacotherapy.* 2010;30(12):1199-206.
- Jick H, Hagberg KW. Measles in the United Kingdom 1990-2008 and the effectiveness of measles vaccines. *Vaccine.* 2010;28(29):4588-92.
- Iversen L, Delaney EK, Hannaford PC, Black C. Menopause-related workload in general practice 1996-2005: a retrospective study in the UK. *Fam Pract.* 2010;27(5):499-506.
- Ishihara L, Webb DJ, Irizarry M, Weil J. Exploring differential prescribing between anti-epileptic drugs in epilepsy patients with a history of mood disorders. *Pharmacoepidemiol Drug Saf.* 2010;19(3):289-95.
- Herrett E, Thomas SL, Schoonen WM, Smeeth L, Hall AJ. Validation and validity of diagnoses in the General Practice Research Database: a systematic review. *British Journal of Clinical Pharmacology.* 2010;69(1):4-14.
- Gulliford MC, Charlton J, Rudd A, Wolfe CD, Toschke AM. Declining 1-year case-fatality of stroke and increasing coverage of vascular risk management: population-based cohort study. *J Neurol Neurosurg Psychiatr.* 2010;81(4):416-22.
- Green J, Czanner G, Reeves G, Watson J, Wise L, Beral V. Oral bisphosphonates and risk of cancer of oesophagus, stomach, and colorectum: case-control analysis within a UK primary care cohort. *BMJ.* 2010;341:c4444.
- Girman CJ, Kou TD, Cai B, Alexander CM, O'Neill EA, Williams-Herman DE, et al. Patients with type 2 diabetes mellitus have higher risk for acute pancreatitis compared with those without diabetes. *Diabetes Obes Metab.* 2010;12(9):766-71.
- Frisher M, Short D, Bashford J. Determining patient characteristics for decision analysis support systems using anonymized electronic patient records. *Health Informatics J.* 2010;16(1):49-57.
- Fleming KM, Aithal GP, Card TR, West J. The rate of decompensation and clinical progression of disease in

people with cirrhosis: a cohort study. *Aliment Pharmacol Ther.* 2010;32(11-12):1343-50.

- Egbring M, Kullak-Ublick GA, Russmann S. Phynx: an open source software solution supporting data management and web-based patient-level data review for drug safety studies in the general practice research database and other health care databases. *Pharmacoepidemiol Drug Saf.* 2010;19(1):38-44.
- Devine S, West S, Andrews E, Tennis P, Hammad TA, Eaton S, et al. The identification of pregnancies within the general practice research database. *Pharmacoepidemiol Drug Saf.* 2010;19(1):45-50.
- de Vries F, Setakis E, Zhang B, van Staa TP. Long-acting β_2 -agonists in adult asthma and the pattern of risk of death and severe asthma outcomes: a study using the GPRD. *Eur Respir J.* 2010;36(3):494-502.
- de Vries F, Setakis E, van Staa T-P. Concomitant use of ibuprofen and paracetamol and the risk of major clinical safety outcomes. *British Journal of Clinical Pharmacology.* 2010;70(3):429-38.
- Currie CJ, Peters JR, Tynan A, Evans M, Heine RJ, Bracco OL, et al. Survival as a function of HbA(1c) in people with type 2 diabetes: a retrospective cohort study. *Lancet.* 2010;375(9713):481-9.
- Culliford DJ, Maskell J, Beard DJ, Murray DW, Price AJ, Arden NK. Temporal trends in hip and knee replacement in the United Kingdom: 1991 to 2006. *J Bone Joint Surg Br.* 2010;92(1):130-5.
- Cornish R, Macleod J, Strang J, Vickerman P, Hickman M. Risk of death during and after opiate substitution treatment in primary care: prospective observational study in UK General Practice Research Database. *BMJ.* 2010;341:c5475.
- Cooper C, Steinbuch M, Stevenson R, Miday R, Watts NB. The epidemiology of osteonecrosis: findings from the GPRD and THIN databases in the UK. *Osteoporos Int.* 2010;21(4):569-77.
- Coleman JJ, McDowell SE, Evans SJW, Gill PS, Ferner RE. Oversight: a retrospective study of biochemical monitoring in patients beginning antihypertensive drug treatment in primary care. *British Journal of Clinical Pharmacology.* 2010;70(1):109-17.
- Charlton RA, Weil JG, Cunningham MC, de Vries CS. Identifying major congenital malformations in the UK General Practice Research Database (GPRD): a study reporting on the sensitivity and added value of photocopied medical records and free text in the GPRD. *Drug Safety.* 2010;33(9):741-50.
- Cardwell CR, Abnet CC, Cantwell MM, Murray LJ. Exposure to oral bisphosphonates and risk of esophageal cancer. *JAMA.* 2010;304(6):657-63.
- Breart G, Cooper C, Meyer O, Speirs C, Deltour N, Reginster JY. Osteoporosis and venous thromboembolism: a retrospective cohort study in the UK General Practice Research Database. *Osteoporos Int.* 2010;21(7):1181-7.
- Bradley MC, Hughes CM, Cantwell MM, Napolitano G, Murray LJ. Non-steroidal anti-inflammatory drugs and pancreatic cancer risk: a nested case-control study. *Br J Cancer.* 2010;102(9):1415-21.
- Bradley MC, Hughes CM, Cantwell MM, Murray LJ. Statins and pancreatic cancer risk: a nested case-control study. *Cancer Causes Control.* 2010;21(12):2093-100.
- Bodmer M, Meier C, Krähenbühl S, Jick SS, Meier CR. Long-term metformin use is associated with decreased risk of breast cancer. *Diabetes Care.* 2010;33(6):1304-8.
- Becker C, Jick SS, Meier CR. Risk of stroke in patients with idiopathic Parkinson disease. *Parkinsonism Relat*

Disord. 2010;16(1):31-5.

- Becker C, Brobert GP, Johansson S, Jick SS, Meier CR. Cancer risk in association with Parkinson disease: a population-based study. *Parkinsonism Relat Disord.* 2010;16(3):186-90.
- Azoulay L, Schneider-Lindner V, Dell'aniello S, Schiffrin A, Suissa S. Combination therapy with sulfonylureas and metformin and the prevention of death in type 2 diabetes: a nested case-control study. *Pharmacoepidemiol Drug Saf.* 2010;19(4):335-42.
- Azoulay L, Schneider-Lindner V, Dell'aniello S, Filion KB, Suissa S. Thiazolidinediones and the risk of incident strokes in patients with type 2 diabetes: a nested case-control study. *Pharmacoepidemiol Drug Saf.* 2010;19(4):343-50.
- Armstrong RG, West J, Card TR. Risk of cancer in inflammatory bowel disease treated with azathioprine: a UK population-based case-control study. *Am J Gastroenterol.* 2010;105(7):1604-9.
- Andersohn F, Waring M, Garbe E. Risk of ischemic stroke in patients with Crohn's disease: a population-based nested case-control study. *Inflamm Bowel Dis.* 2010;16(8):1387-92.
- Andersohn F, Schade R, Willich SN, Garbe E. Use of antiepileptic drugs in epilepsy and the risk of self-harm or suicidal behavior. *Neurology.* 2010;75(4):335-40.
- Alonso A, Logroscino G, Jick SS, Hernán MA. Association of smoking with amyotrophic lateral sclerosis risk and survival in men and women: a prospective study. *BMC neurology.* 2010;10:6.
- Abuabara K, Azfar RS, Shin DB, Neimann AL, Troxel AB, Gelfand JM. Cause-specific mortality in patients with severe psoriasis: a population-based cohort study in the U.K. *Br J Dermatol.* 2010;163(3):586-92.
- Zhang B, de Vries F, Setakis E, van Staa T-P. The pattern of risk of myocardial infarction in patients taking asthma medication: a study with the General Practice Research Database. *Journal of Hypertension.* 2009;27(7):1485-92.
- Wong ICK, Asherson P, Bilbow A, Clifford S, Coghill D, DeSoysa R, et al. Cessation of attention deficit hyperactivity disorder drugs in the young (CADDY)--a pharmacoepidemiological and qualitative study. *Health Technol Assess.* 2009;13(50):iii-iv, ix-xi, 1-120.
- Winchester CC, Macfarlane TV, Thomas M, Price D. Antibiotic prescribing and outcomes of lower respiratory tract infection in UK primary care. *Chest.* 2009;135(5):1163-72.
- Watts RA, Al-Taiar A, Scott DGI, Macgregor AJ. Prevalence and incidence of Wegener's granulomatosis in the UK general practice research database. *Arthritis Rheum.* 2009;61(10):1412-6.
- Watts R, Al-Taiar A, Mooney J, Scott D, Macgregor A. The epidemiology of Takayasu arteritis in the UK. *Rheumatology (Oxford).* 2009;48(8):1008-11.
- Watson L, Baird J, Hösel V, Peveler R. The effect of concurrent pain on the management of patients with depression: an analysis of NHS healthcare resource utilisation using the GPRD database. *Int J Clin Pract.* 2009;63(5):698-706.
- van Staa T-P, Leufkens HG, Zhang B, Smeeth L. A comparison of cost effectiveness using data from randomized trials or actual clinical practice: selective cox-2 inhibitors as an example. *PLoS Med.* 2009;6(12):e1000194.

- van Staa TP, Sprafka JM. Study of adverse outcomes in women using testosterone therapy. *Maturitas*. 2009;62(1):76-80.
- Tzoulaki I, Molokhia M, Curcin V, Little MP, Millett CJ, Ng A, et al. Risk of cardiovascular disease and all cause mortality among patients with type 2 diabetes prescribed oral antidiabetes drugs: retrospective cohort study using UK general practice research database. *BMJ*. 2009;339:b4731.
- Turner S, Thomas M, von Ziegenweidt J, Price D. Prescribing trends in asthma: a longitudinal observational study. *Arch Dis Child*. 2009;94(1):16-22.
- Toschke AM, Wolfe CDA, Heuschmann PU, Rudd AG, Gulliford M. Antihypertensive treatment after stroke and all-cause mortality--an analysis of the General Practitioner Research Database (GPRD). *Cerebrovasc Dis*. 2009;28(2):105-11.
- Thompson PL, Gilbert RE, Long PF, Saxena S, Sharland M, Wong ICK. Effect of antibiotics for otitis media on mastoiditis in children: a retrospective cohort study using the United kingdom general practice research database. *Pediatrics*. 2009;123(2):424-30.
- Thomas M, von Ziegenweidt J, Lee AJ, Price D. High-dose inhaled corticosteroids versus add-on long-acting beta-agonists in asthma: an observational study. *J Allergy Clin Immunol*. 2009;123(1):116-21.e10.
- Thomas M, Price D, Chrystyn H, Lloyd A, Williams AE, von Ziegenweidt J. Inhaled corticosteroids for asthma: impact of practice level device switching on asthma control. *BMC Pulm Med*. 2009;9:1.
- Tate AR, Martin AGR, Murray-Thomas T, Anderson SR, Cassell JA. Determining the date of diagnosis--is it a simple matter? The impact of different approaches to dating diagnosis on estimates of delayed care for ovarian cancer in UK primary care. *BMC Med Res Methodol*. 2009;9:42.
- Tannen RL, Weiner MG, Xie D. Use of primary care electronic medical record database in drug efficacy research on cardiovascular outcomes: comparison of database and randomised controlled trial findings. *BMJ*. 2009;338:b81.
- Sundström A, Seaman H, Kieler H, Alfredsson L. The risk of venous thromboembolism associated with the use of tranexamic acid and other drugs used to treat menorrhagia: a case-control study using the General Practice Research Database. *BJOG*. 2009;116(1):91-7.
- Stowe J, Andrews N, Wise L, Miller E. Investigation of the temporal association of Guillain-Barre syndrome with influenza vaccine and influenzalike illness using the United Kingdom General Practice Research Database. *Am J Epidemiol*. 2009;169(3):382-8.
- Somers EC, Thomas SL, Smeeth L, Hall AJ. Are individuals with an autoimmune disease at higher risk of a second autoimmune disorder? *Am J Epidemiol*. 2009;169(6):749-55.
- Simon TA, Smitten AL, Franklin J, Askling J, Lacaille D, Wolfe F, et al. Malignancies in the rheumatoid arthritis abatacept clinical development programme: an epidemiological assessment. *Annals of the Rheumatic Diseases*. 2009;68(12):1819-26.
- Schoonen WM, Kucera G, Coalson J, Li L, Rutstein M, Mowat F, et al. Epidemiology of immune thrombocytopenic purpura in the General Practice Research Database. *Br J Haematol*. 2009;145(2):235-44.
- Schneider C, Jick SS, Meier CR. Risk of gynecological cancers in users of estradiol/dydrogesterone or other HRT preparations. *Climacteric*. 2009;12(6):514-24.

- Schneider C, Jick SS, Meier CR. Risk of cardiovascular outcomes in users of estradiol/dydrogesterone or other HRT preparations. *Climacteric*. 2009;12(5):445-53.
- Ruigómez A, Wallander M-A, Johansson S, Rodríguez LAG. Irritable bowel syndrome and gastroesophageal reflux disease in primary care: is there a link? *Dig Dis Sci*. 2009;54(5):1079-86.
- Ruigómez A, Massó-González EL, Johansson S, Wallander M-A, García-Rodríguez LA. Chest pain without established ischaemic heart disease in primary care patients: associated comorbidities and mortality. *Br J Gen Pract*. 2009;59(560):e78-86.
- Rodríguez LAG, Tolosa LB, Ruigómez A, Johansson S, Wallander MA. Rheumatoid arthritis in UK primary care: incidence and prior morbidity. *Scand J Rheumatol*. 2009;38(3):173-7.
- Rietbrock S, Plumb JM, Gallagher AM, van Staa TP. How effective are dose-adjusted warfarin and aspirin for the prevention of stroke in patients with chronic atrial fibrillation? An analysis of the UK General Practice Research Database. *Thromb Haemost*. 2009;101(3):527-34.
- Rietbrock S, Olson M, van Staa TP. The potential effects on fracture outcomes of improvements in persistence and compliance with bisphosphonates. *QJM*. 2009;102(1):35-42.
- Punekar YS, Sheikh A. Establishing the sequential progression of multiple allergic diagnoses in a UK birth cohort using the General Practice Research Database. *Clin Exp Allergy*. 2009;39(12):1889-95.
- Punekar YS, Sheikh A. Establishing the incidence and prevalence of clinician-diagnosed allergic conditions in children and adolescents using routinely collected data from general practices. *Clin Exp Allergy*. 2009;39(8):1209-16.
- Phatak H, Wentworth C, Sazonov V, Burke T. Prevalence and predictors of lipid abnormalities in patients treated with statins in the UK general practice. *Atherosclerosis*. 2009;202(1):225-33.
- Moore M, Yuen HM, Dunn N, Mullee MA, Maskell J, Kendrick T. Explaining the rise in antidepressant prescribing: a descriptive study using the general practice research database. *BMJ*. 2009;339:b3999.
- Millett C, Bottle A, Ng A, Curcin V, Molokhia M, Saxena S, et al. Pay for performance and the quality of diabetes management in individuals with and without co-morbid medical conditions. *J R Soc Med*. 2009;102(9):369-77.
- McCarthy S, Cranswick N, Potts L, Taylor E, Wong ICK. Mortality associated with attention-deficit hyperactivity disorder (ADHD) drug treatment: a retrospective cohort study of children, adolescents and young adults using the general practice research database. *Drug Safety*. 2009;32(11):1089-96.
- McCarthy S, Asherson P, Coghill D, Hollis C, Murray M, Potts L, et al. Attention-deficit hyperactivity disorder: treatment discontinuation in adolescents and young adults. *Br J Psychiatry*. 2009;194(3):273-7.
- Martinez C, Sullivan T, Hauser WA. Prevalence of acute repetitive seizures (ARS) in the United Kingdom. *Epilepsy Res*. 2009;87(2-3):137-43.
- Kernick D, Stapley S, Campbell J, Hamilton W. What happens to new-onset headache in children that present to primary care? A case-cohort study using electronic primary care records. *Cephalalgia: An International Journal of Headache*. 2009;29(12):1311-6.
- Jones R, Charlton J, Latinovic R, Gulliford MC. Alarm symptoms and identification of non-cancer diagnoses in primary care: cohort study. *BMJ*. 2009;339:b3094.

- Jick SS, Hagberg KW, Kaye JA, Jick H. The risk of unintended pregnancies in users of the contraceptive patch compared to users of oral contraceptives in the UK General Practice Research Database. *Contraception*. 2009;80(2):142-51.
- Jick H, Hagberg KW, Egger P. Rate of suicide in patients taking montelukast. *Pharmacotherapy*. 2009;29(2):165-6.
- Jick H, Chamberlin DP, Hagberg KW. The origin and spread of a mumps epidemic: United kingdom, 2003-2006. *Epidemiology*. 2009;20(5):656-61.
- Humes DJ, Solaymani-Dodaran M, Fleming KM, Simpson J, Spiller RC, West J. A population-based study of perforated diverticular disease incidence and associated mortality. *Gastroenterology*. 2009;136(4):1198-205.
- Hsia Y, Neubert AC, Rani F, Viner RM, Hindmarsh PC, Wong ICK. An increase in the prevalence of type 1 and 2 diabetes in children and adolescents: results from prescription data from a UK general practice database. *British Journal of Clinical Pharmacology*. 2009;67(2):242-9.
- Hsia Y, Maclennan K. Rise in psychotropic drug prescribing in children and adolescents during 1992-2001: a population-based study in the UK. *Eur J Epidemiol*. 2009;24(4):211-6.
- Heckbert SR, Wiggins KL, Glazer NL, Dublin S, Psaty BM, Smith NL, et al. Antihypertensive treatment with ACE inhibitors or beta-blockers and risk of incident atrial fibrillation in a general hypertensive population. *Am J Hypertens*. 2009;22(5):538-44.
- Hamilton WT, Gallagher AM, Thomas JM, White PD. Risk markers for both chronic fatigue and irritable bowel syndromes: a prospective case-control study in primary care. *Psychol Med*. 2009;39(11):1913-21.
- Gunnell D, Irvine D, Wise L, Davies C, Martin RM. Varenicline and suicidal behaviour: a cohort study based on data from the General Practice Research Database. *BMJ*. 2009;339:b3805.
- Gulliford MC, Charlton J, Ashworth M, Rudd AG, Toshke AM, e CRTRT. Selection of medical diagnostic codes for analysis of electronic patient records. Application to stroke in a primary care database. *PLoS One*. 2009;4(9):e7168.
- Gulliford MC, Charlton J. Is relative mortality of type 2 diabetes mellitus decreasing? *Am J Epidemiol*. 2009;169(4):455-61.
- Gulliford M, Latinovic R, Charlton J, Little P, van Staa T, Ashworth M. Selective decrease in consultations and antibiotic prescribing for acute respiratory tract infections in UK primary care up to 2006. *J Public Health (Oxf)*. 2009;31(4):512-20.
- Grosso A, Douglas I, Hingorani AD, MacAllister R, Hubbard R, Smeeth L. Inhaled tiotropium bromide and risk of stroke. *British Journal of Clinical Pharmacology*. 2009;68(5):731-6.
- Grosso A, Douglas I, Hingorani A, MacAllister R, Smeeth L. Oral bisphosphonates and risk of atrial fibrillation and flutter in women: a self-controlled case-series safety analysis. *PLoS One*. 2009;4(3):e4720.
- Gribbin J, Hubbard R, Smith C, Gladman J, Lewis S. Incidence and mortality of falls amongst older people in primary care in the United Kingdom. *QJM*. 2009;102(7):477-83.
- Gelfand JM, Dommasch ED, Shin DB, Azfar RS, Kurd SK, Wang X, et al. The risk of stroke in patients with psoriasis. *J Invest Dermatol*. 2009;129(10):2411-8.

- Gauthier A, Breuer J, Carrington D, Martin M, Rémy V. Epidemiology and cost of herpes zoster and post-herpetic neuralgia in the United Kingdom. *Epidemiol Infect.* 2009;137(1):38-47.
- García Rodríguez LA, Wallander M-A, Tolosa LB, Johansson S. Chronic obstructive pulmonary disease in UK primary care: incidence and risk factors. *COPD.* 2009;6(5):369-79.
- Gallagher AM, Leighton-Scott J, van Staa TP. Utilization characteristics and treatment persistence in patients prescribed low-dose buprenorphine patches in primary care in the United Kingdom: a retrospective cohort study. *Clin Ther.* 2009;31(8):1707-15.
- Frisher M, Martino O, Crome I, Croft P. Trends in drug misuse recorded in primary care in the UK from 1998 to 2005. *J Public Health (Oxf).* 2009;31(1):69-73.
- Frisher M, Crome I, Martino O, Croft P. Assessing the impact of cannabis use on trends in diagnosed schizophrenia in the United Kingdom from 1996 to 2005. *Schizophr Res.* 2009;113(2-3):123-8.
- Flory JH, Ellenberg S, Szapary PO, Strom BL, Hennessy S. Antidiabetic action of bezafibrate in a large observational database. *Diabetes Care.* 2009;32(4):547-51.
- Fillion KB, Joseph L, Boivin J-F, Suissa S, Brophy JM. Trends in the prescription of anti-diabetic medications in the United Kingdom: a population-based analysis. *Pharmacoepidemiol Drug Saf.* 2009;18(10):973-6.
- Feudjo-Tepie MA, Le Roux G, Beach KJ, Bennett D, Robinson NJ. Comorbidities of idiopathic thrombocytopenic purpura: a population-based study. *Adv Hematol.* 2009;2009:963506.
- Feudjo-Tepie MA, Hall SA, Logie J, Robinson NJ. Risk of cataract among idiopathic thrombocytopenic purpura patients in the UK general practice research database. *Pharmacoepidemiol Drug Saf.* 2009;18(5):380-5.
- Evans HER, Tsourapas A, Mercer CH, Rait G, Bryan S, Hamill M, et al. Primary care consultations and costs among HIV-positive individuals in UK primary care 1995-2005: a cohort study. *Sex Transm Infect.* 2009;85(7):543-9.
- Evans HER, Mercer CH, Rait G, Hamill M, Delpech V, Hughes G, et al. Trends in HIV testing and recording of HIV status in the UK primary care setting: a retrospective cohort study 1995-2005. *Sex Transm Infect.* 2009;85(7):520-6.
- El-Serag H, Hill C, Jones R. Systematic review: the epidemiology of gastro-oesophageal reflux disease in primary care, using the UK General Practice Research Database. *Aliment Pharmacol Ther.* 2009;29(5):470-80.
- Douglas IJ, Evans SJ, Pocock S, Smeeth L. The risk of fractures associated with thiazolidinediones: a self-controlled case-series study. *PLoS Med.* 2009;6(9):e1000154.
- Delaney JAC, Daskalopoulou SS, Suissa S. Traditional versus marginal structural models to estimate the effectiveness of beta-blocker use on mortality after myocardial infarction. *Pharmacoepidemiol Drug Saf.* 2009;18(1):1-6.
- de Vries F, Cooper AL, Cockle SM, van Staa TP, Cooper C. Fracture risk in patients receiving acid-suppressant medication alone and in combination with bisphosphonates. *Osteoporos Int.* 2009;20(12):1989-98.
- De Natale R, Lafuma A, Berdeaux G. Cost effectiveness of travoprost versus a fixed combination of latanoprost/timolol in patients with ocular hypertension or glaucoma: analysis based on the UK general practitioner research database. *Clin Drug Investig.* 2009;29(2):111-20.

- Crooks CJ, West J, Solaymani-Dodaran M, Card TR. The epidemiology of haemochromatosis: a population-based study. *Aliment Pharmacol Ther.* 2009;29(2):183-92.
- Brauchli YB, Jick SS, Miret M, Meier CR. Psoriasis and risk of incident cancer: an inception cohort study with a nested case-control analysis. *J Invest Dermatol.* 2009;129(11):2604-12.
- Brauchli YB, Jick SS, Miret M, Meier CR. Psoriasis and risk of incident myocardial infarction, stroke or transient ischaemic attack: an inception cohort study with a nested case-control analysis. *Br J Dermatol.* 2009;160(5):1048-56.
- Brauchli YB, Jick SS, Curtin F, Meier CR. Lithium, antipsychotics, and risk of psoriasis. *J Clin Psychopharmacol.* 2009;29(2):134-40.
- Bodmer M, Brauchli YB, Krähenbühl S, Jick SS, Meier CR. Statin use and risk of gallstone disease followed by cholecystectomy. *JAMA.* 2009;302(18):2001-7.
- Bernier M-O, Mikaeloff Y, Hudson M, Suissa S. Combined oral contraceptive use and the risk of systemic lupus erythematosus. *Arthritis Rheum.* 2009;61(4):476-81.
- Andersohn F, Schade R, Suissa S, Garbe E. Long-term use of antidepressants for depressive disorders and the risk of diabetes mellitus. *Am J Psychiatry.* 2009;166(5):591-8.
- Alonso A, Rodríguez LAG, Logroscino G, Hernán MA. Use of antidepressants and the risk of Parkinson's disease: a prospective study. *J Neurol Neurosurg Psychiatr.* 2009;80(6):671-4.
- Alonso A, Logroscino G, Jick SS, Hernán MA. Incidence and lifetime risk of motor neuron disease in the United Kingdom: a population-based study. *Eur J Neurol.* 2009;16(6):745-51.
- Ackers R, Besag FMC, Wade A, Murray ML, Wong ICK. Changing trends in antiepileptic drug prescribing in girls of child-bearing potential. *Arch Dis Child.* 2009;94(6):443-7.
- Yurgin N, Secnik K, Lage MJ. Obesity and the use of insulin: a study of patients with type 2 diabetes in the UK. *J Diabetes Complicat.* 2008;22(4):235-40.
- Yang Y-X, Hennessy S, Propert K, Hwang W-T, Sarkar M, Lewis JD. Chronic statin therapy and the risk of colorectal cancer. *Pharmacoepidemiol Drug Saf.* 2008;17(9):869-76.
- Willemen MJ, Mantel-Teeuwisse AK, Straus SM, Leufkens HG, Egberts AC. Psychiatric and cardiovascular comorbidities in patients with diabetes mellitus starting antiobesity drugs. *Obesity (Silver Spring).* 2008;16(10):2331-5.
- Weiner MG, Xie D, Tannen RL. Replication of the Scandinavian Simvastatin Survival Study using a primary care medical record database prompted exploration of a new method to address unmeasured confounding. *Pharmacoepidemiol Drug Saf.* 2008;17(7):661-70.
- Weiner MG, Barnhart K, Xie D, Tannen RL. Hormone therapy and coronary heart disease in young women. *Menopause.* 2008;15(1):86-93.
- Walters K, Rait G, Petersen I, Williams R, Nazareth I. Panic disorder and risk of new onset coronary heart disease, acute myocardial infarction, and cardiac mortality: cohort study using the general practice research database. *Eur Heart J.* 2008;29(24):2981-8.
- van Staa TP, Smeeth L, Persson I, Parkinson J, Leufkens HGM. Evaluating drug toxicity signals: is a hierarchical

classification of evidence useful or a hindrance? *Pharmacoepidemiol Drug Saf.* 2008;17(5):475-84.

- van Staa TP, Smeeth L, Persson I, Parkinson J, Leufkens HGM. What is the harm-benefit ratio of Cox-2 inhibitors? *Int J Epidemiol.* 2008;37(2):405-13.
- van Staa TP, Rietbrock S, Setakis E, Leufkens HGM. Does the varied use of NSAIDs explain the differences in the risk of myocardial infarction? *J Intern Med.* 2008;264(5):481-92.
- van Staa TP, Cooper C, Barlow D, Leufkens HGM. Individualizing the risks and benefits of postmenopausal hormone therapy. *Menopause.* 2008;15(2):374-81.
- Thompson PL, Gilbert RE, Long PF, Saxena S, Sharland M, Wong ICK. Has UK guidance affected general practitioner antibiotic prescribing for otitis media in children? *J Public Health (Oxf).* 2008;30(4):479-86.
- Thomas SL, Edwards CJ, Smeeth L, Cooper C, Hall AJ. How accurate are diagnoses for rheumatoid arthritis and juvenile idiopathic arthritis in the general practice research database? *Arthritis Rheum.* 2008;59(9):1314-21.
- Tannen RL, Weiner MG, Xie D. Replicated studies of two randomized trials of angiotensin-converting enzyme inhibitors: further empiric validation of the 'prior event rate ratio' to adjust for unmeasured confounding by indication. *Pharmacoepidemiol Drug Saf.* 2008;17(7):671-85.
- Solaymani-Dodaran M, Aithal GP, Card T, West J. Risk of cardiovascular and cerebrovascular events in primary biliary cirrhosis: a population-based cohort study. *Am J Gastroenterol.* 2008;103(11):2784-8.
- Shawe J, Mulnier H, Nicholls P, Lawrenson R. Use of hormonal contraceptive methods by women with diabetes. *Primary Care Diabetes.* 2008;2(4):195-9.
- Setakis E, Leufkens HGM, van Staa TP. Changes in the characteristics of patients prescribed selective cyclooxygenase 2 inhibitors after the 2004 withdrawal of rofecoxib. *Arthritis Rheum.* 2008;59(8):1105-11.
- Seaman HE, Ballard KD, Wright JT, de Vries CS. Endometriosis and its coexistence with irritable bowel syndrome and pelvic inflammatory disease: findings from a national case-control study--Part 2. *BJOG.* 2008;115(11):1392-6.
- Rietbrock S, Heeley E, Plumb J, van Staa T. Chronic atrial fibrillation: Incidence, prevalence, and prediction of stroke using the Congestive heart failure, Hypertension, Age >75, Diabetes mellitus, and prior Stroke or transient ischemic attack (CHADS2) risk stratification scheme. *Am Heart J.* 2008;156(1):57-64.
- Renoux C, Dell'aniello S, Garbe E, Suissa S. Hormone replacement therapy use and the risk of stroke. *Maturitas.* 2008;61(4):305-9.
- Rani F, Murray ML, Byrne PJ, Wong ICK. Epidemiologic features of antipsychotic prescribing to children and adolescents in primary care in the United Kingdom. *Pediatrics.* 2008;121(5):1002-9.
- Phatak H, Wentworth C, Burke TA. Lipid testing among patients beginning statin therapy in general practice in the United Kingdom. *Value Health.* 2008;11(5):933-8.
- Osborn D, Levy G, Nazareth I, King M. Suicide and severe mental illnesses. Cohort study within the UK general practice research database. *Schizophr Res.* 2008;99(1-3):134-8.
- Opatrny L, Delaney JAC, Suissa S. Gastro-intestinal haemorrhage risks of selective serotonin receptor antagonist therapy: a new look. *British Journal of Clinical Pharmacology.* 2008;66(1):76-81.

- Nitsch D, Douglas I, Smeeth L, Fletcher A. Age-related macular degeneration and complement activation-related diseases: a population-based case-control study. *Ophthalmology*. 2008;115(11):1904-10.
- Mulnier HE, Seaman HE, Raleigh VS, Soedamah-Muthu SS, Colhoun HM, Lawrenson RA, et al. Risk of myocardial infarction in men and women with type 2 diabetes in the UK: a cohort study using the General Practice Research Database. *Diabetologia*. 2008;51(9):1639-45.
- Molokhia M, McKeigue P, Curcin V, Majeed A. Statin induced myopathy and myalgia: time trend analysis and comparison of risk associated with statin class from 1991-2006. *PloS One*. 2008;3(6):e2522.
- Mikaeloff Y, Kezouh A, Suissa S. Nonsteroidal anti-inflammatory drug use and the risk of severe skin and soft tissue complications in patients with varicella or zoster disease. *British Journal of Clinical Pharmacology*. 2008;65(2):203-9.
- Meropol SB, Chan KA, Chen Z, Finkelstein JA, Hennessy S, Lautenbach E, et al. Adverse events associated with prolonged antibiotic use. *Pharmacoepidemiol Drug Saf*. 2008;17(5):523-32.
- Meier C, Kraenzlin ME, Bodmer M, Jick SS, Jick H, Meier CR. Use of thiazolidinediones and fracture risk. *Arch Intern Med*. 2008;168(8):820-5.
- Lewis JD, Gelfand JM, Troxel AB, Forde KA, Newcomb C, Kim H, et al. Immunosuppressant medications and mortality in inflammatory bowel disease. *Am J Gastroenterol*. 2008;103(6):1428-35; quiz 36.
- Lafuma A, Laurendeau C, Berdeaux G. Costs and persistence of brimonidine versus brinzolamide in everyday glaucoma care: an analysis conducted on the UK General Practitioner Research Database. *J Med Econ*. 2008;11(3):485-97.
- Lafuma A, Berdeaux G. Costs and persistence of carbonic anhydrase inhibitor versus alpha-2 agonists, associated with beta-blockers, in glaucoma and ocular hypertension: an analysis of the UK-GPRD database. *Current Medical Research and Opinion*. 2008;24(5):1519-27.
- Kaye JA, Jick H. Proton pump inhibitor use and risk of hip fractures in patients without major risk factors. *Pharmacotherapy*. 2008;28(8):951-9.
- Jick SS, Li L. Antidepressant drug use and risk of venous thromboembolism. *Pharmacotherapy*. 2008;28(2):144-50.
- Huerta C, Johansson S, Wallander M-A, Rodríguez LAG. Risk of myocardial infarction and overall mortality in survivors of venous thromboembolism. *Thromb J*. 2008;6:10.
- Hammad TA, McAdams MA, Feight A, Iyasu S, Dal Pan GJ. Determining the predictive value of Read/OXMIS codes to identify incident acute myocardial infarction in the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2008;17(12):1197-201.
- Hammad TA, Graham DJ, Staffa JA, Kornegay CJ, Dal Pan GJ. Onset of acute myocardial infarction after use of non-steroidal anti-inflammatory drugs. *Pharmacoepidemiol Drug Saf*. 2008;17(4):315-21.
- Griffin J, Lee S, Caiado M, Kesten S, Price D. Comparison of tiotropium bromide and combined ipratropium/salbutamol for the treatment of COPD: a UK General Practice Research Database 12-month follow-up study. *Prim Care Respir J*. 2008;17(2):104-10.
- García-Rodríguez LA, Massó-González EL, Wallander M-A, Johansson S. The safety of rosuvastatin in comparison with other statins in over 100,000 statin users in UK primary care. *Pharmacoepidemiol Drug Saf*.

2008;17(10):943-52.

- García Rodríguez LA, Ruigómez A, Martín-Merino E, Johansson S, Wallander M-A. Relationship between gastroesophageal reflux disease and COPD in UK primary care. *Chest*. 2008;134(6):1223-30.
- Gallagher AM, Rietbrock S, Plumb J, van Staa TP. Initiation and persistence of warfarin or aspirin in patients with chronic atrial fibrillation in general practice: do the appropriate patients receive stroke prophylaxis? *J Thromb Haemost*. 2008;6(9):1500-6.
- Gallagher AM, Rietbrock S, Olson M, van Staa TP. Fracture outcomes related to persistence and compliance with oral bisphosphonates. *J Bone Miner Res*. 2008;23(10):1569-75.
- Douglas JJ, Smeeth L. Exposure to antipsychotics and risk of stroke: self controlled case series study. *BMJ*. 2008;337:a1227.
- Devine S, West SL, Andrews E, Tennis P, Eaton S, Thorp J, et al. Validation of neural tube defects in the full featured--general practice research database. *Pharmacoepidemiol Drug Saf*. 2008;17(5):434-44.
- Denis P, Lafuma A, Berdeaux G. Costs and persistence of alpha-2 adrenergic agonists versus carbonic anhydrase inhibitors, both associated with prostaglandin analogues, for glaucoma as recorded by The United Kingdom General Practitioner Research Database. *Clin Ophthalmol*. 2008;2(2):321-9.
- Delaney JAC, Schneider-Lindner V, Brassard P, Suissa S. Mortality after infection with methicillin-resistant *Staphylococcus aureus* (MRSA) diagnosed in the community. *BMC Med*. 2008;6:2.
- Delaney JAC, Moodie EEM, Suissa S. Validating the effects of drug treatment on blood pressure in the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2008;17(6):535-45.
- de Abajo FJ, García-Rodríguez LA. Risk of upper gastrointestinal tract bleeding associated with selective serotonin reuptake inhibitors and venlafaxine therapy: interaction with nonsteroidal anti-inflammatory drugs and effect of acid-suppressing agents. *Arch Gen Psychiatry*. 2008;65(7):795-803.
- Daskalopoulou SS, Delaney JAC, Filion KB, Brophy JM, Mayo NE, Suissa S. Discontinuation of statin therapy following an acute myocardial infarction: a population-based study. *Eur Heart J*. 2008;29(17):2083-91.
- Choi CW, Kim GH, Song CS, Wang SG, Lee BJ, I H, et al. Is obesity associated with gastropharyngeal reflux disease? *World J Gastroenterol*. 2008;14(2):265-71.
- Cheeta S, Drummond C, Oyefeso A, Phillips T, Deluca P, Perryman K, et al. Low identification of alcohol use disorders in general practice in England. *Addiction*. 2008;103(5):766-73.
- Charlton RA, Cunningham MC, de Vries CS, Weil JG. Data resources for investigating drug exposure during pregnancy and associated outcomes: the General Practice Research Database (GPRD) as an alternative to pregnancy registries. *Drug Safety*. 2008;31(1):39-51.
- Charlton J, Latinovic R, Gulliford MC. Explaining the decline in early mortality in men and women with type 2 diabetes: a population-based cohort study. *Diabetes Care*. 2008;31(9):1761-6.
- Cardwell CR, McKinney PA, Patterson CC, Murray LJ. Infections in early life and childhood leukaemia risk: a UK case-control study of general practitioner records. *Br J Cancer*. 2008;99(9):1529-33.
- Cardwell CR, Carson DJ, Patterson CC. No association between routinely recorded infections in early life and subsequent risk of childhood-onset Type 1 diabetes: a matched case-control study using the UK General

Practice Research Database. *Diabet Med.* 2008;25(3):261-7.

- Card TR, Solaymani-Dodaran M, West J. Incidence and mortality of primary sclerosing cholangitis in the UK: a population-based cohort study. *J Hepatol.* 2008;48(6):939-44.
- Bremner SA, Carey IM, DeWilde S, Richards N, Maier WC, Hilton SR, et al. Infections presenting for clinical care in early life and later risk of hay fever in two UK birth cohorts. *Allergy.* 2008;63(3):274-83.
- Brauchli YB, Jick SS, Meier CR. Psoriasis and the risk of incident diabetes mellitus: a population-based study. *Br J Dermatol.* 2008;159(6):1331-7.
- Brauchli YB, Jick SS, Curtin F, Meier CR. Association between beta-blockers, other antihypertensive drugs and psoriasis: population-based case-control study. *Br J Dermatol.* 2008;158(6):1299-307.
- Brauchli YB, Jick SS, Curtin F, Meier CR. Association between use of thiazolidinediones or other oral antidiabetics and psoriasis: A population based case-control study. *J Am Acad Dermatol.* 2008;58(3):421-9.
- Becker C, Jick SS, Meier CR. Use of statins and the risk of Parkinson's disease: a retrospective case-control study in the UK. *Drug Safety.* 2008;31(5):399-407.
- Becker C, Brobert GP, Johansson S, Jick SS, Meier CR. Diabetes in patients with idiopathic Parkinson's disease. *Diabetes Care.* 2008;31(9):1808-12.
- Becker C, Brobert GP, Almqvist PM, Johansson S, Jick SS, Meier CR. The risk of newly diagnosed asthma in migraineurs with or without previous triptan prescriptions. *Headache.* 2008;48(4):606-10.
- Becker C, Brobert GP, Almqvist PM, Johansson S, Jick SS, Meier CR. Migraine incidence, comorbidity and health resource utilization in the UK. *Cephalgia: An International Journal of Headache.* 2008;28(1):57-64.
- Abel KM, Heatlie HF, Howard LM, Webb RT. Sex- and age-specific incidence of fractures in mental illness: a historical, population-based cohort study. *The Journal of Clinical Psychiatry.* 2008;69(9):1398-403.
- Yang Y-X, Hennessy S, Probert K, Hwang W-T, Sedarat A, Lewis JD. Chronic proton pump inhibitor therapy and the risk of colorectal cancer. *Gastroenterology.* 2007;133(3):748-54.
- Wurst KE, Ephross SA, Loehr J, Clark DW, Guess HA. The utility of the general practice research database to examine selected congenital heart defects: a validation study. *Pharmacoepidemiol Drug Saf.* 2007;16(8):867-77.
- Wurst KE, Ephross SA, Loehr J, Clark DW, Guess HA. Evaluation of the General Practice Research Database congenital heart defects prevalence: comparison to United Kingdom national systems. *Birth Defects Res Part A Clin Mol Teratol.* 2007;79(4):309-16.
- Wilting I, de Vries F, Thio BMKS, Cooper C, Heerdink ER, Leufkens HGM, et al. Lithium use and the risk of fractures. *Bone.* 2007;40(5):1252-8.
- Wallander M-A, Johansson S, Ruigómez A, García Rodríguez LA, Jones R. Morbidity associated with sleep disorders in primary care: a longitudinal cohort study. *Prim Care Companion J Clin Psychiatry.* 2007;9(5):338-45.
- Wallander MA, Johansson S, Ruigómez A, García Rodríguez LA. Unspecified abdominal pain in primary care: the role of gastrointestinal morbidity. *Int J Clin Pract.* 2007;61(10):1663-70.
- Velthuis-Te Wierik EJM, Hendricks PT, Martinez C. Preferential prescribing of tibolone and combined

estrogen plus progestogen therapy in postmenopausal women. *Menopause*. 2007;14(3 Pt 1):518-27.

- van Staa TP, Geusens P, Zhang B, Leufkens HGM, Boonen A, Cooper C. Individual fracture risk and the cost-effectiveness of bisphosphonates in patients using oral glucocorticoids. *Rheumatology (Oxford)*. 2007;46(3):460-6.
- van de Garde EMW, Souverein PC, Hak E, Deneer VHM, van den Bosch JMM, Leufkens HGM. Angiotensin-converting enzyme inhibitor use and protection against pneumonia in patients with diabetes. *Journal of Hypertension*. 2007;25(1):235-9.
- Tannen RL, Weiner MG, Xie D, Barnhart K. A simulation using data from a primary care practice database closely replicated the women's health initiative trial. *J Clin Epidemiol*. 2007;60(7):686-95.
- Tannen RL, Weiner MG, Xie D, Barnhart K. Estrogen affects post-menopausal women differently than estrogen plus progestin replacement therapy. *Hum Reprod*. 2007;22(6):1769-77.
- Srinivasan R, Yang Y-X, Rubin SC, Morgan MA, Lewis JD. Risk of colorectal cancer in women with a prior diagnosis of gynecologic malignancy. *J Clin Gastroenterol*. 2007;41(3):291-6.
- Somers EC, Thomas SL, Smeeth L, Schoonen WM, Hall AJ. Incidence of systemic lupus erythematosus in the United Kingdom, 1990-1999. *Arthritis Rheum*. 2007;57(4):612-8.
- Smitten AL, Choi HK, Hochberg MC, Suissa S, Simon TA, Testa MA, et al. The risk of herpes zoster in patients with rheumatoid arthritis in the United States and the United Kingdom. *Arthritis Rheum*. 2007;57(8):1431-8.
- Sharland M, Subgroup SP. The use of antibacterials in children: a report of the Specialist Advisory Committee on Antimicrobial Resistance (SACAR) Paediatric Subgroup. *J Antimicrob Chemother*. 2007;60 Suppl 1:i15-26.
- Seaman HE, Langley SEM, Farmer RDT, de Vries CS. Venous thromboembolism and cyproterone acetate in men with prostate cancer: a study using the General Practice Research Database. *BJU international*. 2007;99(6):1398-403.
- Schlienger RG, Fedson DS, Jick SS, Jick H, Meier CR. Statins and the risk of pneumonia: a population-based, nested case-control study. *Pharmacotherapy*. 2007;27(3):325-32.
- Schade R, Andersohn F, Suissa S, Haverkamp W, Garbe E. Dopamine agonists and the risk of cardiac-valve regurgitation. *N Engl J Med*. 2007;356(1):29-38.
- Ruigómez A, Rodríguez LAG, Wallander MA, Johansson S, Dent J. Endoscopic findings in a cohort of newly diagnosed gastroesophageal reflux disease patients registered in a UK primary care database. *Dis Esophagus*. 2007;20(6):504-9.
- Ruigómez A, García Rodríguez LA, Johansson S, Wallander M-A, Edvardsson N. Risk of cerebrovascular accident after a first diagnosis of atrial fibrillation. *Clin Cardiol*. 2007;30(12):624-8.
- Rubino A, Roskell N, Tennis P, Mines D, Weich S, Andrews E. Risk of suicide during treatment with venlafaxine, citalopram, fluoxetine, and dothiepin: retrospective cohort study. *BMJ*. 2007;334(7587):242.
- Pitman RJ, Melegaro A, Gelb D, Siddiqui MR, Gay NJ, Edmunds WJ. Assessing the burden of influenza and other respiratory infections in England and Wales. *J Infect*. 2007;54(6):530-8.
- Petersen I, Johnson AM, Islam A, Duckworth G, Livermore DM, Hayward AC. Protective effect of antibiotics against serious complications of common respiratory tract infections: retrospective cohort study with the UK

General Practice Research Database. *BMJ*. 2007;335(7627):982.

- Petersen I, Hayward AC, Subgroup SS. Antibacterial prescribing in primary care. *J Antimicrob Chemother*. 2007;60 Suppl 1:i43-7.
- Osborn DPJ, Levy G, Nazareth I, Petersen I, Islam A, King MB. Relative risk of cardiovascular and cancer mortality in people with severe mental illness from the United Kingdom's General Practice Research Database. *Arch Gen Psychiatry*. 2007;64(2):242-9.
- Nightingale AL, Farmer RDT, de Vries CS. Systemic lupus erythematosus prevalence in the UK: methodological issues when using the General Practice Research Database to estimate frequency of chronic relapsing-remitting disease. *Pharmacoepidemiol Drug Saf*. 2007;16(2):144-51.
- Margolis DJ, Hoffstad O, Isseroff RR. Association between the use of beta-adrenergic receptor agents and the development of venous leg ulcers. *Arch Dermatol*. 2007;143(10):1275-80.
- MacDonald TM, Morant SV, Mozaffari E. Drug treatment discontinuation and achievement of target blood pressure and cholesterol in United Kingdom primary care. *Current Medical Research and Opinion*. 2007;23(11):2765-74.
- Lafuma A, Berdeaux G. Costs and effectiveness of travoprost versus a dorzolamide + timolol fixed combination in first-line treatment of glaucoma: analysis conducted on the United Kingdom General Practitioner Research Database. *Current Medical Research and Opinion*. 2007;23(12):3009-16.
- Jordan K, Clarke AM, Symmons DPM, Fleming D, Porcheret M, Kadam UT, et al. Measuring disease prevalence: a comparison of musculoskeletal disease using four general practice consultation databases. *Br J Gen Pract*. 2007;57(534):7-14.
- Jones R, Latinovic R, Charlton J, Gulliford MC. Alarm symptoms in early diagnosis of cancer in primary care: cohort study using General Practice Research Database. *BMJ*. 2007;334(7602):1040.
- Jick SS, Kaye JA, Jick H. Diclofenac and acute myocardial infarction in patients with no major risk factors. *British Journal of Clinical Pharmacology*. 2007;64(5):662-7.
- Jackson H, Solaymani-Dodaran M, Card TR, Aithal GP, Logan R, West J. Influence of ursodeoxycholic acid on the mortality and malignancy associated with primary biliary cirrhosis: a population-based cohort study. *Hepatology*. 2007;46(4):1131-7.
- Hughes G, Williams T, Simms I, Mercer C, Fenton K, Cassell J. Use of a primary care database to determine trends in genital chlamydia testing, diagnostic episodes and management in UK general practice, 1990-2004. *Sex Transm Infect*. 2007;83(4):310-3.
- Huerta C, Rivero E, Rodríguez LAG. Incidence and risk factors for psoriasis in the general population. *Arch Dermatol*. 2007;143(12):1559-65.
- Huerta C, Johansson S, Wallander M-A, García Rodríguez LA. Risk factors and short-term mortality of venous thromboembolism diagnosed in the primary care setting in the United Kingdom. *Arch Intern Med*. 2007;167(9):935-43.
- Howard L, Kirkwood G, Leese M. Risk of hip fracture in patients with a history of schizophrenia. *Br J Psychiatry*. 2007;190:129-34.
- Höglér W, Wehl G, van Staa T, Meister B, Klein-Franke A, Kropshofer G. Incidence of skeletal complications

during treatment of childhood acute lymphoblastic leukemia: comparison of fracture risk with the General Practice Research Database. *Pediatr Blood Cancer*. 2007;48(1):21-7.

- Hennessy S, Bilker WB, Leonard CE, Chittams J, Palumbo CM, Karlawish JH, et al. Observed association between antidepressant use and pneumonia risk was confounded by comorbidity measures. *J Clin Epidemiol*. 2007;60(9):911-8.
- Hamilton W, Kernick D. Clinical features of primary brain tumours: a case-control study using electronic primary care records. *Br J Gen Pract*. 2007;57(542):695-9.
- Gelfand JM, Troxel AB, Lewis JD, Kurd SK, Shin DB, Wang X, et al. The risk of mortality in patients with psoriasis: results from a population-based study. *Arch Dermatol*. 2007;143(12):1493-9.
- Fillion KB, Delaney JAC, Brophy JM, Ernst P, Suissa S. The impact of over-the-counter simvastatin on the number of statin prescriptions in the United Kingdom: a view from the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2007;16(1):1-4.
- Edwards CJ, Cooper C, Fisher D, Field M, van Staa TP, Arden NK. The importance of the disease process and disease-modifying antirheumatic drug treatment in the development of septic arthritis in patients with rheumatoid arthritis. *Arthritis Rheum*. 2007;57(7):1151-7.
- Dunn N, Lane D, Everitt H, Little P. Use of antibiotics for sore throat and incidence of quinsy. *Br J Gen Pract*. 2007;57(534):45-9.
- Douglas JJ, Cook C, Chakravarthy U, Hubbard R, Fletcher AE, Smeeth L. A case-control study of drug risk factors for age-related macular degeneration. *Ophthalmology*. 2007;114(6):1164-9.
- Delaney JAC, Daskalopoulou SS, Brophy JM, Steele RJ, Opatrny L, Suissa S. Lifestyle variables and the risk of myocardial infarction in the general practice research database. *BMC Cardiovasc Disord*. 2007;7:38.
- Delaney JA, Opatrny L, Brophy JM, Suissa S. Drug drug interactions between antithrombotic medications and the risk of gastrointestinal bleeding. *CMAJ*. 2007;177(4):347-51.
- de Vries F, Souverein PC, Cooper C, Leufkens HGM, van Staa TP. Use of beta-blockers and the risk of hip/femur fracture in the United Kingdom and The Netherlands. *Calcif Tissue Int*. 2007;80(2):69-75.
- De Vries F, Bracke M, Leufkens HGM, Lammers J-WJ, Cooper C, Van Staa TP. Fracture risk with intermittent high-dose oral glucocorticoid therapy. *Arthritis Rheum*. 2007;56(1):208-14.
- Cook MN, Girman CJ, Stein PP, Alexander CM. Initial monotherapy with either metformin or sulphonylureas often fails to achieve or maintain current glycaemic goals in patients with Type 2 diabetes in UK primary care. *Diabet Med*. 2007;24(4):350-8.
- Burke TA, Sturkenboom MC, Ohman-Strickland PA, Wentworth CE, Rhoads GG. The effect of antihypertensive drugs and drug combinations on the incidence of new-onset type-2 diabetes mellitus. *Pharmacoepidemiol Drug Saf*. 2007;16(9):979-87.
- Bremner SA, Carey IM, DeWilde S, Richards N, Maier WC, Hilton SR, et al. Vaccinations, infections and antibacterials in the first grass pollen season of life and risk of later hayfever. *Clin Exp Allergy*. 2007;37(4):512-7.
- Bowe WP, Hoffstad O, Margolis DJ. Upper respiratory tract infection in household contacts of acne patients. *Dermatology (Basel)*. 2007;215(3):213-8.

- Becker C, Brobert GP, Almqvist PM, Johansson S, Jick SS, Meier CR. Migraine and the risk of stroke, TIA, or death in the UK (CME). *Headache*. 2007;47(10):1374-84.
- Alonso A, Rodríguez LAG, Logroscino G, Hernán MA. Gout and risk of Parkinson disease: a prospective study. *Neurology*. 2007;69(17):1696-700.
- Alonso A, Jick SS, Olek MJ, Hernán MA. Incidence of multiple sclerosis in the United Kingdom : findings from a population-based cohort. *J Neurol*. 2007;254(12):1736-41.
- Ackers R, Murray ML, Besag FMC, Wong ICK. Prioritizing children's medicines for research: a pharmaco-epidemiological study of antiepileptic drugs. *British Journal of Clinical Pharmacology*. 2007;63(6):689-97.
- Aberra FN, Stettler N, Brensinger C, Lichtenstein GR, Lewis JD. Risk for active tuberculosis in inflammatory bowel disease patients. *Clin Gastroenterol Hepatol*. 2007;5(9):1070-5.
- Williamson I, Benge S, Mullee M, Little P. Consultations for middle ear disease, antibiotic prescribing and risk factors for reattendance: a case-linked cohort study. *Br J Gen Pract*. 2006;56(524):170-5.
- West J, Logan RF, Hubbard RB, Card TR. Risk of schizophrenia in people with coeliac disease, ulcerative colitis and Crohn's disease: a general population-based study. *Aliment Pharmacol Ther*. 2006;23(1):71-4.
- Verdel BM, Souverein PC, Egberts AC, Leufkens HG. Difference in risks of allergic reaction to sulfonamide drugs based on chemical structure. *Ann Pharmacother*. 2006;40(6):1040-6.
- van Staa TP, Geusens P, Bijlsma JWJ, Leufkens HGM, Cooper C. Clinical assessment of the long-term risk of fracture in patients with rheumatoid arthritis. *Arthritis Rheum*. 2006;54(10):3104-12.
- van de Garde EMW, Hak E, Souverein PC, Hoes AW, van den Bosch JMM, Leufkens HGM. Statin treatment and reduced risk of pneumonia in patients with diabetes. *Thorax*. 2006;61(11):957-61.
- Tannen RL, Weiner MG, Marcus SM. Simulation of the Syst-Eur randomized control trial using a primary care electronic medical record was feasible. *J Clin Epidemiol*. 2006;59(3):254-64.
- Tam CC, Rodrigues LC, Petersen I, Islam A, Hayward A, O'Brien SJ. Incidence of Guillain-Barré syndrome among patients with *Campylobacter* infection: a general practice research database study. *J Infect Dis*. 2006;194(1):95-7.
- Stowe J, Andrews N, Wise L, Miller E. Bell's palsy and parenteral inactivated influenza vaccine. *Hum Vaccin*. 2006;2(3):110-2.
- Souverein PC, Webb DJ, Weil JG, Van Staa TP, Egberts ACG. Use of antiepileptic drugs and risk of fractures: case-control study among patients with epilepsy. *Neurology*. 2006;66(9):1318-24.
- Solaymani-Dodaran M, Card TR, Aithal GP, West J. Fracture risk in people with primary biliary cirrhosis: a population-based cohort study. *Gastroenterology*. 2006;131(6):1752-7.
- Soedamah-Muthu SS, Fuller JH, Mulnier HE, Raleigh VS, Lawrenson RA, Colhoun HM. High risk of cardiovascular disease in patients with type 1 diabetes in the U.K.: a cohort study using the general practice research database. *Diabetes Care*. 2006;29(4):798-804.
- Soedamah-Muthu SS, Fuller JH, Mulnier HE, Raleigh VS, Lawrenson RA, Colhoun HM. All-cause mortality rates in patients with type 1 diabetes mellitus compared with a non-diabetic population from the UK general practice research database, 1992-1999. *Diabetologia*. 2006;49(4):660-6.

- Smith S, Smith GE, Heatlie H, Bashford JNR, Ashcroft DM, Verlander NQ, et al. Reducing variation in antibacterial prescribing rates for 'cough/cold' and sore throat between 1993 and 2001: regional analyses using the general practice research database. *Public Health*. 2006;120(8):752-9.
- Smeeth L, Cook C, Hall AJ. Incidence of diagnosed polymyalgia rheumatica and temporal arteritis in the United Kingdom, 1990-2001. *Annals of the Rheumatic Diseases*. 2006;65(8):1093-8.
- Shah AD, Martinez C. An algorithm to derive a numerical daily dose from unstructured text dosage instructions. *Pharmacoepidemiol Drug Saf*. 2006;15(3):161-6.
- Ruigómez A, Rodríguez LAG, Wallander M-A, Johansson S, Jones R. Chest pain in general practice: incidence, comorbidity and mortality. *Fam Pract*. 2006;23(2):167-74.
- Ruigómez A, García Rodríguez LA, Wallander M-A, Johansson S, Eklund S. Esophageal stricture: incidence, treatment patterns, and recurrence rate. *Am J Gastroenterol*. 2006;101(12):2685-92.
- Ruigómez A, García Rodríguez LA, Wallander M-A, Johansson S, Dent J. Comparison of gastro-oesophageal reflux disease and heartburn diagnoses in UK primary care. *Current Medical Research and Opinion*. 2006;22(9):1661-8.
- Odeyemi IaO, Dakin HA, O'Donnell RA, Warner J, Jacobs A, Dasgupta P. Epidemiology, prescribing patterns and resource use associated with overactive bladder in UK primary care. *Int J Clin Pract*. 2006;60(8):949-58.
- Noakes K, Pebody RG, Gungabissoon U, Stowe J, Miller E. Pneumococcal polysaccharide vaccine uptake in England, 1989-2003, prior to the introduction of a vaccination programme for older adults. *J Public Health (Oxf)*. 2006;28(3):242-7.
- Nightingale AL, Farmer RDT, de Vries CS. Incidence of clinically diagnosed systemic lupus erythematosus 1992-1998 using the UK General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2006;15(9):656-61.
- Mulnier HE, Seaman HE, Raleigh VS, Soedamah-Muthu SS, Colhoun HM, Lawrenson RA. Mortality in people with type 2 diabetes in the UK. *Diabet Med*. 2006;23(5):516-21.
- Latinovic R, Gulliford MC, Hughes RaC. Incidence of common compressive neuropathies in primary care. *J Neurol Neurosurg Psychiatr*. 2006;77(2):263-5.
- Latinovic R, Gulliford M, Ridsdale L. Headache and migraine in primary care: consultation, prescription, and referral rates in a large population. *J Neurol Neurosurg Psychiatr*. 2006;77(3):385-7.
- Kim J, Evans S, Smeeth L, Pocock S. Hormone replacement therapy and acute myocardial infarction: a large observational study exploring the influence of age. *Int J Epidemiol*. 2006;35(3):731-8.
- Jones R, Latinovic R, Charlton J, Gulliford M. Physical and psychological co-morbidity in irritable bowel syndrome: a matched cohort study using the General Practice Research Database. *Aliment Pharmacol Ther*. 2006;24(5):879-86.
- Jick SS, Lieberman ES, Rahman MU, Choi HK. Glucocorticoid use, other associated factors, and the risk of tuberculosis. *Arthritis Rheum*. 2006;55(1):19-26.
- Jick H, Kaye JA, Russmann S, Jick SS. Nonsteroidal antiinflammatory drugs and acute myocardial infarction in patients with no major risk factors. *Pharmacotherapy*. 2006;26(10):1379-87.

- Hughes RA, Charlton J, Latinovic R, Gulliford MC. No association between immunization and Guillain-Barré syndrome in the United Kingdom, 1992 to 2000. *Arch Intern Med.* 2006;166(12):1301-4.
- Huerta C, Varas-Lorenzo C, Castellsague J, García Rodríguez LA. Non-steroidal anti-inflammatory drugs and risk of first hospital admission for heart failure in the general population. *Heart.* 2006;92(11):1610-5.
- Hernández-Díaz S, García Rodríguez LA. Cardioprotective aspirin users and their excess risk of upper gastrointestinal complications. *BMC Med.* 2006;4:22.
- Hernán MA, Logroscino G, García Rodríguez LA. Nonsteroidal anti-inflammatory drugs and the incidence of Parkinson disease. *Neurology.* 2006;66(7):1097-9.
- Hardy JR, Leaderer BP, Holford TR, Hall GC, Bracken MB. Safety of medications prescribed before and during early pregnancy in a cohort of 81,975 mothers from the UK General Practice Research Database. *Pharmacoepidemiol Drug Saf.* 2006;15(8):555-64.
- Handoko KB, Souverein PC, van Staa TP, Meyboom RHB, Leufkens HGM, Egberts TCG, et al. Risk of aplastic anemia in patients using antiepileptic drugs. *Epilepsia.* 2006;47(7):1232-6.
- Hall GC, Carroll D, Parry D, McQuay HJ. Epidemiology and treatment of neuropathic pain: the UK primary care perspective. *Pain.* 2006;122(1-2):156-62.
- Gupta G, Lautenbach E, Lewis JD. Incidence and risk factors for herpes zoster among patients with inflammatory bowel disease. *Clin Gastroenterol Hepatol.* 2006;4(12):1483-90.
- Gore M, Dukes E, Rowbotham D, Tai K-S, Leslie DL. Prevalence of contraindicated medical conditions and use of precluded medications in patients with painful neuropathic disorders prescribed amitriptyline. *Pain Pract.* 2006;6(4):265-72.
- González-Pérez A, Rodríguez LAG. Upper gastrointestinal complications among users of paracetamol. *Basic Clin Pharmacol Toxicol.* 2006;98(3):297-303.
- Gelfand JM, Shin DB, Neimann AL, Wang X, Margolis DJ, Troxel AB. The risk of lymphoma in patients with psoriasis. *J Invest Dermatol.* 2006;126(10):2194-201.
- Gelfand JM, Neimann AL, Shin DB, Wang X, Margolis DJ, Troxel AB. Risk of myocardial infarction in patients with psoriasis. *JAMA.* 2006;296(14):1735-41.
- García Rodríguez LA, Ruigómez A, Panés J. Acute gastroenteritis is followed by an increased risk of inflammatory bowel disease. *Gastroenterology.* 2006;130(6):1588-94.
- Fox KM, Gerber Pharmd RA, Bolinder B, Chen J, Kumar S. Prevalence of inadequate glycemic control among patients with type 2 diabetes in the United Kingdom general practice research database: A series of retrospective analyses of data from 1998 through 2002. *Clin Ther.* 2006;28(3):388-95.
- de Vries F, de Vries C, Cooper C, Leufkens B, van Staa T-P. Reanalysis of two studies with contrasting results on the association between statin use and fracture risk: the General Practice Research Database. *Int J Epidemiol.* 2006;35(5):1301-8.
- de Vries CS, Bromley SE, Farmer RDT. Myocardial infarction risk and hormone replacement: differences between products. *Maturitas.* 2006;53(3):343-50.
- Cassell JA, Mercer CH, Sutcliffe L, Petersen I, Islam A, Brook MG, et al. Trends in sexually transmitted

infections in general practice 1990-2000: population based study using data from the UK general practice research database. *BMJ*. 2006;332(7537):332-4.

- Carlson C, Hornbuckle K, DeLisle F, Kryzhanovskaya L, Breier A, Cavazzoni P. Diabetes mellitus and antipsychotic treatment in the United Kingdom. *Eur Neuropsychopharmacol*. 2006;16(5):366-75.
- Card TR, Solaymani-Dodaran M, Hubbard R, Logan RFA, West J. Is an internal comparison better than using national data when estimating mortality in longitudinal studies? *J Epidemiol Community Health*. 2006;60(9):819-21.
- Burnham JM, Shults J, Weinstein R, Lewis JD, Leonard MB. Childhood onset arthritis is associated with an increased risk of fracture: a population based study using the General Practice Research Database. *Annals of the Rheumatic Diseases*. 2006;65(8):1074-9.
- Burke TA, Sturkenboom MC, Lu S-e, Wentworth CE, Lin Y, Rhoads GG. Discontinuation of antihypertensive drugs among newly diagnosed hypertensive patients in UK general practice. *Journal of Hypertension*. 2006;24(6):1193-200.
- Brankin E, Walker M, Lynch N, Aspray T, Lis Y, Cowell W. The impact of dosing frequency on compliance and persistence with bisphosphonates among postmenopausal women in the UK: evidence from three databases. *Current Medical Research and Opinion*. 2006;22(7):1249-56.
- Ashworth M, Charlton J, Latinovic R, Gulliford M. Age-related changes in consultations and antibiotic prescribing for acute respiratory infections, 1995-2000. Data from the UK General Practice Research Database. *J Clin Pharm Ther*. 2006;31(5):461-7.
- Arana A, Varas C, González-Pérez A, Gutiérrez L, Bjerrum L, García Rodríguez LA. Hormone therapy and cerebrovascular events: a population-based nested case-control study. *Menopause*. 2006;13(5):730-6.
- Apter AJ, Kinman JL, Bilker WB, Herlim M, Margolis DJ, Lautenbach E, et al. Is there cross-reactivity between penicillins and cephalosporins? *Am J Med*. 2006;119(4):354.e11-19.
- Andersohn F, Suissa S, Garbe E. Use of first- and second-generation cyclooxygenase-2-selective nonsteroidal antiinflammatory drugs and risk of acute myocardial infarction. *Circulation*. 2006;113(16):1950-7.
- Andersohn F, Schade R, Suissa S, Garbe E. Cyclooxygenase-2 selective nonsteroidal anti-inflammatory drugs and the risk of ischemic stroke: a nested case-control study. *Stroke*. 2006;37(7):1725-30.
- Alonso A, Jick SS, Jick H, Hernán MA. Antibiotic use and risk of multiple sclerosis. *Am J Epidemiol*. 2006;163(11):997-1002.
- Alonso A, Jick SS, Hernán MA. Allergy, histamine 1 receptor blockers, and the risk of multiple sclerosis. *Neurology*. 2006;66(4):572-5.
- Aina FO, Smeeth L, Hubbard R, Hurt LS, Fletcher AE. Hormone replacement therapy and cataract: a population-based case-control study. *Eye (Lond)*. 2006;20(4):417-22.
- Yang Y-X, Hennessy S, Lewis JD. Type 2 diabetes mellitus and the risk of colorectal cancer. *Clin Gastroenterol Hepatol*. 2005;3(6):587-94.
- Watson L, Kerstjens HAM, Rabe KF, Kiri V, Visick GT, Postma DS. Obtaining optimal control in mild asthma: theory and practice. *Fam Pract*. 2005;22(3):305-10.

- Wade AG, Toumi I, Hemels MEH. A probabilistic cost-effectiveness analysis of escitalopram, generic citalopram and venlafaxine as a first-line treatment of major depressive disorder in the UK. *Current Medical Research and Opinion*. 2005;21(4):631-42.
- van Staa TP, Geusens P, Pols HaP, de Laet C, Leufkens HGM, Cooper C. A simple score for estimating the long-term risk of fracture in patients using oral glucocorticoids. *QJM*. 2005;98(3):191-8.
- van Staa TP, Card T, Logan RF, Leufkens HGM. 5-Aminosalicylate use and colorectal cancer risk in inflammatory bowel disease: a large epidemiological study. *Gut*. 2005;54(11):1573-8.
- Tata LJ, West J, Smith C, Farrington P, Card T, Smeeth L, et al. General population based study of the impact of tricyclic and selective serotonin reuptake inhibitor antidepressants on the risk of acute myocardial infarction. *Heart*. 2005;91(4):465-71.
- Sullivan CO, Omar RZ, Ambler G, Majeed A. Case-mix and variation in specialist referrals in general practice. *Br J Gen Pract*. 2005;55(516):529-33.
- Souverein PC, Webb DJ, Petri H, Weil J, Van Staa TP, Egberts T. Incidence of fractures among epilepsy patients: a population-based retrospective cohort study in the General Practice Research Database. *Epilepsia*. 2005;46(2):304-10.
- Soriano JB, Visick GT, Muellerova H, Payvandi N, Hansell AL. Patterns of comorbidities in newly diagnosed COPD and asthma in primary care. *Chest*. 2005;128(4):2099-107.
- Solaymani-Dodaran M, Logan RFA, West J, Card T. Mortality associated with Barrett's esophagus and gastroesophageal reflux disease diagnoses-a population-based cohort study. *Am J Gastroenterol*. 2005;100(12):2616-21.
- Smeeth L, Cook C, Chakravarthy U, Hubbard R, Fletcher AE. A case control study of age related macular degeneration and use of statins. *Br J Ophthalmol*. 2005;89(9):1171-5.
- Sharland M, Kendall H, Yeates D, Randall A, Hughes G, Glasziou P, et al. Antibiotic prescribing in general practice and hospital admissions for peritonsillar abscess, mastoiditis, and rheumatic fever in children: time trend analysis. *BMJ*. 2005;331(7512):328-9.
- Shao T, Yang Y-X. Cholecystectomy and the risk of colorectal cancer. *Am J Gastroenterol*. 2005;100(8):1813-20.
- Ryan R, Newnham A, Khunti K, Majeed A. New cases of diabetes mellitus in England and Wales, 1994-1998: database study. *Public Health*. 2005;119(10):892-9.
- Russmann S, Kaye JA, Jick SS, Jick H. Risk of cholestatic liver disease associated with flucloxacillin and flucloxacillin prescribing habits in the UK: cohort study using data from the UK General Practice Research Database. *British Journal of Clinical Pharmacology*. 2005;60(1):76-82.
- Ruigómez A, Rodríguez LAG, Wallander M-A, Johansson S, Thomas M, Price D. Gastroesophageal reflux disease and asthma: a longitudinal study in UK general practice. *Chest*. 2005;128(1):85-93.
- Ruigómez A, Johansson S, Wallander M-A, García Rodríguez LA. Predictors and prognosis of paroxysmal atrial fibrillation in general practice in the UK. *BMC Cardiovasc Disord*. 2005;5:20.
- Mines D, Hill D, Yu H, Novelli L. Prevalence of risk factors for suicide in patients prescribed venlafaxine, fluoxetine, and citalopram. *Pharmacoepidemiol Drug Saf*. 2005;14(6):367-72.

- Mikuls TR, Farrar JT, Bilker WB, Fernandes S, Schumacher HR, Saag KG. Gout epidemiology: results from the UK General Practice Research Database, 1990-1999. *Annals of the Rheumatic Diseases*. 2005;64(2):267-72.
- Mikuls TR, Farrar JT, Bilker WB, Fernandes S, Saag KG. Suboptimal physician adherence to quality indicators for the management of gout and asymptomatic hyperuricaemia: results from the UK General Practice Research Database (GPRD). *Rheumatology (Oxford)*. 2005;44(8):1038-42.
- Maru S, Koch GG, Stender M, Clark D, Gibowski L, Petri H, et al. Antidiabetic drugs and heart failure risk in patients with type 2 diabetes in the U.K. primary care setting. *Diabetes Care*. 2005;28(1):20-6.
- Martinez C, Rietbrock S, Wise L, Ashby D, Chick J, Moseley J, et al. Antidepressant treatment and the risk of fatal and non-fatal self harm in first episode depression: nested case-control study. *BMJ*. 2005;330(7488):389.
- Margolis DJ, Bowe WP, Hoffstad O, Berlin JA. Antibiotic treatment of acne may be associated with upper respiratory tract infections. *Arch Dermatol*. 2005;141(9):1132-6.
- Logie J, Clifford GM, Farmer RDT. Incidence, prevalence and management of lower urinary tract symptoms in men in the UK. *BJU international*. 2005;95(4):557-62.
- Lewis JD, Bilker WB, Weinstein RB, Strom BL. The relationship between time since registration and measured incidence rates in the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2005;14(7):443-51.
- Lawson A, West J, Aithal GP, Logan RFA. Autoimmune cholestatic liver disease in people with coeliac disease: a population-based study of their association. *Aliment Pharmacol Ther*. 2005;21(4):401-5.
- Koro CE, Bowlin SJ, Weiss SR. Antidiabetic therapy and the risk of heart failure in type 2 diabetic patients: an independent effect or confounding by indication. *Pharmacoepidemiol Drug Saf*. 2005;14(10):697-703.
- Kiri VA, Pride NB, Soriano JB, Vestbo J. Inhaled corticosteroids in chronic obstructive pulmonary disease: results from two observational designs free of immortal time bias. *Am J Respir Crit Care Med*. 2005;172(4):460-4.
- Kiri VA, Bettoncelli G, Testi R, Viegi G. Inhaled corticosteroids are more effective in COPD patients when used with LABA than with SABA. *Respir Med*. 2005;99(9):1115-24.
- Kaye JA, Jick H. Antibiotics and the risk of breast cancer. *Epidemiology*. 2005;16(5):688-90.
- Joseph C, Goddard N, Gelb D. Influenza vaccine uptake and distribution in England and Wales using data from the General Practice Research Database, 1989/90-2003/04. *J Public Health (Oxf)*. 2005;27(4):371-7.
- Hugenholtz GWK, Heerdink ER, van Staa TP, Nolen WA, Egberts ACG. Risk of hip/femur fractures in patients using antipsychotics. *Bone*. 2005;37(6):864-70.
- Huerta C, Lanes SF, García Rodríguez LA. Respiratory medications and the risk of cardiac arrhythmias. *Epidemiology*. 2005;16(3):360-6.
- Huerta C, Castellsague J, Varas-Lorenzo C, García Rodríguez LA. Nonsteroidal anti-inflammatory drugs and risk of ARF in the general population. *Am J Kidney Dis*. 2005;45(3):531-9.
- Hernán MA, Jick SS, Logroscino G, Olek MJ, Ascherio A, Jick H. Cigarette smoking and the progression of multiple sclerosis. *Brain*. 2005;128(Pt 6):1461-5.
- Helms PJ, Ekins Daukes S, Taylor MW, Simpson CR, McLay JS. Utility of routinely acquired primary care data for paediatric disease epidemiology and pharmacoepidemiology. *British Journal of Clinical Pharmacology*.

2005;59(6):684-90.

- Hamilton WT, Gallagher AM, Thomas JM, White PD. The prognosis of different fatigue diagnostic labels: a longitudinal survey. *Fam Pract*. 2005;22(4):383-8.
- Gupta G, Gelfand JM, Lewis JD. Increased risk for demyelinating diseases in patients with inflammatory bowel disease. *Gastroenterology*. 2005;129(3):819-26.
- Gulliford MC, Charlton J, Latinovic R. Trends in antihypertensive and lipid-lowering therapy in subjects with type II diabetes: clinical effectiveness or clinical discretion? *J Hum Hypertens*. 2005;19(2):111-7.
- Gulliford MC, Charlton J, Latinovic R. Increased utilization of primary care 5 years before diagnosis of type 2 diabetes: a matched cohort study. *Diabetes Care*. 2005;28(1):47-52.
- Gulliford MC, Adams G, Ukoumunne OC, Latinovic R, Chinn S, Campbell MJ. Intraclass correlation coefficient and outcome prevalence are associated in clustered binary data. *J Clin Epidemiol*. 2005;58(3):246-51.
- González-Pérez A, García Rodríguez LA. Prostate cancer risk among men with diabetes mellitus (Spain). *Cancer Causes Control*. 2005;16(9):1055-8.
- González-Pérez A, García Rodríguez LA. Breast cancer risk among users of antidepressant medications. *Epidemiology*. 2005;16(1):101-5.
- Gelfand JM, Weinstein R, Porter SB, Neimann AL, Berlin JA, Margolis DJ. Prevalence and treatment of psoriasis in the United Kingdom: a population-based study. *Arch Dermatol*. 2005;141(12):1537-41.
- Gasse C, Hollowell J, Meier CR, Haefeli WE. Drug interactions and risk of acute bleeding leading to hospitalisation or death in patients with chronic atrial fibrillation treated with warfarin. *Thromb Haemost*. 2005;94(3):537-43.
- García Rodríguez LA, González-Pérez A. Use of antibiotics and risk of breast cancer. *Am J Epidemiol*. 2005;161(7):616-9.
- Frisher M, Crome I, Macleod J, Millson D, Croft P. Substance misuse and psychiatric illness: prospective observational study using the general practice research database. *J Epidemiol Community Health*. 2005;59(10):847-50.
- Fleming DM, Ross AM, Cross KW, Kendall H, Elliot AJ. Concerning: 'Why has antibiotic prescribing for respiratory illness declined in primary care? A longitudinal study using the General Practice Research Database'. *J Public Health (Oxf)*. 2005;27(2):228-9; author reply 9-31.
- Fischer LM, Schlienger RG, Matter CM, Jick H, Meier CR. Current use of nonsteroidal antiinflammatory drugs and the risk of acute myocardial infarction. *Pharmacotherapy*. 2005;25(4):503-10.
- Edwards CJ, Arden NK, Fisher D, Saperia JC, Reading I, Van Staa TP, et al. The changing use of disease-modifying anti-rheumatic drugs in individuals with rheumatoid arthritis from the United Kingdom General Practice Research Database. *Rheumatology (Oxford)*. 2005;44(11):1394-8.
- Dunn N, Mullee M, Perry VH, Holmes C. Association between dementia and infectious disease: evidence from a case-control study. *Alzheimer Dis Assoc Disord*. 2005;19(2):91-4.
- Dunn N, Holmes C, Mullee M. Does lithium therapy protect against the onset of dementia? *Alzheimer Dis Assoc Disord*. 2005;19(1):20-2.

- Dial S, Delaney JaC, Barkun AN, Suissa S. Use of gastric acid-suppressive agents and the risk of community-acquired *Clostridium difficile*-associated disease. *JAMA*. 2005;294(23):2989-95.
- de Vries F, van Staa TP, Bracke MSGM, Cooper C, Leufkens HGM, Lammers JWJ. Severity of obstructive airway disease and risk of osteoporotic fracture. *Eur Respir J*. 2005;25(5):879-84.
- de Vries CS, Bromley SE, Thomas H, Farmer RDT. Tibolone and endometrial cancer: a cohort and nested case-control study in the UK. *Drug Safety*. 2005;28(3):241-9.
- de Giuli F, Khaw K-T, Cowie MR, Sutton GC, Ferrari R, Poole-Wilson PA. Incidence and outcome of persons with a clinical diagnosis of heart failure in a general practice population of 696,884 in the United Kingdom. *Eur J Heart Fail*. 2005;7(3):295-302.
- Currin L, Schmidt U, Treasure J, Jick H. Time trends in eating disorder incidence. *Br J Psychiatry*. 2005;186:132-5.
- Cook MN, Girman CJ, Stein PP, Alexander CM, Holman RR. Glycemic control continues to deteriorate after sulfonylureas are added to metformin among patients with type 2 diabetes. *Diabetes Care*. 2005;28(5):995-1000.
- Clifford GM, Logie J, Farmer RDT. No risk of drug-associated liver injury with alpha1-adrenoreceptor blocking agents in men with BPH: results from an observational study using the GPRD. *Pharmacoepidemiol Drug Saf*. 2005;14(2):75-80.
- Bradbury BD, Wilk JB, Kaye JA. Obesity and the risk of prostate cancer (United States). *Cancer Causes Control*. 2005;16(6):637-41.
- Bradbury BD, Lash TL, Kaye JA, Jick SS. Tamoxifen-treated breast carcinoma patients and the risk of acute myocardial infarction and newly-diagnosed angina. *Cancer*. 2005;103(6):1114-21.
- Black C, Kaye JA, Jick H. Cesarean delivery in the United Kingdom: time trends in the general practice research database. *Obstet Gynecol*. 2005;106(1):151-5.
- Ashworth M, Charlton J, Ballard K, Latinovic R, Gulliford M. Variations in antibiotic prescribing and consultation rates for acute respiratory infection in UK general practices 1995-2000. *Br J Gen Pract*. 2005;55(517):603-8.
- Alonso A, Jick SS, Olek MJ, Ascherio A, Jick H, Hernán MA. Recent use of oral contraceptives and the risk of multiple sclerosis. *Arch Neurol*. 2005;62(9):1362-5.
- Aberra FN, Brensinger CM, Bilker WB, Lichtenstein GR, Lewis JD. Antibiotic use and the risk of flare of inflammatory bowel disease. *Clin Gastroenterol Hepatol*. 2005;3(5):459-65.
- Zhou Z, Althin R, Sforzolini BS, Dhawan R. Persistency and treatment failure in newly diagnosed open angle glaucoma patients in the United Kingdom. *Br J Ophthalmol*. 2004;88(11):1391-4.
- Yang Y-X, Hennessy S, Lewis JD. Insulin therapy and colorectal cancer risk among type 2 diabetes mellitus patients. *Gastroenterology*. 2004;127(4):1044-50.
- West J, Logan RFA, Smith CJ, Hubbard RB, Card TR. Malignancy and mortality in people with coeliac disease: population based cohort study. *BMJ*. 2004;329(7468):716-9.
- West J, Logan RFA, Card TR, Smith C, Hubbard R. Risk of vascular disease in adults with diagnosed coeliac

disease: a population-based study. *Aliment Pharmacol Ther.* 2004;20(1):73-9.

- Varney SJ, Guest JF. Relative cost effectiveness of Depo-Provera, Implanon, and Mirena in reversible long-term hormonal contraception in the UK. *Pharmacoeconomics.* 2004;22(17):1141-51.
- van Staa T-P, Bishop N, Leufkens HGM, Cooper C. Are inhaled corticosteroids associated with an increased risk of fracture in children? *Osteoporos Int.* 2004;15(10):785-91.
- Van Staa TP, Travis S, Leufkens HGM, Logan RF. 5-aminosalicylic acids and the risk of renal disease: a large British epidemiologic study. *Gastroenterology.* 2004;126(7):1733-9.
- Van De Vijver DAMC, Walley T, Petri H. Epidemiology of restless legs syndrome as diagnosed in UK primary care. *Sleep Med.* 2004;5(5):435-40.
- Torgerson DJ, Sykes D, Puffer S, Brown P, Cooper C. Pharmaceutical treatment of symptomatic vertebral fractures in primary care. *Annals of the Rheumatic Diseases.* 2004;63(7):853-6.
- Souverein PC, Berard A, Van Staa TP, Cooper C, Egberts ACG, Leufkens HGM, et al. Use of oral glucocorticoids and risk of cardiovascular and cerebrovascular disease in a population based case-control study. *Heart.* 2004;90(8):859-65.
- Solaymani-Dodaran M, Logan RFA, West J, Card T, Coupland C. Risk of extra-oesophageal malignancies and colorectal cancer in Barrett's oesophagus and gastro-oesophageal reflux. *Scand J Gastroenterol.* 2004;39(7):680-5.
- Smith GE, Smith S, Heatlie H, Bashford JNR, Hawker J, Ashcroft D, et al. What has happened to antimicrobial usage in primary care in the United Kingdom since the SMAC report? - description of trends in antimicrobial usage using the General Practice Research Database. *J Public Health (Oxf).* 2004;26(4):359-64.
- Smeeth L, Thomas SL, Hall AJ, Hubbard R, Farrington P, Vallance P. Risk of myocardial infarction and stroke after acute infection or vaccination. *N Engl J Med.* 2004;351(25):2611-8.
- Smeeth L, Cook C, Fombonne PE, Heavey L, Rodrigues LC, Smith PG, et al. Rate of first recorded diagnosis of autism and other pervasive developmental disorders in United Kingdom general practice, 1988 to 2001. *BMC Med.* 2004;2:39.
- Smeeth L, Cook C, Fombonne E, Heavey L, Rodrigues LC, Smith PG, et al. MMR vaccination and pervasive developmental disorders: a case-control study. *Lancet.* 2004;364(9438):963-9.
- Seaman HE, de Vries CS, Farmer RDT. Venous thromboembolism associated with cyproterone acetate in combination with ethinyloestradiol (Dianette): observational studies using the UK General Practice Research Database. *Pharmacoepidemiol Drug Saf.* 2004;13(7):427-36.
- Schlienger RG, Kraenzlin ME, Jick SS, Meier CR. Use of beta-blockers and risk of fractures. *JAMA.* 2004;292(11):1326-32.
- Schlienger RG, Jick SS, Meier CR. Inhaled corticosteroids and the risk of fractures in children and adolescents. *Pediatrics.* 2004;114(2):469-73.
- Schlienger RG, Fischer LM, Jick H, Meier CR. Current use of selective serotonin reuptake inhibitors and risk of acute myocardial infarction. *Drug Safety.* 2004;27(14):1157-65.
- Ruigómez A, García Rodríguez LA, Wallander MA, Johansson S, Graffner H, Dent J. Natural history of gastro-

oesophageal reflux disease diagnosed in general practice. *Aliment Pharmacol Ther.* 2004;20(7):751-60.

- Ronquist G, Rodríguez LAG, Ruigómez A, Johansson S, Wallander M-A, Frithz G, et al. Association between captopril, other antihypertensive drugs and risk of prostate cancer. *Prostate.* 2004;58(1):50-6.
- Peng MM, Jick H. A population-based study of the incidence, cause, and severity of anaphylaxis in the United Kingdom. *Arch Intern Med.* 2004;164(3):317-9.
- Nightingale AL, Farmer RDT. Ischemic stroke in young women: a nested case-control study using the UK General Practice Research Database. *Stroke.* 2004;35(7):1574-8.
- Murray ML, de Vries CS, Wong ICK. A drug utilisation study of antidepressants in children and adolescents using the General Practice Research Database. *Arch Dis Child.* 2004;89(12):1098-102.
- Morant SV, Pettitt D, MacDonald TM, Burke TA, Goldstein JL. Application of a propensity score to adjust for channelling bias with NSAIDs. *Pharmacoepidemiol Drug Saf.* 2004;13(6):345-53.
- Meier CR, Wilcock K, Jick SS. The risk of severe depression, psychosis or panic attacks with prophylactic antimalarials. *Drug Safety.* 2004;27(3):203-13.
- McKeever TM, Lewis SA, Smith C, Hubbard R. Vaccination and allergic disease: a birth cohort study. *Am J Public Health.* 2004;94(6):985-9.
- Margolis DJ, Knauss J, Bilker W. Medical conditions associated with venous leg ulcers. *Br J Dermatol.* 2004;150(2):267-73.
- Mangtani P, Cumberland P, Hodgson CR, Roberts JA, Cutts FT, Hall AJ. A cohort study of the effectiveness of influenza vaccine in older people, performed using the United Kingdom general practice research database. *J Infect Dis.* 2004;190(1):1-10.
- Lewis JD, Brensinger C. Agreement between GPRD smoking data: a survey of general practitioners and a population-based survey. *Pharmacoepidemiol Drug Saf.* 2004;13(7):437-41.
- Lewis JD, Aberra FN, Lichtenstein GR, Bilker WB, Brensinger C, Strom BL. Seasonal variation in flares of inflammatory bowel disease. *Gastroenterology.* 2004;126(3):665-73.
- Kaye JA, Jick H. Statin use and cancer risk in the General Practice Research Database. *Br J Cancer.* 2004;90(3):635-7.
- Kaye JA, Jick H. Epidemiology of lower limb fractures in general practice in the United Kingdom. *Inj Prev.* 2004;10(6):368-74.
- Jick SS, Bradbury BD. Statins and newly diagnosed diabetes. *British Journal of Clinical Pharmacology.* 2004;58(3):303-9.
- Jick H, Kaye JA, Jick SS. Antidepressants and the risk of suicidal behaviors. *JAMA.* 2004;292(3):338-43.
- Jick H, Kaye JA, Black C. Incidence and prevalence of drug-treated attention deficit disorder among boys in the UK. *Br J Gen Pract.* 2004;54(502):345-7.
- Jick H, Kaye JA. Autism and DPT vaccination in the United Kingdom. *N Engl J Med.* 2004;350(26):2722-3.
- Howard LM, Goss C, Leese M, Appleby L, Thornicroft G. The psychosocial outcome of pregnancy in women with psychotic disorders. *Schizophr Res.* 2004;71(1):49-60.

- Hernán MA, Logroscino G, Rodríguez LAG. A prospective study of alcoholism and the risk of Parkinson's disease. *J Neurol*. 2004;251 Suppl 7:VII14-7.
- Hernán MA, Jick SS, Olek MJ, Jick H. Recombinant hepatitis B vaccine and the risk of multiple sclerosis: a prospective study. *Neurology*. 2004;63(5):838-42.
- Hennessy S, Bilker WB, Knauss JS, Kimmel SE, Margolis DJ, Morrison MF, et al. Comparative cardiac safety of low-dose thioridazine and low-dose haloperidol. *British Journal of Clinical Pharmacology*. 2004;58(1):81-7.
- Hardy JR, Holford TR, Hall GC, Bracken MB. Strategies for identifying pregnancies in the automated medical records of the General Practice Research Database. *Pharmacoepidemiol Drug Saf*. 2004;13(11):749-59.
- Hansell AL, Lam KA, Richardson S, Visick G, Soriano JB. Medical event profiling of COPD patients. *Pharmacoepidemiol Drug Saf*. 2004;13(8):547-55.
- Hall GC, Brown MM, Mo J, MacRae KD. Triptans in migraine: the risks of stroke, cardiovascular disease, and death in practice. *Neurology*. 2004;62(4):563-8.
- González-Pérez A, Ronquist G, García Rodríguez LA. Breast cancer incidence and use of antihypertensive medication in women. *Pharmacoepidemiol Drug Saf*. 2004;13(8):581-5.
- Geoghegan JM, Clark DI, Bainbridge LC, Smith C, Hubbard R. Risk factors in carpal tunnel syndrome. *J Hand Surg Br*. 2004;29(4):315-20.
- García Rodríguez LA, González-Pérez A. Risk of breast cancer among users of aspirin and other anti-inflammatory drugs. *Br J Cancer*. 2004;91(3):525-9.
- García Rodríguez LA, González-Pérez A. Inverse association between nonsteroidal anti-inflammatory drugs and prostate cancer. *Cancer Epidemiol Biomarkers Prev*. 2004;13(4):649-53.
- Gallagher AM, Thomas JM, Hamilton WT, White PD. Incidence of fatigue symptoms and diagnoses presenting in UK primary care from 1990 to 2001. *J R Soc Med*. 2004;97(12):571-5.
- Gaitatzis A, Carroll K, Majeed A, W Sander J. The epidemiology of the comorbidity of epilepsy in the general population. *Epilepsia*. 2004;45(12):1613-22.
- Frisher M, Collins J, Millson D, Crome I, Croft P. Prevalence of comorbid psychiatric illness and substance misuse in primary care in England and Wales. *J Epidemiol Community Health*. 2004;58(12):1036-41.
- Fombonne E, Heavey L, Smeeth L, Rodrigues LC, Cook C, Smith PG, et al. Validation of the diagnosis of autism in general practitioner records. *BMC Public Health*. 2004;4:5.
- Fischer LM, Schlienger RG, Matter CM, Jick H, Meier CR. Discontinuation of nonsteroidal anti-inflammatory drug therapy and risk of acute myocardial infarction. *Arch Intern Med*. 2004;164(22):2472-6.
- Fischer LM, Schlienger RG, Matter C, Jick H, Meier CR. Effect of rheumatoid arthritis or systemic lupus erythematosus on the risk of first-time acute myocardial infarction. *Am J Cardiol*. 2004;93(2):198-200.
- De Wilde S, Carey IM, Bremner SA, Richards N, Hilton SR, Strachan DP, et al. A comparison of the recording of 30 common childhood conditions in the Doctor's Independent Network and General Practice Research Databases. *Health Stat Q*. 2004(22):21-31.
- de Abajo FJ, Montero D, Madurga M, García Rodríguez LA. Acute and clinically relevant drug-induced liver injury: a population based case-control study. *British Journal of Clinical Pharmacology*. 2004;58(1):71-80.

- Cooper C, Dennison EM, Leufkens HGM, Bishop N, van Staa TP. Epidemiology of childhood fractures in Britain: a study using the general practice research database. *J Bone Miner Res.* 2004;19(12):1976-81.
- Cleary P, Shorvon S, Tallis R. Late-onset seizures as a predictor of subsequent stroke. *Lancet.* 2004;363(9416):1184-6.
- Carey IM, Cook DG, De Wilde S, Bremner SA, Richards N, Caine S, et al. Developing a large electronic primary care database (Doctors' Independent Network) for research. *Int J Med Inform.* 2004;73(5):443-53.
- Card T, West J, Hubbard R, Logan RFA. Hip fractures in patients with inflammatory bowel disease and their relationship to corticosteroid use: a population based cohort study. *Gut.* 2004;53(2):251-5.
- Card T, Logan RFA, Rodrigues LC, Wheeler JG. Antibiotic use and the development of Crohn's disease. *Gut.* 2004;53(2):246-50.
- Bromley SE, de Vries CS, Farmer RDT. Utilisation of hormone replacement therapy in the United Kingdom. A descriptive study using the general practice research database. *BJOG.* 2004;111(4):369-76.
- Bradbury BD, Lash TL, Kaye JA, Jick SS. Tamoxifen and cataracts: a null association. *Breast Cancer Res Treat.* 2004;87(2):189-96.
- Bradbury BD, Jick H. In vitro fertilization and childhood retinoblastoma. *British Journal of Clinical Pharmacology.* 2004;58(2):209-11.
- Boyle P, Roehrborn C, Harkaway R, Logie J, de la Rosette J, Emberton M. 5-Alpha reductase inhibition provides superior benefits to alpha blockade by preventing AUR and BPH-related surgery. *European Urology.* 2004;45(5):620-6; discussion 6-7.
- Ashworth M, Latinovic R, Charlton J, Cox K, Rowlands G, Gulliford M. Why has antibiotic prescribing for respiratory illness declined in primary care? A longitudinal study using the General Practice Research Database. *J Public Health (Oxf).* 2004;26(3):268-74.
- Apter AJ, Kinman JL, Bilker WB, Herlim M, Margolis DJ, Lautenbach E, et al. Represcription of penicillin after allergic-like events. *J Allergy Clin Immunol.* 2004;113(4):764-70.
- Andrews N, Miller E, Grant A, Stowe J, Osborne V, Taylor B. Thimerosal exposure in infants and developmental disorders: a retrospective cohort study in the United kingdom does not support a causal association. *Pediatrics.* 2004;114(3):584-91.
- Yang C-C, Jick SS, Testa MA. Discontinuation and switching of therapy after initiation of lipid-lowering drugs: the effects of comorbidities and patient characteristics. *British Journal of Clinical Pharmacology.* 2003;56(1):84-91.
- Yang C-C, Jick SS, Testa MA. Who receives lipid-lowering drugs: the effects of comorbidities and patient characteristics on treatment initiation. *British Journal of Clinical Pharmacology.* 2003;55(3):288-98.
- Yang C-C, Jick SS, Jick H. Lipid-lowering drugs and the risk of depression and suicidal behavior. *Arch Intern Med.* 2003;163(16):1926-32.
- Wood SL, Jick H, Sauve R. The risk of stillbirth in pregnancies before and after the onset of diabetes. *Diabet Med.* 2003;20(9):703-7.
- West J, Logan RFA, Card TR, Smith C, Hubbard R. Fracture risk in people with celiac disease: a population-

based cohort study. *Gastroenterology*. 2003;125(2):429-36.

- Watson DJ, Rhodes T, Guess HA. All-cause mortality and vascular events among patients with rheumatoid arthritis, osteoarthritis, or no arthritis in the UK General Practice Research Database. *J Rheumatol*. 2003;30(6):1196-202.
- van Staa TP, Cooper C, Leufkens HGM, Lammers JW, Suissa S. The use of inhaled corticosteroids in the United Kingdom and the Netherlands. *Respir Med*. 2003;97(5):578-85.
- van Staa TP, Cooper C, Leufkens HGM, Bishop N. Children and the risk of fractures caused by oral corticosteroids. *J Bone Miner Res*. 2003;18(5):913-8.
- van Staa TP, Boulton F, Cooper C, Hagenbeek A, Inskip H, Leufkens HGM. Neutropenia and agranulocytosis in England and Wales: incidence and risk factors. *Am J Hematol*. 2003;72(4):248-54.
- van der Linden PD, Sturkenboom MCJM, Herings RMC, Leufkens HGM, Rowlands S, Stricker BHC. Increased risk of achilles tendon rupture with quinolone antibacterial use, especially in elderly patients taking oral corticosteroids. *Arch Intern Med*. 2003;163(15):1801-7.
- Toledano MB, Hansell AL, Jarup L, Quinn M, Jick S, Elliott P. Temporal trends in orchidopexy, Great Britain, 1992-1998. *Environ Health Perspect*. 2003;111(1):129-32.
- Tata LJ, West J, Harrison T, Farrington P, Smith C, Hubbard R. Does influenza vaccination increase consultations, corticosteroid prescriptions, or exacerbations in subjects with asthma or chronic obstructive pulmonary disease? *Thorax*. 2003;58(10):835-9.
- Strom BL, Schinnar R, Apter AJ, Margolis DJ, Lautenbach E, Hennessy S, et al. Absence of cross-reactivity between sulfonamide antibiotics and sulfonamide nonantibiotics. *N Engl J Med*. 2003;349(17):1628-35.
- Souverein PC, Van Staa TP, Egberts ACG, De la Rosette JJMCH, Cooper C, Leufkens HGM. Use of alpha-blockers and the risk of hip/femur fractures. *J Intern Med*. 2003;254(6):548-54.
- Soriano JB, Kiri VA, Maier WC, Strachan D. Increasing prevalence of asthma in UK primary care during the 1990s. *Int J Tuberc Lung Dis*. 2003;7(5):415-21.
- Soriano JB, Davis KJ, Coleman B, Visick G, Mannino D, Pride NB. The proportional Venn diagram of obstructive lung disease: two approximations from the United States and the United Kingdom. *Chest*. 2003;124(2):474-81.
- Smith S, Smith G, Heatlie H, Bashford J, Ashcroft D, Millson D. Head lice diagnosed in general practice in the West Midlands between 1993 and 2000: a survey using the General Practice Research Database. *Commun Dis Public Health*. 2003;6(2):139-43.
- Smeeth L, Hubbard R, Fletcher AE. Cataract and the use of statins: a case-control study. *QJM*. 2003;96(5):337-43.
- Smeeth L, Boulis M, Hubbard R, Fletcher AE. A population based case-control study of cataract and inhaled corticosteroids. *Br J Ophthalmol*. 2003;87(10):1247-51.
- Seaman HE, de Vries CS, Farmer RDT. Differences in the use of combined oral contraceptives amongst women with and without acne. *Hum Reprod*. 2003;18(3):515-21.
- Seaman HE, de Vries CS, Farmer RDT. The risk of venous thromboembolism in women prescribed

cyproterone acetate in combination with ethinyl estradiol: a nested cohort analysis and case-control study. *Hum Reprod.* 2003;18(3):522-6.

- Seaman HE, de Vries CS, Farmer RD. The risk of liver disorders in women prescribed cyproterone acetate in combination with ethinylloestradiol (Dianette): a nested case-control study using the GPRD. *Pharmacoepidemiol Drug Saf.* 2003;12(7):541-50.
- Ruigómez A, García Rodríguez LA, Johansson S, Wallander M-A. Is hormone replacement therapy associated with an increased risk of irritable bowel syndrome? *Maturitas.* 2003;44(2):133-40.
- Rowlands S, Hannaford P. The incidence of sterilisation in the UK. *BJOG.* 2003;110(9):819-24.
- McInnes GT. The differences between ACE inhibitor-treated and calcium channel blocker-treated hypertensive patients. *J Clin Hypertens (Greenwich).* 2003;5(5):337-44.
- Margolis DJ, Knauss J, Bilker W, Baumgarten M. Medical conditions as risk factors for pressure ulcers in an outpatient setting. *Age Ageing.* 2003;32(3):259-64.
- MacDonald TM, Morant SV, Goldstein JL, Burke TA, Pettitt D. Channelling bias and the incidence of gastrointestinal haemorrhage in users of meloxicam, coxibs, and older, non-specific non-steroidal anti-inflammatory drugs. *Gut.* 2003;52(9):1265-70.
- Kaye JA, Jick H. Incidence of erectile dysfunction and characteristics of patients before and after the introduction of sildenafil in the United Kingdom: cross sectional study with comparison patients. *BMJ.* 2003;326(7386):424-5.
- Kaye JA, Bradbury BD, Jick H. Changes in antipsychotic drug prescribing by general practitioners in the United Kingdom from 1991 to 2000: a population-based observational study. *British Journal of Clinical Pharmacology.* 2003;56(5):569-75.
- Jick SS, Kaye JA, Vasilakis-Scaramozza C, Garcia Rodríguez LA, Ruigómez A, Meier CR, et al. Validity of the general practice research database. *Pharmacotherapy.* 2003;23(5):686-9.
- Jick H, Kaye JA. Epidemiology and possible causes of autism. *Pharmacotherapy.* 2003;23(12):1524-30.
- Huerta C, García Rodríguez LA, Wallander M-A, Johansson S. Users of oral steroids are at a reduced risk of developing irritable bowel syndrome. *Pharmacoepidemiol Drug Saf.* 2003;12(7):583-8.
- Hubbard R, Farrington P, Smith C, Smeeth L, Tattersfield A. Exposure to tricyclic and selective serotonin reuptake inhibitor antidepressants and the risk of hip fracture. *Am J Epidemiol.* 2003;158(1):77-84.
- Howard LM, Goss C, Leese M, Thornicroft G. Medical outcome of pregnancy in women with psychotic disorders and their infants in the first year after birth. *Br J Psychiatry.* 2003;182:63-7.
- Hollowell J, Ruigómez A, Johansson S, Wallander M-A, García-Rodríguez LA. The incidence of bleeding complications associated with warfarin treatment in general practice in the United Kingdom. *Br J Gen Pract.* 2003;53(489):312-4.
- Hoare J, Bruce M, Majeed A. Prevalence of treated asthma and its management in general practice in England and Wales, 1994–1998. *Health Stat Q.* 2003;17:15-22.
- Hoare J. Comparison of area-based inequality measures and disease morbidity in England, 1994–1998. *Health Stat Q.* 2003;18:18-24.

- Gelfand JM, Berlin J, Van Voorhees A, Margolis DJ. Lymphoma rates are low but increased in patients with psoriasis: results from a population-based cohort study in the United Kingdom. *Arch Dermatol.* 2003;139(11):1425-9.
- García Rodríguez LA, Hernández-Díaz S. Nonsteroidal antiinflammatory drugs as a trigger of clinical heart failure. *Epidemiology.* 2003;14(2):240-6.
- Forrest CB, Majeed A, Weiner JP, Carroll K, Bindman AB. Referral of children to specialists in the United States and the United Kingdom. *Arch Pediatr Adolesc Med.* 2003;157(3):279-85.
- Deschaseaux-Voinet C, Lafuma A, Berdeaux G. Cost and effectiveness of brinzolamide versus dorzolamide in current practice: an analysis based on the UK-GPRD database. *J Med Econ.* 2003;6(1-4):69-78.
- Carey IM, Cook DG, De Wilde S, Bremner SA, Richards N, Caine S, et al. Implications of the problem orientated medical record (POMR) for research using electronic GP databases: a comparison of the Doctors Independent Network Database (DIN) and the General Practice Research Database (GPRD). *BMC Fam Pract.* 2003;4:14.
- Card T, Hubbard R, Logan RFA. Mortality in inflammatory bowel disease: a population-based cohort study. *Gastroenterology.* 2003;125(6):1583-90.
- Bremner SA, Carey IM, DeWilde S, Richards N, Maier WC, Hilton SR, et al. Early-life exposure to antibacterials and the subsequent development of hayfever in childhood in the UK: case-control studies using the General Practice Research Database and the Doctors' Independent Network. *Clin Exp Allergy.* 2003;33(11):1518-25.
- Blumentals WA, Brown RR, Gomez-Caminero A. Antihypertensive treatment and erectile dysfunction in a cohort of type II diabetes patients. *Int J Impot Res.* 2003;15(5):314-7.
- Black C, Kaye JA, Jick H. MMR vaccine and idiopathic thrombocytopenic purpura. *British Journal of Clinical Pharmacology.* 2003;55(1):107-11.
- Yang C-C, Jick SS, Jick H. Statins and the risk of idiopathic venous thromboembolism. *British Journal of Clinical Pharmacology.* 2002;53(1):101-5.
- Wyatt KM, Dimmock PW, Frischer M, Jones PW, O'Brien SPM. Prescribing patterns in premenstrual syndrome. *BMC Womens Health.* 2002;2(1):4.
- Wrigley T, Tinto A, Majeed A. Age and sex specific antibiotic prescribing patterns in general practice in England and Wales, 1994 to 1998. *Health Stat Q.* 2002;14:14-20.
- Watson DJ, Rhodes T, Cai B, Guess HA. Lower risk of thromboembolic cardiovascular events with naproxen among patients with rheumatoid arthritis. *Arch Intern Med.* 2002;162(10):1105-10.
- van Staa TP, Selby P, Leufkens HGM, Lyles K, Sprafka JM, Cooper C. Incidence and natural history of Paget's disease of bone in England and Wales. *J Bone Miner Res.* 2002;17(3):465-71.
- van Staa TP, Leufkens HGM, Cooper C. Utility of medical and drug history in fracture risk prediction among men and women. *Bone.* 2002;31(4):508-14.
- van Staa TP, Leufkens HGM, Cooper C. Does a fracture at one site predict later fractures at other sites? A British cohort study. *Osteoporos Int.* 2002;13(8):624-9.
- Soriano JB, Vestbo J, Pride NB, Kiri V, Maden C, Maier WC. Survival in COPD patients after regular use of

fluticasone propionate and salmeterol in general practice. *Eur Respir J.* 2002;20(4):819-25.

- Shorvon SD, Tallis RC, Wallace HK. Antiepileptic drugs: coprescription of proconvulsant drugs and oral contraceptives: a national study of antiepileptic drug prescribing practice. *J Neurol Neurosurg Psychiatr.* 2002;72(1):114-5.
- Sheppard MC, Holder R, Franklyn JA. Levothyroxine treatment and occurrence of fracture of the hip. *Arch Intern Med.* 2002;162(3):338-43.
- Schlienger RG, Jick H, Meier CR. Use of nonsteroidal anti-inflammatory drugs and the risk of first-time acute myocardial infarction. *British Journal of Clinical Pharmacology.* 2002;54(3):327-32.
- Ryan R, Majeed A. Prevalence of treated hypertension in general practice in England and Wales, 1994 to 1998. *Health Stat Q.* 2002(16):14-8.
- Ruigómez A, Johansson S, Wallander M-A, García Rodríguez LA. Risk of mortality in a cohort of patients newly diagnosed with chronic atrial fibrillation. *BMC Cardiovasc Disord.* 2002;2:5.
- Ruigómez A, Johansson S, Wallander MA, Rodríguez LAG. Incidence of chronic atrial fibrillation in general practice and its treatment pattern. *J Clin Epidemiol.* 2002;55(4):358-63.
- Rowlands S, Moser K. Consultation rates from the general practice research database. *Br J Gen Pract.* 2002;52(481):658-60.
- Rowlands S, Hooper R, Hughes R, Burney P. The epidemiology and treatment of Bell's palsy in the UK. *Eur J Neurol.* 2002;9(1):63-7.
- Purcell B, Gaitatzis A, Sander JW, Majeed A. Epilepsy prevalence and prescribing patterns in England and Wales. *Health Stat Q.* 2002;15(Autumn):23-30.
- Newnham A, Ryan R, Khunti K, Majeed A. Prevalence of diagnosed diabetes mellitus in general practice in England and Wales, 1994 to 1998. *Health Stat Q.* 2002(14):5-13.
- Meier CR, Schmitz S, Jick H. Association between acetaminophen or nonsteroidal antiinflammatory drugs and risk of developing ovarian, breast, or colon cancer. *Pharmacotherapy.* 2002;22(3):303-9.
- Meier CR, Krähenbühl S, Schlienger RG, Jick H. Association between body mass index and liver disorders: an epidemiological study. *J Hepatol.* 2002;37(6):741-7.
- McKeever TM, Lewis SA, Smith C, Hubbard R. The importance of prenatal exposures on the development of allergic disease: a birth cohort study using the West Midlands General Practice Database. *Am J Respir Crit Care Med.* 2002;166(6):827-32.
- McKeever TM, Lewis SA, Smith C, Hubbard R. Mode of delivery and risk of developing allergic disease. *J Allergy Clin Immunol.* 2002;109(5):800-2.
- McKeever TM, Lewis SA, Smith C, Collins J, Heatlie H, Frischer M, et al. Early exposure to infections and antibiotics and the incidence of allergic disease: a birth cohort study with the West Midlands General Practice Research Database. *J Allergy Clin Immunol.* 2002;109(1):43-50.
- McCarney R, Fisher P, van Haselen R. Accruing large numbers of patients in primary care trials by retrospective recruitment methods. *Complement Ther Med.* 2002;10(2):63-8.
- Martin RM, Hemingway H, Gunnell D, Karsch KR, Baumbach A, Frankel S. Population need for coronary

revascularisation: are national targets for England credible? *Heart*. 2002;88(6):627-33.

- Margolis DJ, Knauss J, Bilker W. Hormone replacement therapy and prevention of pressure ulcers and venous leg ulcers. *Lancet*. 2002;359(9307):675-7.
- Margolis DJ, Bilker W, Santanna J, Baumgarten M. Venous leg ulcer: incidence and prevalence in the elderly. *J Am Acad Dermatol*. 2002;46(3):381-6.
- Margolis DJ, Bilker W, Knauss J, Baumgarten M, Strom BL. The incidence and prevalence of pressure ulcers among elderly patients in general medical practice. *Ann Epidemiol*. 2002;12(5):321-5.
- Macfarlane A, Hockley C, McCandlish R, McNiece R. Can the General Practice Research Database be used to monitor the health of babies and their mothers? *Health Stat Q*. 2002(13):5-15.
- Lewis JD, Brensinger C, Bilker WB, Strom BL. Validity and completeness of the General Practice Research Database for studies of inflammatory bowel disease. *Pharmacoepidemiol Drug Saf*. 2002;11(3):211-8.
- Lanes SF, García Rodríguez LA, Huerta C. Respiratory medications and risk of asthma death. *Thorax*. 2002;57(8):683-6.
- Koro CE, Fedder DO, L'Italien GJ, Weiss SS, Magder LS, Kreyenbuhl J, et al. Assessment of independent effect of olanzapine and risperidone on risk of diabetes among patients with schizophrenia: population based nested case-control study. *BMJ*. 2002;325(7358):243.
- Koro CE, Fedder DO, L'Italien GJ, Weiss S, Magder LS, Kreyenbuhl J, et al. An assessment of the independent effects of olanzapine and risperidone exposure on the risk of hyperlipidemia in schizophrenic patients. *Arch Gen Psychiatry*. 2002;59(11):1021-6.
- Kornegay CJ, Vasilakis-Scaramozza C, Jick H. Incident diabetes associated with antipsychotic use in the United Kingdom general practice research database. *The Journal of Clinical Psychiatry*. 2002;63(9):758-62.
- Kennerfalk A, Ruigómez A, Wallander M-A, Wilhelmsen L, Johansson S. Geriatric drug therapy and healthcare utilization in the United Kingdom. *Ann Pharmacother*. 2002;36(5):797-803.
- Kaye JA, Meier CR, Walker AM, Jick H. Statin use, hyperlipidaemia, and the risk of breast cancer. *Br J Cancer*. 2002;86(9):1436-9.
- Kang JY, Tinto A, Higham J, Majeed A. Peptic ulceration in general practice in England and Wales 1994-98: period prevalence and drug management. *Aliment Pharmacol Ther*. 2002;16(6):1067-74.
- Johansson S, Wallander M-A, Ruigómez A, García R. Treatment patterns among newly diagnosed heart failure patients in general practice. *Eur J Clin Pharmacol*. 2002;57(11):813-7.
- Huerta C, Zhao SZ, García Rodríguez LA. Risk of acute liver injury in patients with diabetes. *Pharmacotherapy*. 2002;22(9):1091-6.
- Huerta C, García Rodríguez LA, Wallander M-A, Johansson S. Risk of irritable bowel syndrome among asthma patients. *Pharmacoepidemiol Drug Saf*. 2002;11(1):31-5.
- Huerta C, García Rodríguez LA. Risk of clinical blood dyscrasia in a cohort of antibiotic users. *Pharmacotherapy*. 2002;22(5):630-6.
- Hubbard RB, Smith CJP, Smeeth L, Harrison TW, Tattersfield AE. Inhaled corticosteroids and hip fracture: a population-based case-control study. *Am J Respir Crit Care Med*. 2002;166(12 Pt 1):1563-6.

- Hubbard R, Venn A. The impact of coexisting connective tissue disease on survival in patients with fibrosing alveolitis. *Rheumatology (Oxford)*. 2002;41(6):676-9.
- Howard LM, Kumar C, Leese M, Thornicroft G. The general fertility rate in women with psychotic disorders. *Am J Psychiatry*. 2002;159(6):991-7.
- Hajat S, Haines A. Associations of cold temperatures with GP consultations for respiratory and cardiovascular disease amongst the elderly in London. *Int J Epidemiol*. 2002;31(4):825-30.
- Hajat S, Anderson HR, Atkinson RW, Haines A. Effects of air pollution on general practitioner consultations for upper respiratory diseases in London. *Occup Environ Med*. 2002;59(5):294-9.
- Gaitatzis A, Purcell B, Carroll K, Sander JWaS, Majeed A. Differences in the use of health services among people with and without epilepsy in the United Kingdom: socio-economic and disease-specific determinants. *Epilepsy Res*. 2002;50(3):233-41.
- Forrest CB, Majeed A, Weiner JP, Carroll K, Bindman AB. Comparison of specialty referral rates in the United Kingdom and the United States: retrospective cohort analysis. *BMJ*. 2002;325(7360):370-1.
- Davey G, Sedgwick P, Maier W, Visick G, Strachan DP, Anderson HR. Association between migraine and asthma: matched case-control study. *Br J Gen Pract*. 2002;52(482):723-7.
- Clifford GM, Farmer RDT. Drug or symptom-induced depression in men treated with alpha 1-blockers for benign prostatic hyperplasia? A nested case-control study. *Pharmacoepidemiol Drug Saf*. 2002;11(1):55-61.
- Castellsague J, García-Rodríguez L-A, Duque A, Pérez S. Risk of serious skin disorders among users of oral antifungals: a population-based study. *BMC Dermatol*. 2002;2:14.
- Bradbury BD, Jick SS. Itraconazole and fluconazole and certain rare, serious adverse events. *Pharmacotherapy*. 2002;22(6):697-700.
- Black C, Kaye JA, Jick H. Relation of childhood gastrointestinal disorders to autism: nested case-control study using data from the UK General Practice Research Database. *BMJ*. 2002;325(7361):419-21.
- Black C, Kaye JA, Jick H. Clinical risk factors for venous thromboembolism in users of the combined oral contraceptive pill. *British Journal of Clinical Pharmacology*. 2002;53(6):637-40.
- Black C, Jick H. Etiology and frequency of rhabdomyolysis. *Pharmacotherapy*. 2002;22(12):1524-6.
- Bailey K, Ellis C, Majeed A. Trends in the treatment of osteoporosis and types of drug treatments prescribed in general practice in England and Wales, 1994 to 1998. *Health Stat Q*. 2002(16):5-13.
- Vasilakis-Scaramozza C, Jick H. Risk of venous thromboembolism with cyproterone or levonorgestrel contraceptives. *Lancet*. 2001;358(9291):1427-9.
- van Staa TP, Wegman S, de Vries F, Leufkens B, Cooper C. Use of statins and risk of fractures. *JAMA*. 2001;285(14):1850-5.
- van Staa TP, Leufkens HG, Cooper C. Use of inhaled corticosteroids and risk of fractures. *J Bone Miner Res*. 2001;16(3):581-8.
- van Staa TP, Dennison EM, Leufkens HG, Cooper C. Epidemiology of fractures in England and Wales. *Bone*. 2001;29(6):517-22.

- van Staa TP, Abenhaim L, Cooper C, Zhang B, Leufkens HG. Public health impact of adverse bone effects of oral corticosteroids. *British Journal of Clinical Pharmacology*. 2001;51(6):601-7.
- van Bommel MJ, Numans ME, de Wit NJ, Stalman WA. Consultations and referrals for dyspepsia in general practice--a one year database survey. *Postgrad Med J*. 2001;77(910):514-8.
- Smeeth L, Hall AJ, Fombonne E, Rodrigues LC, Huang X, Smith PG. A case-control study of autism and mumps-measles-rubella vaccination using the general practice research database: design and methodology. *BMC Public Health*. 2001;1:2.
- Seshadri S, Zornberg GL, Derby LE, Myers MW, Jick H, Drachman DA. Postmenopausal estrogen replacement therapy and the risk of Alzheimer disease. *Arch Neurol*. 2001;58(3):435-40.
- Seaman HE, Lawrenson RA, Williams TJ, MacRae KD, Farmer RD. The risk of liver damage associated with minocycline: a comparative study. *J Clin Pharmacol*. 2001;41(8):852-60.
- Schlienger RG, Haefeli WE, Jick H, Meier CR. Risk of cataract in patients treated with statins. *Arch Intern Med*. 2001;161(16):2021-6.
- Ryan R, Majeed A. Prevalence of ischaemic heart disease and its management with statins and aspirin in general practice in England and Wales, 1994 and 1998. *12 Winter 2001*. 2001;1329:34.
- Ruigómez A, Johansson S, Wallander MA, García Rodríguez LA. Gender and drug treatment as determinants of mortality in a cohort of heart failure patients. *Eur J Epidemiol*. 2001;17(4):329-35.
- Rowlands S, Devalia H, Smith C, Hubbard R, Dean A. Otitis externa in UK general practice: a survey using the UK General Practice Research Database. *Br J Gen Pract*. 2001;51(468):533-8.
- Rowlands S, Devalia H, Lawrenson R. Use of the combined oral contraceptive pill by under 16s. *J Fam Plann Reprod Health Care*. 2001;27(1):17-9.
- Rowlands S, Devalia H, Lawrenson R. Preferential prescribing of type of combined oral contraceptive pill by general practitioners to teenagers with acne. *Eur J Contracept Reprod Health Care*. 2001;6(1):9-13.
- Moser K. Inequalities in treated heart disease and mental illness in England and Wales, 1994-1998. *Br J Gen Pract*. 2001;51(467):438-44.
- Meier CR, Schlienger RG, Jick H. Use of selective serotonin reuptake inhibitors and risk of developing first-time acute myocardial infarction. *British Journal of Clinical Pharmacology*. 2001;52(2):179-84.
- McKeever TM, Lewis SA, Smith C, Collins J, Heatlie H, Frischer M, et al. Siblings, multiple births, and the incidence of allergic disease: a birth cohort study using the West Midlands general practice research database. *Thorax*. 2001;56(10):758-62.
- McGuire A, Irwin DE, Fenn P, Gray A, Anderson P, Lovering A, et al. The excess cost of acute exacerbations of chronic bronchitis in patients aged 45 and older in England and Wales. *Value Health*. 2001;4(5):370-5.
- Majeed A, Moser K, Carroll K. Trends in the prevalence and management of atrial fibrillation in general practice in England and Wales, 1994-1998: analysis of data from the general practice research database. *Heart*. 2001;86(3):284-8.
- Logie JW, Clifford GM, Farmer RD, Meesen BP. Lower urinary tract symptoms suggestive of benign prostatic obstruction--Triumph: the role of general practice databases. *European Urology*. 2001;39 Suppl 3:42-7.

- Lewis JD, Bilker WB, Brensinger C, Deren JJ, Vaughn DJ, Strom BL. Inflammatory bowel disease is not associated with an increased risk of lymphoma. *Gastroenterology*. 2001;121(5):1080-7.
- Lawrenson RA, Logie JW. Antibiotic failure in the treatment of urinary tract infections in young women. *J Antimicrob Chemother*. 2001;48(6):895-901.
- Lawrenson R, Wyndaele JJ, Vlachonikolis I, Farmer C, Glickman S. Renal failure in patients with neurogenic lower urinary tract dysfunction. *Neuroepidemiology*. 2001;20(2):138-43.
- Kaye JA, Myers MW, Jick H. Acetaminophen and the risk of renal and bladder cancer in the general practice research database. *Epidemiology*. 2001;12(6):690-4.
- Kaye JA, del Mar Melero-Montes M, Jick H. Mumps, measles, and rubella vaccine and the incidence of autism recorded by general practitioners: a time trend analysis. *BMJ*. 2001;322(7284):460-3.
- Johansson S, Wallander MA, Ruigómez A, García Rodríguez LA. Incidence of newly diagnosed heart failure in UK general practice. *Eur J Heart Fail*. 2001;3(2):225-31.
- Jick SS, Vasilakis-Scaramozza C, Maier WC. The risk of cataract among users of inhaled steroids. *Epidemiology*. 2001;12(2):229-34.
- Jick H, Vasilakis-Scaramozza C, Jick SS. Live attenuated polio vaccine and the risk of intussusception. *British Journal of Clinical Pharmacology*. 2001;52(4):451-3.
- Huerta C, Rodríguez LA. Incidence of ocular melanoma in the general population and in glaucoma patients. *J Epidemiol Community Health*. 2001;55(5):338-9.
- Huerta C, García Rodríguez LA, Möller CS, Arellano FM. The risk of obstructive airways disease in a glaucoma population. *Pharmacoepidemiol Drug Saf*. 2001;10(2):157-63.
- Hodgson C, Ellis C. Time trends in GP outpatient referrals. *Health Statistics Quartely*. 2001;10:14-9.
- Higuchi LM, Joffe S, Neufeld EJ, Weisdorf S, Rosh J, Murch S, et al. Inflammatory bowel disease associated with immune thrombocytopenic purpura in children. *J Pediatr Gastroenterol Nutr*. 2001;33(5):582-7.
- Hernández-Díaz S, Rodríguez LA. Steroids and risk of upper gastrointestinal complications. *Am J Epidemiol*. 2001;153(11):1089-93.
- Hernández-Díaz S, García-Rodríguez LA. Epidemiologic assessment of the safety of conventional nonsteroidal anti-inflammatory drugs. *Am J Med*. 2001;110 Suppl 3A:20S-7S.
- Hajat S, Haines A, Atkinson RW, Bremner SA, Anderson HR, Emberlin J. Association between air pollution and daily consultations with general practitioners for allergic rhinitis in London, United Kingdom. *Am J Epidemiol*. 2001;153(7):704-14.
- Gibbs RG, Newson R, Lawrenson R, Greenhalgh RM, Davies AH. Diagnosis and initial management of stroke and transient ischemic attack across UK health regions from 1992 to 1996: experience of a national primary care database. *Stroke*. 2001;32(5):1085-90.
- García-Rodríguez LA, Huerta-Alvarez C. Reduced risk of colorectal cancer among long-term users of aspirin and nonaspirin nonsteroidal antiinflammatory drugs. *Epidemiology*. 2001;12(1):88-93.
- García Rodríguez LA, Hernández-Díaz S. Relative risk of upper gastrointestinal complications among users of acetaminophen and nonsteroidal anti-inflammatory drugs. *Epidemiology*. 2001;12(5):570-6.

- Garcia Rodríguez LA, Hernández-Díaz S. The risk of upper gastrointestinal complications associated with nonsteroidal anti-inflammatory drugs, glucocorticoids, acetaminophen, and combinations of these agents. *Arthritis Res.* 2001;3(2):98-101.
- Gale CR, Edington J, Coles SJ, Martyn CN. Patterns of prescribing of nutritional supplements in the United Kingdom. *Clin Nutr.* 2001;20(4):333-7.
- Gaist D, Rodríguez LA, Huerta C, Hallas J, Sindrup SH. Lipid-lowering drugs and risk of myopathy: a population-based follow-up study. *Epidemiology.* 2001;12(5):565-9.
- Frischer M, Heatlie H, Norwood J, Bashford J, Millson D, Chapman S. Trends in antibiotic prescribing and associated indications in primary care from 1993 to 1997. *J Public Health Med.* 2001;23(1):69-73.
- Ellis C, Gnani S, Majeed A. Prevalence and management of heart failure in general practice in England and Wales, 1994-98. *Health Stat Q.* 2001(11):17-24.
- de Abajo FJ, García Rodríguez LA. Risk of upper gastrointestinal bleeding and perforation associated with low-dose aspirin as plain and enteric-coated formulations. *BMC Clin Pharmacol.* 2001;1:1.
- Chapple CR. Lower urinary tract symptoms suggestive of benign prostatic obstruction--Triumph: design and implementation. *Eur Urol.* 2001;39 Suppl 3:31-6.
- Andrews N, Miller E, Waight P, Farrington P, Crowcroft N, Stowe J, et al. Does oral polio vaccine cause intussusception in infants? Evidence from a sequence of three self-controlled cases series studies in the United Kingdom. *Eur J Epidemiol.* 2001;17(8):701-6.
- Akram MFG. Prevalence of comorbid mental illness and drug use recorded in general practice: preliminary findings from the general practice research database. *Drugs: education, prevention and policy.* 2001;8(3):275-80.
- Zornberg GL, Jick H. Antipsychotic drug use and risk of first-time idiopathic venous thromboembolism: a case-control study. *Lancet.* 2000;356(9237):1219-23.
- Varas-Lorenzo C, García-Rodríguez LA, Perez-Gutthann S, Duque-Oliart A. Hormone replacement therapy and incidence of acute myocardial infarction. A population-based nested case-control study. *Circulation.* 2000;101(22):2572-8.
- van Staa TP, Leufkens HG, Cooper C. Use of nonsteroidal anti-inflammatory drugs and risk of fractures. *Bone.* 2000;27(4):563-8.
- Van Staa TP, Leufkens HG, Abenhaim L, Zhang B, Cooper C. Use of oral corticosteroids and risk of fractures. *J Bone Miner Res.* 2000;15(6):993-1000.
- van Staa TP, Leufkens HG, Abenhaim L, Zhang B, Cooper C. Oral corticosteroids and fracture risk: relationship to daily and cumulative doses. *Rheumatology (Oxford).* 2000;39(12):1383-9.
- van Staa TP, Leufkens HG, Abenhaim L, Begaud B, Zhang B, Cooper C. Use of oral corticosteroids in the United Kingdom. *QJM.* 2000;93(2):105-11.
- Van Staa TP, Abenhaim L, Cooper C, Zhang B, Leufkens HG. The use of a large pharmacoepidemiological database to study exposure to oral corticosteroids and risk of fractures: validation of study population and results. *Pharmacoepidemiol Drug Saf.* 2000;9(5):359-66.

- Tooley PJ, Cookson RF, Jones M, Nuyts GD. Medical events and resource utilisation in cancer-pain patients treated with strong opioids: an analysis of the UK General Practice Research Database. *Journal of Drug Assessment*. 2000;3(3):221-33.
- Soriano JB, Maier WC, Egger P, Visick G, Thakrar B, Sykes J, et al. Recent trends in physician diagnosed COPD in women and men in the UK. *Thorax*. 2000;55(9):789-94.
- Ruigómez A, García Rodríguez LA, Hasselgren G, Johansson S, Wallander MA. Overall mortality among patients surviving an episode of peptic ulcer bleeding. *J Epidemiol Community Health*. 2000;54(2):130-3.
- Ruigómez A, García Rodríguez LA, Dev VJ, Arellano F, Raniwala J. Are schizophrenia or antipsychotic drugs a risk factor for cataracts? *Epidemiology*. 2000;11(6):620-3.
- Rowlands S, Devalia H, Lawrenson R, Logie J, Ineichen B. Repeated use of hormonal emergency contraception by younger women in the UK. *Br J Fam Plann*. 2000;26(3):138-43.
- Nightingale AL, Lawrenson RA, Simpson EL, Williams TJ, MacRae KD, Farmer RD. The effects of age, body mass index, smoking and general health on the risk of venous thromboembolism in users of combined oral contraceptives. *Eur J Contracept Reprod Health Care*. 2000;5(4):265-74.
- Myers MW, Vasilakis C, Kaufman MR, Jick H. Antihypertensive drugs and the risk of idiopathic aplastic anaemia. *British Journal of Clinical Pharmacology*. 2000;49(6):604-8.
- Millson D, Frischer M, Croft P, Goadsby PJ. Are triptans with enhanced lipophilicity used for the acute treatment of migraine associated with an increased consulting rate for depressive illness? *Cephalgia: An International Journal of Headache*. 2000;20(8):732-7.
- Meier CR, Schlienger RG, Kraenzlin ME, Schlegel B, Jick H. HMG-CoA reductase inhibitors and the risk of fractures. *JAMA*. 2000;283(24):3205-10.
- Meier CR, Napalkov PN, Wegmüller Y, Jefferson T, Jick H. Population-based study on incidence, risk factors, clinical complications and drug utilisation associated with influenza in the United Kingdom. *Eur J Clin Microbiol Infect Dis*. 2000;19(11):834-42.
- Meier CR, Derby LE, Jick SS, Jick H. Angiotensin-converting enzyme inhibitors, calcium channel blockers, and breast cancer. *Arch Intern Med*. 2000;160(3):349-53.
- Meier CR. Antibiotics in the prevention and treatment of coronary heart disease. *J Infect Dis*. 2000;181 Suppl 3:S558-62.
- Marrie RA, Wolfson C, Sturkenboom MC, Gout O, Heinzlef O, Roullet E, et al. Multiple sclerosis and antecedent infections: a case-control study. *Neurology*. 2000;54(12):2307-10.
- Majeed A, Moser K, Maxwell R. Age, sex and practice variations in the use of statins in general practice in England and Wales. *J Public Health Med*. 2000;22(3):275-9.
- Lawrenson R, Wyndaele JJ, Vlachonikolis I, Farmer C, Glickman S. A UK general practice database study of prevalence and mortality of people with neural tube defects. *Clin Rehabil*. 2000;14(6):627-30.
- Lawrenson R, Todd JC, Leydon GM, Williams TJ, Farmer RD. Validation of the diagnosis of venous thromboembolism in general practice database studies. *British Journal of Clinical Pharmacology*. 2000;49(6):591-6.

- Lawrenson R, Farmer R. Venous thromboembolism and combined oral contraceptives: does the type of progestogen make a difference? *Contraception*. 2000;62(2 Suppl):21S-8S; discussion 37S-8S.
- Langman MJ, Cheng KK, Gilman EA, Lancashire RJ. Effect of anti-inflammatory drugs on overall risk of common cancer: case-control study in general practice research database. *BMJ*. 2000;320(7250):1642-6.
- Lanes SF, Rodríguez LA, Hwang E. Baseline risk of gastrointestinal disorders among new users of meloxicam, ibuprofen, diclofenac, naproxen and indomethacin. *Pharmacoepidemiol Drug Saf*. 2000;9(2):113-7.
- Kaye JA, Derby LE, del Mar Melero-Montes M, Quinn M, Jick H. The incidence of breast cancer in the General Practice Research Database compared with national cancer registration data. *Br J Cancer*. 2000;83(11):1556-8.
- Jick SS, Kremers HM, Vasilakis-Scaramozza C. Isotretinoin use and risk of depression, psychotic symptoms, suicide, and attempted suicide. *Arch Dermatol*. 2000;136(10):1231-6.
- Jick SS. The risk of gastrointestinal bleed, myocardial infarction, and newly diagnosed hypertension in users of meloxicam, diclofenac, naproxen, and piroxicam. *Pharmacotherapy*. 2000;20(7):741-4.
- Jick H, Zornberg GL, Jick SS, Seshadri S, Drachman DA. Statins and the risk of dementia. *Lancet*. 2000;356(9242):1627-31.
- Jick H, Kaye JA, Vasilakis-Scaramozza C, Jick SS. Risk of venous thromboembolism among users of third generation oral contraceptives compared with users of oral contraceptives with levonorgestrel before and after 1995: cohort and case-control analysis. *BMJ*. 2000;321(7270):1190-5.
- Jick H. Incidence of venous thromboembolism in users of combined oral contraceptives. Methods for identifying cases and estimating person time at risk must be detailed. *BMJ*. 2000;320(7226):57;-author reply - 8.
- Ineichen B, Logie J, Rowlands S, Lawrenson R. Patterns of prescription of PC4 by general practitioners in England and Wales. *Eur J Contracept Reprod Health Care*. 2000;5(4):241-7.
- Hubbard R, Venn A, Lewis S, Britton J. Lung cancer and cryptogenic fibrosing alveolitis. A population-based cohort study. *Am J Respir Crit Care Med*. 2000;161(1):5-8.
- Hubbard R, Venn A, Britton J. Exposure to antidepressants and the risk of cryptogenic fibrosing alveolitis: a case-control study. *Eur Respir J*. 2000;16(3):409-13.
- Hubbard R, Venn A. Adult height and cryptogenic fibrosing alveolitis: a case-control study using the UK general practice research database. *Thorax*. 2000;55(10):864-6.
- Gupta D, Hansell A, Nichols T, Duong T, Ayres JG, Strachan D. Epidemiology of pneumothorax in England. *Thorax*. 2000;55(8):666-71.
- Gasse C, Derby LE, Vasilakis C, Jick H. Risk of suicide among users of calcium channel blockers: population based, nested case-control study. *BMJ*. 2000;320(7244):1251.
- Gasse C, Derby L, Vasilakis-Scaramozza C, Jick H. Incidence of first-time idiopathic seizures in users of tramadol. *Pharmacotherapy*. 2000;20(6):629-34.
- García Rodríguez LA, Varas C, Patrono C. Differential effects of aspirin and non-aspirin nonsteroidal antiinflammatory drugs in the primary prevention of myocardial infarction in postmenopausal women.

Epidemiology. 2000;11(4):382-7.

- García Rodríguez LA, Ruigómez A, Wallander MA, Johansson S, Olbe L. Detection of colorectal tumor and inflammatory bowel disease during follow-up of patients with initial diagnosis of irritable bowel syndrome. *Scand J Gastroenterol.* 2000;35(3):306-11.
- García Rodríguez LA, Huerta-Alvarez C. Reduced incidence of colorectal adenoma among long-term users of nonsteroidal antiinflammatory drugs: a pooled analysis of published studies and a new population-based study. *Epidemiology.* 2000;11(4):376-81.
- Frischer M, Norwood J, Heatlie H, Millson D, Lowdell J, Hickman M, et al. A comparison of trends in problematic drug misuse from two reporting systems. *J Public Health Med.* 2000;22(3):362-7.
- Frischer M, Heatlie H, Chapman S, Millson D. Defining the spectrum of appropriate prescribing: implications for effective medicines management. *International journal of pharmaceutical medicine.* 2000;14(1):17-21.
- Frischer M, Heatlie H, Chapman S, Bashford J, Norwood J, Millson D. Ratio of inhaled corticosteroid to bronchodilator treatment and asthma hospitalisation. *Current Medical Research and Opinion.* 2000;16(1):8-13.
- Farmer RD, Williams TJ, Simpson EL, Nightingale AL. Effect of 1995 pill scare on rates of venous thromboembolism among women taking combined oral contraceptives: analysis of general practice research database. *BMJ.* 2000;321(7259):477-9.
- Farmer RD, Lawrenson RA, Todd JC, Williams TJ, MacRae KD, Tyrer F, et al. A comparison of the risks of venous thromboembolic disease in association with different combined oral contraceptives. *British Journal of Clinical Pharmacology.* 2000;49(6):580-90.
- Eland IA, Alvarez CH, Stricker BH, Rodríguez LA. The risk of acute pancreatitis associated with acid-suppressing drugs. *British Journal of Clinical Pharmacology.* 2000;49(5):473-8.
- Derby L, Maier WC. Risk of cataract among users of intranasal corticosteroids. *J Allergy Clin Immunol.* 2000;105(5):912-6.
- del Mar Melero-Montes M, Jick H. Hyperemesis Gravidarum and the Sex of the Offspring. *Epidemiology.* 2000;12(1):123-4.
- de Abajo FJ, Jick H, Derby L, Jick S, Schmitz S. Intracranial haemorrhage and use of selective serotonin reuptake inhibitors. *British Journal of Clinical Pharmacology.* 2000;50(1):43-7.
- Clifford GM, Logie J, Farmer RD. How do symptoms indicative of BPH progress in real life practice? The UK experience. *European Urology.* 2000;38 Suppl 1:48-53.
- Vasilakis C, Jick SS, Jick H. The risk of venous thromboembolism in users of postcoital contraceptive pills. *Contraception.* 1999;59(2):79-83.
- Vasilakis C, Jick H, del Mar Melero-Montes M. Risk of idiopathic venous thromboembolism in users of progestagens alone. *Lancet.* 1999;354(9190):1610-1.
- Sturkenboom MC, Meier CR, Jick H, Stricker BH. Minocycline and lupuslike syndrome in acne patients. *Arch Intern Med.* 1999;159(5):493-7.
- Ruigómez A, Wallander MA, Johansson S, García Rodríguez LA. One-year follow-up of newly diagnosed

irritable bowel syndrome patients. *Aliment Pharmacol Ther.* 1999;13(8):1097-102.

- Ruigómez A, García Rodríguez LA, Cattaruzzi C, Troncon MG, Agostinis L, Wallander MA, et al. Use of cimetidine, omeprazole, and ranitidine in pregnant women and pregnancy outcomes. *Am J Epidemiol.* 1999;150(5):476-81.
- Rodríguez LA, Ruigómez A. Increased risk of irritable bowel syndrome after bacterial gastroenteritis: cohort study. *BMJ.* 1999;318(7183):565-6.
- Pérez-Gutthann S, García-Rodríguez LA, Duque-Oliart A, Varas-Lorenzo C. Low-dose diclofenac, naproxen, and ibuprofen cohort study. *Pharmacotherapy.* 1999;19(7):854-9.
- Meier CR, Derby LE, Jick SS, Vasilakis C, Jick H. Antibiotics and risk of subsequent first-time acute myocardial infarction. *JAMA.* 1999;281(5):427-31.
- Majeed A, Moser K. Age- and sex-specific antibiotic prescribing patterns in general practice in England and Wales in 1996. *Br J Gen Pract.* 1999;49(446):735-6.
- Lawrence M, Jones L, Lancaster T, Daly E, Banks E. Hormone replacement therapy: patterns of use studied through British general practice computerized records. *Fam Pract.* 1999;16(4):335-42.
- Jick SS, Stender M, Myers MW. Frequency of liver disease in type 2 diabetic patients treated with oral antidiabetic agents. *Diabetes Care.* 1999;22(12):2067-71.
- Jick SS, Myers MW, Jick H. Risk of idiopathic cerebral haemorrhage in women on oral contraceptives with differing progestagen components. *Lancet.* 1999;354(9175):302-3.
- Jick SS. Pregnancy outcomes after maternal exposure to fluconazole. *Pharmacotherapy.* 1999;19(2):221-2.
- Hansell A, Hollowell J, Nichols T, McNiece R, Strachan D. Use of the General Practice Research Database (GPRD) for respiratory epidemiology: a comparison with the 4th Morbidity Survey in General Practice (MSGP4). *Thorax.* 1999;54(5):413-9.
- Hajat S, Haines A, Goubet SA, Atkinson RW, Anderson HR. Association of air pollution with daily GP consultations for asthma and other lower respiratory conditions in London. *Thorax.* 1999;54(7):597-605.
- García Rodríguez LA, Ruigómez A. Secondary prevention of upper gastrointestinal bleeding associated with maintenance acid-suppressing treatment in patients with peptic ulcer bleed. *Epidemiology.* 1999;10(3):228-32.
- García Rodríguez LA, Duque A, Castellsague J, Pérez-Gutthann S, Stricker BH. A cohort study on the risk of acute liver injury among users of ketoconazole and other antifungal drugs. *British Journal of Clinical Pharmacology.* 1999;48(6):847-52.
- Frischer M, Heatlie H, Chapman S, Norwood J, Bashford J, Millson D. Should the corticosteroid to bronchodilator ratio be promoted as a quality prescribing marker? *Public Health.* 1999;113(5):247-50.
- Frischer M, Heatlie H, Chapman S, Bashford J, Norwood J. Switching between metered dose inhalers (MDIs) and dry powder inhalers (DPIs) in airways disease: An analysis of agespecific rates using the general practice research database. *Journal of Applied Therapeutic Research.* 1999;2:253-9.
- Farmer RD, Lawrenson RA, Todd JC, Williams TJ, MacRae K. Oral contraceptives and venous thromboembolic disease. Analyses of the UK General Practice Research Database and the UK Mediplus database. *Hum Reprod*

Update. 1999;5(6):688-706.

- Edington J, Winter PD, Coles SJ, Gale CR, Martyn CN. Outcomes of undernutrition in patients in the community with cancer or cardiovascular disease. *Proc Nutr Soc.* 1999;58(3):655-61.
- Dolan P, Torgerson DJ, Wolstenholme J. Costs of breast cancer treatment in the United Kingdom. *Breast.* 1999;8(4):205-7.
- Derby LE, Myers MW, Jick H. Use of dexfenfluramine, fenfluramine and phentermine and the risk of stroke. *British Journal of Clinical Pharmacology.* 1999;47(5):565-9.
- de Abajo FJ, Rodríguez LA, Montero D. Association between selective serotonin reuptake inhibitors and upper gastrointestinal bleeding: population based case-control study. *BMJ.* 1999;319(7217):1106-9.
- de Abajo FJ, Rodríguez LA. Risk of ventricular arrhythmias associated with nonsedating antihistamine drugs. *British Journal of Clinical Pharmacology.* 1999;47(3):307-13.
- Wallace H, Shorvon S, Tallis R. Age-specific incidence and prevalence rates of treated epilepsy in an unselected population of 2,052,922 and age-specific fertility rates of women with epilepsy. *Lancet.* 1998;352(9145):1970-3.
- van Staa TP, Leufkens H, Abenham L, Cooper C. Postmarketing surveillance of the safety of cyclic etidronate. *Pharmacotherapy.* 1998;18(5):1121-8.
- van Staa TP, Abenham L, Cooper C. Use of cyclical etidronate and prevention of non-vertebral fractures. *Br J Rheumatol.* 1998;37(1):87-94.
- Torgerson DJ, Dolan P. Prescribing by general practitioners after an osteoporotic fracture. *Annals of the Rheumatic Diseases.* 1998;57(6):378-9.
- Thorogood M, Armstrong B, Nichols T, Hollowell J. Mortality in people taking selegiline: observational study. *BMJ.* 1998;317(7153):252-4.
- Ruigómez A, García Rodríguez LA. Presence of diabetes related complication at the time of NIDDM diagnosis: an important prognostic factor. *Eur J Epidemiol.* 1998;14(5):439-45.
- Myers MW, Jick H. Terfenadine and risk of acute liver disease. *British Journal of Clinical Pharmacology.* 1998;46(3):251-3.
- Meier CR, Sturkenboom MC, Cohen AS, Jick H. Postmenopausal estrogen replacement therapy and the risk of developing systemic lupus erythematosus or discoid lupus. *J Rheumatol.* 1998;25(8):1515-9.
- Meier CR, Jick SS, Derby LE, Vasilakis C, Jick H. Acute respiratory-tract infections and risk of first-time acute myocardial infarction. *Lancet.* 1998;351(9114):1467-71.
- Meier CR, Jick H. Tamoxifen and risk of idiopathic venous thromboembolism. *British Journal of Clinical Pharmacology.* 1998;45(6):608-12.
- Martyn CN, Winter PD, Coles SJ, Edington J. Effect of nutritional status on use of health care resources by patients with chronic disease living in the community. *Clin Nutr.* 1998;17(3):119-23.
- Lancaster T, Swart AM, Jick H. Risk of serious haematological toxicity with use of chloramphenicol eye drops in a British general practice database. *BMJ.* 1998;316(7132):667.

- Jick SS, Vasilakis C, Martinez C, Jick H. A study of the relation of exposure to quinolones and suicidal behaviour. *British Journal of Clinical Pharmacology*. 1998;45(1):77-81.
- Jick SS, Vasilakis C, Jick H. Pregnancies and terminations after 1995 warning about third-generation oral contraceptives. *Lancet*. 1998;351(9113):1404-5.
- Jick H, Vasilakis C, Weinrauch LA, Meier CR, Jick SS, Derby LE. A population-based study of appetite-suppressant drugs and the risk of cardiac-valve regurgitation. *N Engl J Med*. 1998;339(11):719-24.
- Jick H, Derby LE, Vasilakis C, Fife D. The risk of seizures associated with tramadol. *Pharmacotherapy*. 1998;18(3):607-11.
- Hubbard R, Venn A, Smith C, Cooper M, Johnston I, Britton J. Exposure to commonly prescribed drugs and the etiology of cryptogenic fibrosing alveolitis: a case-control study. *Am J Respir Crit Care Med*. 1998;157(3 Pt 1):743-7.
- Hubbard R, Venn A, Lewis S, Britton J. Antidepressant exposure and risk of cryptogenic fibrosing alveolitis (CFA): a case-control study using the General Practice Research Database (GPRD). *THORAX-LONDON*. 1998;53:S44-S.
- Haste F, Charlton J, Jenkins R. Potential for suicide prevention in primary care? An analysis of factors associated with suicide. *Br J Gen Pract*. 1998;48(436):1759-63.
- Gibbs RG, Todd JC, Irvine C, Lawrenson R, Newson R, Greenhalgh RM, et al. Relationship between the regional and national incidence of transient ischaemic attack and stroke and performance of carotid endarterectomy. *Eur J Vasc Endovasc Surg*. 1998;16(1):47-52.
- García Rodríguez LA, Ruigómez A, Hasselgren G, Wallander MA, Johansson S. Comparison of mortality from peptic ulcer bleed between patients with or without peptic ulcer antecedents. *Epidemiology*. 1998;9(4):452-6.
- García Rodríguez LA, Pérez Gutthann S. Use of the UK General Practice Research Database for pharmacoepidemiology. *British Journal of Clinical Pharmacology*. 1998;45(5):419-25.
- García Rodríguez LA, Cattaruzzi C, Troncon MG, Agostinis L. Risk of hospitalization for upper gastrointestinal tract bleeding associated with ketorolac, other nonsteroidal anti-inflammatory drugs, calcium antagonists, and other antihypertensive drugs. *Arch Intern Med*. 1998;158(1):33-9.
- Dolan P, Torgerson DJ. The cost of treating osteoporotic fractures in the United Kingdom female population. *Osteoporos Int*. 1998;8(6):611-7.
- Derby LE, Jick H. Appendectomy protects against ulcerative colitis. *Epidemiology*. 1998;9(2):205-7.
- De Lusignan S, Brown A, Majeed FA. Recent developments in general practice computing. *Clinician in Management*. 1998;7:148-54.
- Castellsague J, Pérez Gutthann S, García Rodríguez LA. Recent epidemiological studies of the association between hormone replacement therapy and venous thromboembolism. A review. *Drug Safety*. 1998;18(2):117-23.
- Blackburn SC, Oliart AD, García Rodríguez LA, Pérez Gutthann S. Antiepileptics and blood dyscrasias: a cohort study. *Pharmacotherapy*. 1998;18(6):1277-83.
- Bashford JN, Norwood J, Chapman SR. Why are patients prescribed proton pump inhibitors? Retrospective

analysis of link between morbidity and prescribing in the General Practice Research Database. *BMJ*. 1998;317(7156):452-6.

- Walker AM, Szneke P, Bianchi LA, Field LG, Sutherland LR, Dreyer NA. 5-Aminosalicylates, sulfasalazine, steroid use, and complications in patients with ulcerative colitis. *Am J Gastroenterol*. 1997;92(5):816-20.
- van Staa T, Abenhaim L, Monette J. Rates of hypoglycemia in users of sulfonylureas. *J Clin Epidemiol*. 1997;50(6):735-41.
- van Staa T, Abenhaim L, Cooper C. Upper gastrointestinal adverse events and cyclical etidronate. *Am J Med*. 1997;103(6):462-7.
- Pérez Gutthann S, García Rodríguez LA, Castellsague J, Duque Oliart A. Hormone replacement therapy and risk of venous thromboembolism: population based case-control study. *BMJ*. 1997;314(7083):796-800.
- Meier CR, Jick H. Drug use and pulmonary death rates in increasingly symptomatic asthma patients in the UK. *Thorax*. 1997;52(7):612-7.
- Meier CR, Jick H. Omeprazole, H2 blockers, and polyarthralgia: case-control study. *BMJ*. 1997;315(7118):1283.
- Meier CR, Jick H. Omeprazole, other antiulcer drugs and newly diagnosed gout. *British Journal of Clinical Pharmacology*. 1997;44(2):175-8.
- Jick SS, Terris BZ. Anticonvulsants and congenital malformations. *Pharmacotherapy*. 1997;17(3):561-4.
- Jick S. Ciprofloxacin safety in a pediatric population. *The Pediatric infectious disease journal*. 1997;16(1):130-4.
- Jick H, Vasilakis C, Derby LE. Antihypertensive drugs and fatal myocardial infarction in persons with uncomplicated hypertension. *Epidemiology*. 1997;8(4):446-8.
- Jick H. Calcium-channel blockers and risk of cancer. *Lancet*. 1997;349(9066):1699-700.
- Hollowell J. The General Practice Research Database: quality of morbidity data. *Popul Trends*. 1997(87):36-40.
- Hajat S, Goubet SA, Haines A. Thunderstorm-associated asthma: the effect on GP consultations. *Br J Gen Pract*. 1997;47(423):639-41.
- García Rodríguez LA, Wallander MA, Stricker BH. The risk of acute liver injury associated with cimetidine and other acid-suppressing anti-ulcer drugs. *British Journal of Clinical Pharmacology*. 1997;43(2):183-8.
- García Rodríguez LA, Wallander MA, Johanson S, Björck S. Renal disease and acid-suppressing drugs. *Pharmacoepidemiol Drug Saf*. 1997;6(4):247-51.
- García Rodríguez LA, Ruigómez A, Jick H. A review of epidemiologic research on drug-induced acute liver injury using the general practice research data base in the United Kingdom. *Pharmacotherapy*. 1997;17(4):721-8.
- García Rodríguez LA, Ruigómez A. Gastric acid, acid-suppressing drugs, and bacterial gastroenteritis: how much of a risk? *Epidemiology*. 1997;8(5):571-4.
- Turnbull S, Ward A, Treasure J, Jick H, Derby L. The demand for eating disorder care. An epidemiological study using the general practice research database. *Br J Psychiatry*. 1996;169(6):705-12.
- Jick H, Derby LE, Gurewich V, Vasilakis C. The risk of myocardial infarction associated with antihypertensive

drug treatment in persons with uncomplicated essential hypertension. *Pharmacotherapy*. 1996;16(3):321-6.

- Henry D, Lim LL, Garcia Rodriguez LA, Perez Gutthann S, Carson JL, Griffin M, et al. Variability in risk of gastrointestinal complications with individual non-steroidal anti-inflammatory drugs: results of a collaborative meta-analysis. *BMJ*. 1996;312(7046):1563-6.
- García Rodríguez LA, Stricker BH, Zimmerman HJ. Risk of acute liver injury associated with the combination of amoxicillin and clavulanic acid. *Arch Intern Med*. 1996;156(12):1327-32.
- García Rodríguez LA, Mannino S, Wallander MA, Lindblom B. A cohort study of the ocular safety of anti-ulcer drugs. *British Journal of Clinical Pharmacology*. 1996;42(2):213-6.
- Donovan S. The use of the General Practice Research Database (GPRD) to examine potential links between antidepressant medication and the incidence of suicide. *Drug information journal*. 1996;30(3):785-92.
- Derby LE, Tennis P, Jick H. Sudden unexplained death among subjects with refractory epilepsy. *Epilepsia*. 1996;37(10):931-5.
- Nazareth I, King M, Davies S. Care of schizophrenia in general practice: the general practitioner and the patient. *Br J Gen Pract*. 1995;45(396):343-7.
- Lloyd DC, Harris CM, Roberts DJ. Specific therapeutic group age-sex related prescribing units (STAR-PU): weightings for analysing general practices' prescribing in England. *BMJ*. 1995;311(7011):991-4.
- Lancaster T, Surman G, Lawrence M, Mant D, Vessey M, Thorogood M, et al. Hormone replacement therapy: characteristics of users and non-users in a British general practice cohort identified through computerised prescribing records. *J Epidemiol Community Health*. 1995;49(4):389-94.
- Jick SS, Dean AD, Jick H. Antidepressants and suicide. *BMJ*. 1995;310(6974):215-8.
- Jick H, Myers MW, Dean AD. The risk of sulfasalazine- and mesalazine-associated blood disorders. *Pharmacotherapy*. 1995;15(2):176-81.
- Jick H, Jick SS, Gurewich V, Myers MW, Vasilakis C. Risk of idiopathic cardiovascular death and nonfatal venous thromboembolism in women using oral contraceptives with differing progestagen components. *Lancet*. 1995;346(8990):1589-93.
- Jick H, Derby LE. A large population-based follow-up study of trimethoprim-sulfamethoxazole, trimethoprim, and cephalexin for uncommon serious drug toxicity. *Pharmacotherapy*. 1995;15(4):428-32.
- García Rodríguez LA, Mannino S, Wallander MA. Ocular safety of antiulcer drugs. *Lancet*. 1995;345(8956):1059-60.
- Van Staa TP, Abenhaim L. The quality of information recorded on a UK database of primary care records: a study of hospitalizations due to hypoglycemia and other conditions. *Pharmacoepidemiol Drug Saf*. 1994;3(1):15-21.
- Roberts SJ, Bateman DN. The use of nitrates, calcium channel blockers and ACE inhibitors in primary care in the Northern Region: a pharmacoepidemiological study. *British Journal of Clinical Pharmacology*. 1994;38(6):489-97.
- Mann RD, Lis Y, Chukwujindu J, Chanter DO. A study of the association between hormone replacement therapy, smoking and the occurrence of myocardial infarction in women. *J Clin Epidemiol*. 1994;47(3):307-12.

- Mann RD. The VAMP system and the detection of risk, some examples. *Pharmacoepidemiol Drug Saf.* 1994;3(6):313-9.
- Jick H, Derby LE, Dean AD, Henry DA. Flucloxacillin and cholestatic hepatitis. *Med J Aust.* 1994;160(8):525.
- García Rodríguez LA, Williams R, Derby LE, Dean AD, Jick H. Acute liver injury associated with nonsteroidal anti-inflammatory drugs and the role of risk factors. *Arch Intern Med.* 1994;154(3):311-6.
- García Rodríguez LA, Jick H. Risk of gynaecomastia associated with cimetidine, omeprazole, and other antiulcer drugs. *BMJ.* 1994;308(6927):503-6.
- García Rodríguez LA, Jick H. Risk of upper gastrointestinal bleeding and perforation associated with individual non-steroidal anti-inflammatory drugs. *Lancet.* 1994;343(8900):769-72.
- Fleming DM. Practice based information systems for monitoring medicines. *Pharmaceutical medicine.* 1994;8(3-4):161-76.
- Sleator DJ. Towards accurate prescribing analysis in general practice: accounting for the effects of practice demography. *Br J Gen Pract.* 1993;43(368):102-6.
- Nazareth I, King M, Haines A, Tai SS, Hall G. Care of schizophrenia in general practice. *BMJ.* 1993;307(6909):910.
- Nazareth I, King M, Haines A, Rangel L, Myers S. Accuracy of diagnosis of psychosis on general practice computer system. *BMJ.* 1993;307(6895):32-4.
- Lis Y, Spitzer WO, Mann RD, Chukwujindu J, Thorogood M, Evans S, et al. A concurrent cohort study of oral contraceptive users from the VAMP research bank. *Pharmacoepidemiol Drug Saf.* 1993;2(1):51-63.
- Jick SS, Jick H, Dean AD. A follow-up safety study of ciprofloxacin users. *Pharmacotherapy.* 1993;13(5):461-4.
- Jick H, Derby LE, García Rodríguez LA, Jick SS, Dean AD. Nonsteroidal antiinflammatory drugs and certain rare, serious adverse events: a cohort study. *Pharmacotherapy.* 1993;13(3):212-7.
- Hall GC, Chukwujindu J, Richardson J, Lis Y, Wild RN. Micturin (terodiline hydrochloride), torsades de pointe and other arrhythmias—a study using the VAMP Database. *Pharmacoepidemiol Drug Saf.* 1993;2(2):127-32.
- Derby LE, Jick H, Henry DA, Dean AD. Erythromycin-associated cholestatic hepatitis. *Med J Aust.* 1993;158(9):600-2.
- Derby LE, Jick H, Henry DA, Dean AD. Cholestatic hepatitis associated with flucloxacillin. *Med J Aust.* 1993;158(9):596-600.
- Derby LE, Gutthann SP, Jick H, Dean AD. Liver disorders in patients receiving chlorpromazine or isoniazid. *Pharmacotherapy.* 1993;13(4):353-8.
- Wild RN. Micturin and torsades de pointe—experience of a post-marketing alert. *Pharmacoepidemiol Drug Saf.* 1992;1(3):147-50.
- Orlander JD, Jick SS, Dean AD, Jick H. Urinary tract infections and estrogen use in older women. *Journal of the American Geriatrics Society.* 1992;40(8):817-20.
- Mannino S, Garcia-Rodriguez LA, Jick SS. NSAIDs, quinolones and convulsions: an epidemiologic approach. *Post Marketing Surveillance.* 1992;6(2):119-.

- Mann RD, Hall G, Chukwujindu J. Research implications of computerised primary care. *Post Marketing Surveillance*. 1992;5:259-68.
- Jick SS, Jick H, Knauss TA, Dean AD. Antidepressants and convulsions. *J Clin Psychopharmacol*. 1992;12(4):241-5.
- Jick H, Ulcickas M, Dean A. Comparison of frequencies of suicidal tendencies among patients receiving fluoxetine, lofepramine, mianserin, or trazodone. *Pharmacotherapy*. 1992;12(6):451-4.
- Jick H, Terris BZ, Derby LE, Jick SS. Further validation of information recorded on a general practitioner based computerized data resource in the United Kingdom. *Pharmacoepidemiol Drug Saf*. 1992;1(6):347-9.
- Jick H, Derby LE, García Rodríguez LA, Jick SS, Dean AD. Liver disease associated with diclofenac, naproxen, and piroxicam. *Pharmacotherapy*. 1992;12(3):207-12.
- Derby LE, Jick H, Dean AD. Antidepressant drugs and suicide. *J Clin Psychopharmacol*. 1992;12(4):235-40.
- Tallis R, Hall G, Craig I, Dean A. How common are epileptic seizures in old age? *Age Ageing*. 1991;20(6):442-8.
- Jick H, Jick SS, Derby LE. Validation of information recorded on general practitioner based computerised data resource in the United Kingdom. *BMJ*. 1991;302(6779):766-8.
- Lawrence M, Coulter A, Jones L. A total audit of preventive procedures in 45 practices caring for 430,000 patients. *BMJ*. 1990;300(6738):1501-3.
- Jick H, Hall GC, Dean AD, Jick SS, Derby LE. A comparison of the risk of hypoglycemia between users of human and animal insulins. 1. Experience in the United Kingdom. *Pharmacotherapy*. 1990;10(6):395-7.
- Hall GC, Luscombe DK, Walker SR. Post-marketing surveillance using a computerised general practice data base. *Pharmaceutical medicine*. 1988;2(4):345-51.